

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent	Classification	6	:
---------------------------	----------------	---	---

# C12N 15/31, C07K 14/35, 16/12, C12Q 1/68, C12N 15/62, G01N 33/53

(11) International Publication Number:

WO 98/16645

A2

(43) International Publication Date:

23 April 1998 (23.04.98)

(21) International Application Number:

PCT/US97/18214

(22) International Filing Date:

7 October 1997 (07.10.97)

(74) Agents: MAKI, David, J. et al., Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).

(30) Priority Data:

08/729,622 08/818,111

11 October 1996 (11.10.96) 13 March 1997 (13.03.97)

US US

(71) Applicant: CORIXA CORPORATION [US/US]; 1124 Columbia Street, Seattle, WA 98104 (US).


(72) Inventors: REED, Steven, G.; 2843 - 122nd Place N.E., Bellevue, WA 98005 (US). SKEIKY, Yasir, A., W.; 8327 - 25th Avenue N.W., Seattle, WA 98107 (US). DILLON, Davin, C.; 21607 N.E. 24th Street, Redmond, WA 98053 (US). CAMPOS-NETO, Antonio; 9308 Midship Court N.E., Bainbridge Island, WA 98021 (US). HOUGHTON, Raymond; 2636 - 242nd Place S.E., Bothell, WA 98021 (US). VEDVICK, Thomas, S., 124 South 300th Place, Federal Way, WA 98003 (US). TWARDZIK, Daniel, R.; 10195 South Beach Drive, Bambridge Island, WA 98110 (US). LODES, Michael, J.; 9223 - 36th Avenue S.W., Seattle, WA 98126 (US).

(81) Designated States: AL, AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OARI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished .:pon receipt of that report.

(54) Title: COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUBERCULOSIS



where the term of Markov and street of DNA symmetric terms of the street of the symmetric production of the street of the stree

الهابي في المحاجب المساول في الأواجب

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of panighlets publishing international applications under the PCT.

۸ī	$\Lambda^{+} \sim (4)$	ES	$S_1^{r}(\Gamma)$	1.5	1000	SI	Slovema
AM	Afternak	1:1	Empand	1.1	Lathigania	SK	Sorvania
AT	Nostria	FR	Eran.	1.1	i sensi any	88	Semegal
11	Numerican	GA	Galeon	1 V	lains ia	SZ	Swäziland
AZ	Azsabanjun	GB	United Kingdom	MC	Months o	TD	Chad
BA	Bossia and Herzek is in a	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tapkistan
BE	Belgnun	GN	Guinea	MK	The former Yugoslas	TM	burkmenistay.
BF	Birkin i Fasi	GR	Citer. r		Rapubli of Micedonia	TR	Lakes
BG	Bulgaria	HU	Hompary	311	Maria	1 1	Itimidad and 1 Sagar
BJ	Brian.	D.	Ire and	MN	Miling illia	LA	Larisme
BR	30.4.1	Ħ	15,5 401	MR	*,*: - *;		
HY	· v · · · ·	1.					

		1
. *.	Sq. et	1
* 1	ls z* · · · · ·	1, 1
- /	1.4	21

1

# COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUBERCULOSIS

# TECHNICAL FIELD

5

10

The present invention relates generally to the detection of *Mycobacterium* tuberculosis infection. The invention is more particularly related to polypeptides comprising a *Mycobacterium tuberculosis* antigen, or a portion or other variant thereof, and the use of such polypeptides for the serodiagnosis of *Mycobacterium tuberculosis* infection.

### BACKGROUND OF THE INVENTION

Tuberculosis is a chronic, infectious disease, that is generally caused by infection with *Mycobacterium tuberculosis*. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world, with about 8 million new cases and 3 million deaths each year. Although the infection may be asymptomatic for a considerable period of time, the disease is most commonly manifested as an acute inflammation of the lungs, resulting in fever and a nonproductive cough. If left untreated, serious complications and death typically result.

Although tuberculosis can generally be controlled using extended antibiotic therapy, such treatment is not sufficient to prevent the spread of the disease. Infected individuals may be asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behavior is difficult to monitor. Some patients do not complete the course of treatment, which can lead to ineffective treatment and the development of drug resistance.

Inhibiting the spread of tuberculosis will require effective vaccination and accurate, early diagnosis of the disease. Currently, vaccination with live bacteria is the most efficient method for inducing protective immunity. The most common Mycobacterium for this purpose is Bacillus Calmette Canada (Beck).

a filosoficio de la companya de la companya de la filosoficia de la PPD epretena parametra de la companya del companya de la companya de la companya del companya de la companya del companya de la companya de la companya de la companya del companya de la companya de la companya de la companya del co

15

.10

site by 48-72 hours after injection, which indicates exposure to Mycobacterial antigens. Sensitivity and specificity have, however, been a problem with this test, and individuals vaccinated with BCG cannot be distinguished from infected individuals.

While macrophages have been shown to act as the principal effectors of *M. tuberculosis* immunity. T cells are the predominant inducers of such immunity. The essential role of T cells in protection against *M. tuberculosis* infection is illustrated by the frequent occurrence of *M. tuberculosis* in AIDS patients, due to the depletion of CD4 T cells associated with human immunodeficiency virus (HIV) infection. Mycobacterium-reactive CD4 T cells have been shown to be potent producers of gamma-interferon (IFN- $\gamma$ ), which, in turn, has been shown to trigger the anti-mycobacterial effects of macrophages in mice. While the role of IFN- $\gamma$  in humans is less clear, studies have shown that 1,25-dihydroxy-vitamin D3, either alone or in combination with IFN- $\gamma$  or tumor necrosis factor-alpha, activates human macrophages to inhibit *M. tuberculosis* infection. Furthermore, it is known that IFN- $\gamma$  stimulates human macrophages to make 1,25-dihydroxy-vitamin D3. Similarly, IL-12 has been shown to play a role in stimulating resistance to *M. tuberculosis* infection. For a review of the immunology of *M. tuberculosis* infection see Chan and Kaufmann, in *Tuberculosis*: *Pathogenesis*. *Protection and Control*, Bloom (ed.), ASM Press, Washington, DC, 1994.

Accordingly, there is a need in the art for improved diagnostic methods for detecting tuberculosis. The present invention fulfills this need and further provides other related advantages.

#### SUMMARY OF THE INVENTION

Briefly stated, the present invention provides compositions and methods for diagnosing tuberculosis. In one aspect, polypeptides are provided comprising an antigenic portion of a soluble *M tuberculosis* antigen, or a variant of such an antigen that differs only in concernation of the concernation.

	(b)	Ala-Val-Glu-Scr-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser
		(SEQ ID NO: 116);
	(c)	Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-
		Lys-Glu-Gly-Arg (SEQ ID NO: 117);
5	(d)	Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro
		(SEQ ID NO: 118);
	·(e)	Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID
	* <b>*</b>	NO: 119);
	(f)	Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID
10		NO: 120);
	(g)	Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-
		Ser (SEQ ID NO: 121);
	(h)	Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly
		(SEQ ID NO: 122);
15	(i)	Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Leu-Thr-Ser-
		Leu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn-Val-Ser-Phe-Ala-Asn (SEQ
		ID NO: 123);
	(j)	Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-He-Lys-Val-Thr-Asp-Ala-Ser;
		(SEQ ID NO: 129)
20	(k)	Ala Gly Asp Thr Xaa Ile Tvr-Ee-Val-Glv-Asn-Leu-Thr-Ala-Asp:
		(SFQ ID NO. 130) or
	(1)	Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu Gly Gly Thr-Val Gln-Ala-Gly;
		(SEQ ID NO: 131)
25	where	zin Xaa may be any amino acid

- (m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132) or
- (n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124)
- 5 wherein Xaa may be any amino acid.

In another embodiment, the soluble *M. tuberculosis* antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96 or a complement thereof under moderately stringent conditions.<sup>12</sup>

In a related aspect, the polypeptides comprise an antigenic portion of a *M. tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and/or modifications, wherein the antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 26-51, 133, 134, 158-178 and 196, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 26-51, 133, 134, 158-178 and 196 or a complement thereof under moderately stringent conditions.

In related aspects, DNA sequences encoding the above polypeptides, recombinant expression vectors comprising these DNA sequences and host cells transformed or transfected with such expression vectors are also provided.

In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known M tuberculosis antigen.

In further aspects of the subject invention, methods and diagnostic kits are provided for detecting tuberculosis in a patient. The methods comprise: (a) contacting a book of the subject invention is a patient.

the state with a value category matter and urms. The diagnostic for

The present invention also provides methods for detecting *M. tuberculosis* infection comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with at least one oligonucleotide primer in a polymerase chain reaction, the oligonucleotide primer being specific for a DNA sequence encoding the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In one embodiment, the oligonucleotide primer comprises at least about 10 contiguous nucleotides of such a DNA sequence.

In a further aspect, the present invention provides a method for detecting *M. tuberculosis* infection in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a DNA sequence encoding the above polypeptides; and (c) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe. In one embodiment, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of such a DNA sequence.

In yet another aspect, the present invention provides antibodies, both polyclonal and monoclonal, that bind to the polypeptides described above, as well as methods for their use in the detection of *M. tuberculosis* infection.

15

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

# BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE IDENTIFIERS

Figure 1A and B illustrate the stimulation of proliferation and interferon γ production in 1 cells derived from a first and a second *M tuberculosis*-immune donor, respectively, by the 14 Kd, 20 Kd and 26 Kd antigens described in Example 1.

Section motors of and research infinantic BSC Countries are embinaried BUDG carries of ending

Figure 3A illustrates the stimulation of proliferation in a TbH-9-specific T cell clone by secretory *M. tuberculosis* proteins, recombinant TbH-9 and a control antigen, TbRa11.

Figure 3B illustrates the stimulation of interferon-y production in a TbH-9-specific T cell clone by secretory *M. tubercutosis* proteins, PPD and recombinant TbH-9.

Figure 4 illustrates the reactivity of two representative polypeptides with sera from *M tuberculosis*-infected and uninfected individuals, as compared to the reactivity of bacterial lysate.

Figure 5 shows the reactivity of four representative polypeptides with sera from *M. tuberculosis*-infected and uninfected individuals, as compared to the reactivity of the 38 kD antigen.

Figure 6 shows the reactivity of recombinant 38 kD and TbRa11 antigens with sera from *M. tuberculosis* patients, PPD positive donors and normal donors.

Figure 7 shows the reactivity of the antigen TbRa2A with 38 kD negative sera.

Figure 8 shows the reactivity of the antigen of SEQ ID NO: 60 with sera from *M. tuberculosis* patients and normal donors.

Figure 9 illustrates the reactivity of the recombinant antigen TbH-29 (SEQ ID NO: 137) with sera from *M. tuberculosis* patients, PPD positive donors and normal donors as determined by indirect FLISA.

20 Figure 10 illustrates the reactivity of the recombinant antigen TbH-33 (SEQ ID NO: 140) with sera from *M. tuberculosis* patients and from normal donors, and with a pool of sera from *M. tuberculosis* patients, as determined both by direct and indirect ELISA

Figure 11 illustrates the reactivity of increasing concentrations of the recombinant antigen 1bH-33 (SEQ ID NO: 140) with sera from M. tuberculosis patients and from normal donors as determined by ELISA.

	SEQ. ID NO. 5 is the DNA sequence of TbRa13.
	SEQ. ID NO. 6 is the DNA sequence of TbRa16.
	SEQ. ID NO. 7 is the DNA sequence of TbRa17.
	SEQ. ID NO. 8 is the DNA sequence of TbRa18.
5	SEQ. ID NO. 9 is the DNA sequence of TbRa19.
	SEQ. ID NO. 10 is the DNA sequence of TbRa24.
	SEQ. ID NO. 11 is the DNA sequence of TbRa26.
	SEQ. ID NO. 12 is the DNA sequence of TbRa28.
	SEQ. ID NO. 13 is the DNA sequence of TbRa29.
10	SEQ. ID NO. 14 is the DNA sequence of TbRa2A.
	SEQ. ID NO. 15 is the DNA sequence of TbRa3.
	SEQ. ID NO. 16 is the DNA sequence of TbRa32.
	SEQ. ID NO. 17 is the DNA sequence of TbRa35.
	SEQ. ID NO. 18 is the DNA sequence of TbRa36.
15	SEQ. ID NO. 19 is the DNA sequence of TbRa4.
	SEQ. ID NO. 20 is the DNA sequence of TbRa9.
	SEQ. ID NO. 21 is the DNA sequence of TbRaB.
	SEQ. ID NO. 22 is the DNA sequence of TbRaC.
	SFQ. ID NO. 23 is the DNA sequence of TbRaD
20	SEQ. ID NO. 24 is the DNA sequence of YYWCPC
	SEQ. ID NO. 25 is the DNA sequence of AAMK.
	SEQ. ID NO. 26 is the DNA sequence of TbL-23.
	SEQ. ID NO. 27 is the DNA sequence of TbL-24.
	SFQ. ID NO. 28 is the DNA sequence of Tb1 -25
25	SLQ, ID NO. 29 is the DNA sequence of Tb1. 28.

SEQ.	ID	NO.	35	is the	DNA	sequence	of TbM-3.

- SEQ. ID NO. 36 is the DNA sequence of TbM-6.
- SEQ. ID NO. 37 is the DNA sequence of TbM-7.
- SEQ. ID NO. 38 is the DNA sequence of TbM-9.
- 5 SEQ. ID NO. 39 is the DNA sequence of TbM-12.
  - SEQ. ID NO. 40 is the DNA sequence of TbM-13
  - SEQ. ID NO. 41 is the DNA sequence of TbM-14.
  - SEQ. ID NO. 42 is the DNA sequence of TbM-15.
  - SEQ. ID NO. 43 is the DNA sequence of TbH-4.
- SEQ. ID NO. 44 is the DNA sequence of TbH-4-FWD.
  - SEQ. ID NO. 45 is the DNA sequence of TbH-12.
  - SEQ. ID NO. 46 is the DNA sequence of Tb38-1.
  - SEQ. ID NO. 47 is the DNA sequence of Tb38-4.
  - SFQ. ID NO. 48 is the DNA sequence of TbL-17.
- 15 SEQ. ID NO. 49 is the DNA sequence of TbL-20.
  - SEQ. ID NO. 50 is the DNA sequence of TbL-21.
  - SEQ. ID NO. 51 is the DNA sequence of TbH-16.
  - SFQ. ID NO. 52 is the DNA sequence of DPEP.
  - SEQ. ID NO. 53 is the deduced amino acid sequence of DPFP.
- 20 SEQ. ID NO. 54 is the protein sequence of DPV N-terminal Antigen.
  - SEQ. ID NO. 55 is the protein sequence of AVGS N-terminal Antigen.
  - SFQ. ID NO. 56 is the protein sequence of AAMK N terminal Antigen.
  - SEQ. ID NO. 57 is the protein sequence of YYWC N-terminal Antigen.
  - SFQ. ID NO. 58 is the protein sequence of DIGS N-terminal Antigen.
- 25 SEQ. ID NO. 59 is the protein sequence of AFES N-terminal Antigen

Sing ID Notes in the leaffect and each range contribution in the second section of the section of the second section of the section of the

	SEQ. ID NO. 65 is the deduced amino acid sequence of TbRa10.
	SEQ. ID NO. 66 is the deduced amino acid sequence of TbRa11.
	SEQ. ID NO. 67 is the deduced amino acid sequence of TbRa12.
	SEQ. ID NO. 68 is the deduced amino acid sequence of TbRa13.
5	SEQ. ID NO. 69 is the deduced amino acid sequence of TbRa16.
	SEQ. ID NO 70 is the deduced amino acid sequence of TbRa17.
	SEQ. ID NO. 71 is the deduced amino acid sequence of TbRa18.
	SEQ. ID NO. 72 is the deduced amino acid sequence of TbRa19.
	SEQ. ID NO. 73 is the deduced amino acid sequence of TbRa24.
10	SEQ. ID NO. 74 is the deduced amino acid sequence of TbRa26.
	SFQ. ID NO. 75 is the deduced amino acid sequence of TbRa28.
	SEQ. ID NO. 76 is the deduced amino acid sequence of TbRa29.
	SEQ. ID NO. 77 is the deduced amino acid sequence of TbRa2A.
	SFQ. ID NO. 78 is the deduced amino acid sequence of TbRa3.
15	SFQ. ID NO. 79 is the deduced amino acid sequence of TbRa32.
	SFQ. ID NO. 80 is the deduced amino acid sequence of TbRa35.
	SEQ. ID NO. 81 is the deduced amino acid sequence of TbRa36.
	SEQ. ID NO. 82 is the deduced amino acid sequence of TbRa4.
	SFQ. ID NO. 83 is the deduced amino acid sequence of TbRa9.
20	SFQ, ID NO, 84 is the deduced amino acid sequence of TbRaB-
	SEQ. ID NO. 85 is the deduced amino acid sequence of TbRaC.
	SEQ. ID NO. 86 is the deduced amino acid sequence of TbRaD.
	SEQ. ID NO. 87 is the deduced amino acid sequence of YYWCPG
	SEQ. ID NO. 88 is the deduced amino acid sequence of TbAAMK.
25	SEQ. ID NO. 89 is the deduced amino acid sequence of Tb38-1

Sto Honores of a transfer for the or with

PCT/US97/18214

SEQ. ID NO. 95 is the deduced amino acid sequence of DPAS.

SEQ. ID NO. 96 is the DNA sequence of DPV.

SEQ. ID NO. 97 is the deduced amino acid sequence of DPV.

SEO. ID NO. 98 is the DNA sequence of ESAT-6.

5 SEQ. ID NO. 99 is the deduced amino acid sequence of ESAT-6.

SEQ. ID NO. 100 is the DNA sequence of TbH-8-2.

SEQ. ID NO. 101 is the DNA sequence of TbH-9FL.

SEQ. ID NO. 102 is the deduced amino acid sequence of TbH-9FL.

SEQ. ID NO. 103 is the DNA sequence of TbH-9-1.

SEQ. ID NO. 104 is the deduced amino acid sequence of TbH-9-1. 10

SEQ. ID NO. 105 is the DNA sequence of TbH-9-4.

SEQ. ID NO. 106 is the deduced amino acid sequence of TbH-9-4.

SEQ. ID NO. 107 is the DNA sequence of Tb38-1F2 IN.

SEQ. ID NO. 108 is the DNA sequence of Tb38-1F2 RP.

15 SEQ. ID NO. 109 is the deduced amino acid sequence of Tb37-FL.

SEQ. ID NO. 110 is the deduced amino acid sequence of Tb38-IN.

SEQ. ID NO. 111 is the DNA sequence of Tb38-1F3.

SEQ. ID NO. 112 is the deduced amino acid sequence of Tb38-1F3.

SEQ. ID NO. 113 is the DNA sequence of Tb38-1F5.

20 SEQ. ID NO. 114 is the DNA sequence of Tb38-1F6.

SEQ. ID NO. 115 is the deduced N terminal amino acid sequence of DPV.

SEQ. ID NO. 116 is the deduced N terminal amino acid sequence of AVGS.

SEQ. ID NO. 117 is the deduced N-terminal amino acid sequence of AAMK.

SEQ. ID NO. 118 is the deduced N-terminal amino acid sequence of YYWC

25 SEQ. ID NO. 119 is the deduced Naterminal amino acid sequence of DIGS.

SLO HINNING CONTRACTOR

Social and the control of the control of the community DPAS Same the state of the state of

SEQ ID NO. 125-128 are the protein sequences of four DPPD cyanogen bromide fragments.

SEQ ID NO. 129 is the N-terminal protein sequence of XDS antigen.

SEQ ID NO. 130 is the N-terminal protein sequence of AGD antigen.

5 SEQ ID NO. 131 is the N-terminal protein sequence of APE antigen.

SEQ ID NO. 132 is the N-terminal protein sequence of XYI antigen.

SEQ ID NO. 133 is the DNA sequence of TbH-29.

SEQ ID NO. 134 is the DNA sequence of TbH-30.

SEQ ID NO. 135 is the DNA sequence of TbH-32.

SEQ ID NO. 136 is the DNA sequence of TbH-33.

SEQ ID NO. 137 is the predicted amino acid sequence of TbH-29.

SEQ ID NO. 138 is the predicted amino acid sequence of TbH-30.

SEQ ID NO. 139 is the predicted amino acid sequence of TbH-32.

SEQ ID NO. 140 is the predicted amino acid sequence of TbH-33.

SEQ ID NO: 141-146 are PCR primers used in the preparation of a fusion protein containing TbRa3, 38 kD and Tb38-1.

SEQ ID NO: 147 is the DNA sequence of the fusion protein containing TbRa3, 38 kD and Tb38-1.

SEQ ID NO: 148 is the amino acid sequence of the fusion protein containing TbRa3.

20 38 kD and Tb38-1.

SEQ ID NO: 149 is the DNA sequence of the M. tuberculosis antigen 38 kD.

SEQ ID NO: 150 is the amino acid sequence of the M. tuberculosis antigen 38 kD.

SEQ ID NO: 151 is the DNA sequence of XP14.

SEQ ID NO: 152 is the DNA sequence of XP24.

25 SEQ ID NO: 153 is the DNA sequence of XP31

SEQUID NOTE: The translation of the expression and the contraction of the contraction of

	SEQ ID NO: 158 is the DNA sequence of XP27.
	SEQ ID NO: 159 is the DNA sequence of XP36.
	SEQ ID NO: 160 is the 5' DNA sequence of XP4.
	SEQ ID NO: 161 is the 5' DNA sequence of XP5.
5	SEQ ID NO: 162 is the 5' DNA sequence of XP17.
	SEQ ID NO: 163 is the 5° DNA sequence of XP30
	SEQ ID NO: 164 is the 5° DNA sequence of XP2.
	SEQ ID NO: 165 is the 3° DNA sequence of XP2.
	SEQ ID NO: 166 is the 5' DNA sequence of XP3.
10	SEQ ID NO: 167 is the 3° DNA sequence of XP3.
	SEQ ID NO: 168 is the 5' DNA sequence of XP6.
	SEQ ID NO: 169 is the 3' DNA sequence of XP6.
	SEQ ID NO: 170 is the 5° DNA sequence of XP18.
	SFQ ID NO: 171 is the 3° DNA sequence of XP18.
15	SFQ ID NO: 172 is the 5° DNA sequence of XP19.
	SEQ ID NO: 173 is the 3' DNA sequence of XP19.
	SEQ ID NO: 174 is the 5' DNA sequence of XP22.
	SEQ ID NO: 175 is the 3' DNA sequence of XP22.
	SEQ ID NO: 176 is the 5' DNA sequence of XP25
20	SEQ ID NO: 177 is the 31 DNA sequence of XP25.
	SEQ ID NO: 178 is the full-length DNA sequence of TbH4-XPL.
	SEQ ID NO: 179 is the predicted amino acid sequence of TbH4-XP1.
	SEQ ID NO: 180 is the predicted amino acid sequence encoded by the revers
	complement of TbH4-XP1.

SEQ ID NO: 181 is a first predicted amino acid sequence encoded by XP36. SEQ ID NO: 1824(2) and sequence encoded by XP36.

Scould's Six District

30 StO HO 20 15 1 16 DNA Agrence of 2011.

SEQ ID NO: 186 is the DNA sequence of RDIF8.

SEQ ID NO: 187 is the DNA sequence of RDIF10.

SEQ ID NO: 188 is the DNA sequence of RDIF11.

SEQ ID NO: 189 is the predicted amino acid sequence of RDIF2.

5 SEQ ID NO: 190 is the predicted amino acid sequence of RDIF5.

SEQ ID NO: 191 is the predicted amino acid sequence of RDIF8.

SEQ ID NO: 192 is the predicted amino acid sequence of RDIF10.

SEQ ID NO: 193 is the predicted amino acid sequence of RDIF11.

SEQ ID NO: 194 is the 5' DNA sequence of RDIF12.

SEQ ID NO: 195 is the 3' DNA sequence of RDIF12.

SEQ ID NO: 196 is the DNA sequence of RDIF7.

SEQ ID NO: 197 is the predicted amino acid sequence of RDIF7.

SEQ ID NO: 198 is the DNA sequence of DIF2-1.

SEQ ID NO: 199 is the predicted amino acid sequence of DIF2-1.

SEQ ID NO: 200-207 are PCR primers used in the preparation of a fusion protein containing TbRa3, 38 kD, Tb38-1 and DPEP (hereinafter referred to as TbF-2).

SEQ ID NO: 208 is the DNA sequence of the fusion protein TbF-2.

SEQ ID NO: 209 is the amino acid sequence of the fusion protein TbF-2.

20

### DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for diagnosing tuberculosis. The compositions of the subject invention include polypeptides that comprise at least one antigenic portion of a *M tuberculosis* antigen, or a variant of such an antigen that differs only in conservative substitutions and or modifications

the first of the first of the second of the second will be selected as a fill condition of the second secon

14

a polypeptide comprising an antigenic portion of one of the above antigens may consist entirely of the antigenic portion, or may contain additional sequences. The additional sequences may be derived from the native *M. tuberculosis* antigen or may be heterologous, and such sequences may (but need not) be antigenic.

5

15

An "antigenic portion" of an antigen (which may or may not be soluble) is a portion that is capable of reacting with sera obtained from an *M tuberculosis*-infected individual (*i.e.*, generates an absorbance reading with sera from infected individuals that is at least three standard deviations above the absorbance obtained with sera from uninfected individuals, in a representative ELISA assay described herein). An "*M. tuberculosis*-infected individual" is a human who has been infected with *M. tuberculosis* (*e.g.*, has an intradermal skin test response to PPD that is at least 0.5 cm in diameter). Infected individuals may display symptoms of tuberculosis or may be free of disease symptoms. Polypeptides comprising at least an antigenic portion of one or more *M. tuberculosis* antigens as described herein may generally be used, alone or in combination, to detect tuberculosis in a patient.

The compositions and methods of this invention also encompass variants of the above polypeptides. A "variant," as used herein, is a polypeptide that differs from the native antigen only in conservative substitutions and/or modifications, such that the antigenic properties of the polypeptide are retained. Such variants may generally be identified by modifying one of the above polypeptide sequences, and evaluating the antigenic properties of the modified polypeptide using, for example, the representative procedures described herein.

A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids represent conservative changes: (1) ala, pro, ply, plu, asp, gln, asn, ser, thr: (2) eys, ser, tyr, thr. (3) values.

the charge and involve partice matrix of the propagation of the exclupping of non-negative may be assumed that the charge of the form

translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

In a related aspect, combination polypeptides are disclosed. A "combination polypeptide" is a polypeptide comprising at least one of the above antigenic portions and one or more additional antigenic *M. tuberculosis* sequences, which are joined via a peptide linkage into a single amino acid chain. The sequences may be joined directly (*i.e.*, with no intervening amino acids) or may be joined by way of a linker sequence (*e.g.*, Gly-Cys-Gly) that does not significantly diminish the antigenic properties of the component polypeptides.

In general, *M. tuberculosis* antigens, and DNA sequences encoding such antigens, may be prepared using any of a variety of procedures. For example, soluble antigens may be isolated from *M. tuberculosis* culture filtrate by procedures known to those of ordinary skill in the art, including anion-exchange and reverse phase chromatography. Purified antigens may then be evaluated for a desired property, such as the ability to react with sera obtained from an *M. tuberculosis*-infected individual. Such screens may be performed using the representative methods described herein. Antigens may then be partially sequenced using, for example, traditional Edman chemistry. *Sec* Edman and Berg, *Fur J. Biochem.* 80:116-132, 1967.

15

20

m. 1. přístí v Sarger († 1967) M. 1. pokovane († 1967) v Sar († 1967) v Coli Sprin.
 Plarbye Larieratorica C. Št. pro (\* Data v S. ) († 1967)

16

DNA sequences encoding soluble antigens may also be obtained by screening an appropriate *M. tuberculosis* eDNA or genomic DNA library for DNA sequences that hybridize to degenerate oligonucleotides derived from partial amino acid sequences of isolated soluble antigens. Degenerate oligonucleotide sequences for use in such a screen may be designed and synthesized, and the screen may be performed, as described (for example) in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY (and references cited therein). Polymerase chain reaction (PCR) may also be employed, using the above oligonucleotides in methods well known in the art, to isolate a nucleic acid probe from a cDNA or genomic library. The library screen may then be performed using the isolated probe.

Regardless of the method of preparation, the antigens described herein are "antigenic." More specifically, the antigens have the ability to react with sera obtained from an *M. tuberculosis*-infected individual. Reactivity may be evaluated using, for example, the representative ELISA assays described herein, where an absorbance reading with sera from infected individuals that is at least three standard deviations above the absorbance obtained with sera from uninfected individuals is considered positive.

Antigenic portions of M tuberculosis antigens may be prepared and identified using well known techniques, such as those summarized in Paul, Fundamental Immunology, 3d ed., Raven Press, 1993, pp. 243-247 and references cited therein. Such techniques include screening polypeptide portions of the native antigen for antigenic properties. The representative FLISAs described herein may generally be employed in these screens. An antigenic portion of a polypeptide is a portion that, within such representative assays, generates a signal in such assays that is substantially similar to that generated by the full length antigen. In other words, an antigenic portion of a M tuberculosis antigen generates at least about  $20^{6}$ , and preferably about  $100^{6}$ , of the signal induced by the full length antigen

and the first of the second of the about the problem and the may be expected using the integral and the content of the second of

commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, J. Am. Chem. Soc. 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied BioSystems, Inc., Foster City, CA, and may be operated according to the manufacturer's instructions. Variants of a native antigen may generally be prepared using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis. Sections of the DNA sequence may also be removed using standard techniques to permit preparation of truncated polypeptides.

Recombinant polypeptides containing portions and/or variants of a native antigen may be readily prepared from a DNA sequence encoding the polypeptide using a variety of techniques well known to those of ordinary skill in the art. For example, supernatants from suitable host/vector systems which secrete recombinant protein into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant protein.

10

15

25

MH: -1

Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides as described herein. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher enkaryotic cells. Preferably, the host cells employed are *E-coli*, yeast or a mammalian cell line, such as COS or CHO. The DNA sequences expressed in this manner may encode naturally occurring antigens, portions of naturally occurring antigens, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in substantially possibles. Productive of the disclosed

In certain specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a soluble *M. tuberculosis* antigen (or a variant of such an antigen), where the antigen has one of the following N-terminal sequences:

- (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 115),
- (b) Ala-Val-Glu-Ser-Gly-Mct-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 116);
- (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg (SEQ ID NO: 117);
- 10 (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 118);

5

- (c) Asp-lle-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID NO: 119);
- (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SEQ ID NO: 120);
- (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser (SEO ID NO: 121);
- (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly (SFQ ID NO. 122);
- 20 Asp Pro Ala Ser-Ala Pro-Asp-Val-Pro-Hhr-Ala-Ala-Gln-Gln-Fhr-Ser-Leu Leu-Asm-Ser-Leu-Ala Asp Pro Asn Val Ser Phe-Ala-Asn (SEQ ID NO: 123);
  - (j) Xaa-Asp-Ser-Giu-Lys-Ser-Ala-Thr-He-Lys-Val-Thr-Asp-Ala-Ser;(SEQ ID NO 129)
- 25 (k) Ala-Gly-Asp-Thr-Naa-He-Tyr-He-Val-Gly-Asn-Leu-Thr-Ala-Asp;

Figure with the second of the

15

amino acid sequence of which is provided in SEQ ID NO: 53. A DNA sequence encoding the antigen identified as (a) above is provided in SEQ ID NO: 96; its deduced amino acid sequence is provided in SEQ ID NO: 97. A DNA sequence corresponding to antigen (d) above is provided in SEQ ID NO: 24, a DNA sequence corresponding to antigen (c) is provided in SEQ ID NO: 25 and a DNA sequence corresponding to antigen (I) is disclosed in SEQ ID NO: 94 and its deduced amino acid sequence is provided in SEQ ID NO: 95.

In a further specific embodiment, the subject invention discloses polypeptides comprising at least an immunogenic portion of an *M. tuberculosis* antigen having one of the following N-terminal sequences, or a variant thereof that differs only in conservative substitutions and/or modifications:

- (m) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132) or
- (n) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124)

wherein Xaa may be any amino acid, preferably a cysteine residue.

In other specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a soluble *M. tuberculosis* antigen (or a variant of such an antigen) that comprises one or more of the amino acid sequences encoded by (a) the DNA sequences of SEQ ID NOS: 1, 2, 4 10, 13-25, 52, 94 and 96, (b) the complements of such DNA sequences, or (c) DNA sequences substantially homologous to a sequence in (a) or (b).

In further specific embodiments, the subject invention discloses polypeptides comprising at least an antigenic portion of a *M-tuberculosis* antigen (or a variant of such an antigen), which may or may not be soluble, that comprises one or more of the main antigen.

15

or more of DNA sequences specifically recited herein. "Substantial homology," as used herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the event of cross-species homology, at 45°C with 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS). Such hybridizing DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known *M tuberculosis* antigen, such as the 38 kD antigen described above or ESAT-6 (SEQ ID NOS: 98 and 99), together with variants of such fusion proteins. The fusion proteins of the present invention may also include a linker peptide between the first and second polypeptides.

A DNA sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate DNA sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a DNA sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its

of a first sentermation. If the membrate it is depthic a similar concernation could interact the system of a Laprace country text on the sentence of the sente

or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene 40*:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA 83*:8258-8562, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric hindrance.

10

In another aspect, the present invention provides methods for using the polypeptides described above to diagnose tuberculosis. In this aspect, methods are provided for detecting *M. tuberculosis* infection in a biological sample, using one or more of the above polypeptides, alone or in combination. In embodiments in which multiple polypeptides are employed, polypeptides other than those specifically described herein, such as the 38 ktD antigen described in Andersen and Hansen, *Infect. Immun.* 57:2481-2488, 1989, may be included. As used herein, a "biological sample" is any antibody-containing sample obtained from a patient. Preferably, the sample is whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid or urine. More preferably, the sample is a blood, serum or plasma sample obtained from a patient or a blood supply. The polypeptide(s) are used in an assay, as described below, to determine the presence or absence of antibodies to the polypeptide(s) in the sample, relative to a predetermined cut off value. The presence of such antibodies indicates previous sensitization to mycobacterial antigens which may be indicative of tuberculosis.

In embodiments in which more than one polypeptide is employed, the polypeptides used are preferably complementary (i.e., one component polypeptide will tend

utroat tota interpretatives. More versus of America maining semiliar into the progression of the progression

formulated that are capable of detecting infection in most, or all, of the samples tested. Such polypeptides are complementary. For example, approximately 25-30% of sera from tuberculosis-infected individuals are negative for antibodies to any single protein, such as the 38 kD antigen mentioned above. Complementary polypeptides may, therefore, be used in combination with the 38 kD antigen to improve sensitivity of a diagnostic test.

There are a variety of assay formats known to those of ordinary skill in the art for using one or more polypeptides to detect antibodies in a sample. See, e.g., Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988, which is incorporated herein by reference. In a preferred embodiment, the assay involves the use of polypeptide immobilized on a solid support to bind to and remove the antibody from the sample. The bound antibody may then be detected using a detection reagent that contains a reporter group. Suitable detection reagents include antibodies that bind to the antibody/polypeptide complex and free polypeptide labeled with a reporter group (e.g., in a semi-competitive assay). Alternatively, a competitive assay may be utilized, in which an antibody that binds to the polypeptide is labeled with a reporter group and allowed to bind to the immobilized antigen after incubation of the antigen with the sample. The extent to which components of the sample inhibit the binding of the labeled antibody to the polypeptide is indicative of the reactivity of the sample with the immobilized polypeptide

The solid support may be any solid material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681.

20

25 The polypeptides may be bound to the solid support using a variety of

tract follows without the first open may be a linkage

membrane is preferred. In such cases, adsorption may be achieved by contacting the polypeptide, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of polypeptide ranging from about 10 ng to about 1  $\mu$ g, and preferably about 100 ng, is sufficient to bind an adequate amount of antigen.

Covalent attachment of polypeptide to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the polypeptide. For example, the polypeptide may be bound to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the polypeptide (*see*, *e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is an enzyme linked immunosorbent assay (ELISA). This assay may be performed by first contacting a polypeptide antigen that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that antibodies to the polypeptide within the sample are allowed to bind to the immobilized polypeptide. Unbound sample is then removed from the immobilized polypeptide and a detection reagent capable of binding to the immobilized antibody-polypeptide complex is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific detection reagent.

15

25

More specifically, once the polypeptide is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 2018 (Sigma Chemical Co., St. Louis, MO) may be employed. The

The second period of tractifiants afficient to despet the presence of intribute expining a

of binding that is at least 95% of that achieved at equilibrium between bound and unbound antibody. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20<sup>TM</sup>. Detection reagent may then be added to the solid support. An appropriate detection reagent is any compound that binds to the immobilized antibody-polypeptide complex and that can be detected by any of a variety of means known to those in the art. Preferably, the detection reagent contains a binding agent (such as, for example, Protein A, Protein G, immunoglobulin, lectin or free antigen) conjugated to a reporter group. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of binding agent to reporter group may be achieved using standard methods known to those of ordinary skill in the art. Common binding agents may also be purchased conjugated to a variety of reporter groups from many commercial sources (e.g., Zymed Laboratories, San Francisco, CA, and Pierce, Rockford, IL).

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound antibody. An appropriate amount of time may generally be determined from the manufacturer's instructions or by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the teporter group. For radioactive groups, scintillation counting or autoradiographic methods

reporter around mass sectorally by three-terrors in the externor to resituate elemenable for a sector of the externor of the e

To determine the presence or absence of anti-M. tuberculosis antibodies in the sample, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value is the average mean signal obtained when the immobilized antigen is incubated with samples from an uninfected patient. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for tuberculosis. In an alternate preferred embodiment, the cutoff value is determined using a Receiver Operator Curve, according to the method of Sackett et al., Clinical Epidemiology. A Basic Science for Clinical Medicine, Little Brown and Co., 1985, pp. 106-107. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for tuberculosis.

10

15

20

In a related embodiment, the assay is performed in a rapid flow-through or strip test format, wherein the antigen is immobilized on a membrane, such as nitrocellulose. In the flow-through test, antibodies within the sample bind to the immobilized polypeptide as the sample passes through the membrane. A detection reagent (e.g., protein A-colloidal gold) then binds to the antibody polypeptide complex as the solution containing the detection reagent flows through the membrane. The detection of bound detection reagent may then be

the state of the second state of the second state of the polypeptide redicates the

detection reagent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of polypeptide immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of antibodies that would be sufficient to generate a positive signal in an ELISA, as discussed above. Preferably, the amount of polypeptide immobilized on the membrane ranges from about 25 ng to about 1 µg, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount (e.g., one drop) of patient serum or blood.

Of course, numerous other assay protocols exist that are suitable for use with the polypeptides of the present invention. The above descriptions are intended to be exemplary only.

In yet another aspect, the present invention provides antibodies to the inventive polypeptides. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

15

from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Antibodies may be used in diagnostic tests to detect the presence of *M tuberculosis* antigens using assays similar to those detailed above and other techniques well known to those of skill in the art, thereby providing a method for detecting *M tuberculosis* infection in a patient.

Diagnostic reagents of the present invention may also comprise DNA sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain

ONAN them is to to a fine technique much anomal in the interface coefficients for executionnoises.

10

present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 80%, preferably at least about 90% and more preferably at least about 95%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule encoding one of the polypeptides disclosed herein. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule encoding one of the polypeptides disclosed herein. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al. Ibid; Ehrlich, Ibid). Primers or probes may thus be used to detect M. tuberculosis-specific sequences in biological samples. DNA probes or primers comprising oligonucleotide sequences described above may be used alone, in combination with each other, or with previously identified sequences, such as the 38 kD antigen discussed above.

The following Examples are offered by way of illustration and not by way of 20. limitation.

# EXAMPLES

### EXAMPLE 1

25

PERIFICATION AND CHARACTERIZATION OF POLYPEPTIDES

ipositi mana and a company

List of the state of the state

*M. tuberculosis* (either H37Ra, ATCC No. 25177, or H37Rv, ATCC No. 25618) was cultured in sterile GAS media at 37°C for fourteen days. The media was then vacuum filtered (leaving the bulk of the cells) through a 0.45  $\mu$  filter into a sterile 2.5 L bottle. The media was then filtered through a 0.2  $\mu$  filter into a sterile 4 L bottle. NaN<sub>3</sub> was then added to the culture filtrate to a concentration of 0.04%. The bottles were then placed in a 4°C cold room.

The culture filtrate was concentrated by placing the filtrate in a 12 L reservoir that had been autoclaved and feeding the filtrate into a 400 ml Amicon stir cell which had been rinsed with ethanol and contained a 10,000 kDa MWCO membrane. The pressure was maintained at 60 psi using nitrogen gas. This procedure reduced the 12 L volume to approximately 50 ml.

The culture filtrate was then dialyzed into 0.1% ammonium bicarbonate using a 8,000 kDa MWCO cellulose ester membrane, with two changes of ammonium bicarbonate solution. Protein concentration was then determined by a commercially available BCA assay (Pierce, Rockford, IL).

15

The dialyzed culture filtrate was then lyophilized, and the polypeptides resuspended in distilled water. The polypeptides were then dialyzed against 0.01 mM 1,3 bis[tris(hydroxymethyl)-methylamino]propane, pH 7.5 (Bis-Tris propane buffer), the initial conditions for anion exchange chromatography. Fractionation was performed using gel profusion chromatography on a POROS 146 H Q M anion exchange column 4.6 mm x 100 mm (Perseptive BioSystems, Framingham, MA) equilibrated in 0.01 mM Bis-Tris propane buffer pH 7.5. Polypeptides were cluted with a linear 0-0.5 M NaCl gradient in the above buffer system. The column eluent was monitored at a wavelength of 220 nm.

The pools of polypeptides eluting from the ion exchange column were dialyzed against distilled water and lyophilized. The resulting material was dissolved in 0.1% triplyoropastics (1.1.1) (1.5.11.1.3)

interest to the second HAM is a mortal to the state of the wave of main material field HPLC and plants a magnificating MA report to the contract of the second seco

to maximize the purity of the individual samples. Approximately 200 purified polypeptides were obtained.

The purified polypeptides were then screened for the ability to induce T-cell proliferation in PBMC preparations. The PBMCs from donors known to be PPD skin test positive and whose T cells were shown to proliferate in response to PPD and crude soluble proteins from MTB were cultured in medium comprising RPMI 1640 supplemented with 10% pooled human serum and 50 µg/ml gentamicin. Purified polypeptides were added in duplicate at concentrations of 0.5 to 10 µg/ml. After six days of culture in 96-well round-bottom plates in a volume of 200 µl, 50 µl of medium was removed from each well for determination of IFN-γ levels, as described below. The plates were then pulsed with 1 µCi/well of tritiated thymidine for a further 18 hours, harvested and tritium uptake determined using a gas scintillation counter. Fractions that resulted in proliferation in both replicates three fold greater than the proliferation observed in cells cultured in medium alone were considered positive.

IFN-γ was measured using an enzyme-linked immunosorbent assay (ELISA). ELISA plates were coated with a mouse monoclonal antibody directed to human IFN-γ (Chemicon) in PBS for four hours at room temperature. Wells were then blocked with PBS containing 5% (W/V) non-fat dried milk for 1 hour at room temperature. The plates were then washed six times in PBS 0.2% TWFEN-20 and samples diluted 1½ in culture medium in the ELISA plates were incubated overnight at room temperature. The plates were again washed and a polyelonal rabbit anti-human IFN-γ serum diluted 1:3000 in PBS/10% normal goat serum was added to each well. The plates were then incubated for two hours at room temperature, washed and horseradish peroxidase coupled anti-rabbit IgG (Jackson Labs.) was added at a 1:2000 dilution in PBS/5% non-fat dried milk. After a further two hour incubation at room temperature, the plates were washed and IMB substrate added. The reaction was

20

For sequencing, the polypeptides were individually dried onto Biobrene<sup>TM</sup> (Perkin Elmer/Applied BioSystems Division, Foster City, CA) treated glass fiber filters. The filters with polypeptide were loaded onto a Perkin Elmer/Applied BioSystems Division Procise 492 protein sequencer. The polypeptides were sequenced from the amino terminal and using traditional Edman chemistry. The amino acid sequence was determined for each polypeptide by comparing the retention time of the PTH amino acid derivative to the appropriate PTH derivative standards.

Using the procedure described above, antigens having the following N-terminal sequences were isolated:

- 10 (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Thr-Xaa-Asn-Tyr-Gly-Gln- (b) Val-Val-Ala-Ala-Leu (SEQ ID NO: 54);
  - (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala-Leu-Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 55);
  - (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg (SEQ ID NO: 56);
  - (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 57);
  - (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Xaa-Ala-Val (SEQ ID NO 58):
  - (f) Ala-Glu-Glu-Ser Ile-Ser-Thr-Xaa-Glu-Xaa-Ile-Val-Pro (SFQ ID) NO: 59):
    - (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Ala-Ala-Ala-Ala-Pro-Pro-Ala (SEQ ID NO: 60); and
- (h) Ala-Pro I ys Thr Tyr-Xaa-Glu-Glu-Leu-I ys-Gly-Thr-Asp-Thr-Gly (SLQ ID NO: 61):

the second of th

maying the first of the control of the material for although previously described was to the control of the con

City, CA) with a 7 micron pore size, column size 1 mm x 100 mm, in a Perkin Elmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted from the column with a linear gradient of 1%/minute of acetonitrile (containing 0.05% TFA) in water (0.05% TFA) at a flow rate of 80 µl/minute. The eluent was monitored at 250 nm. The original fraction was separated into 4 major peaks plus other smaller components and a polypeptide was obtained which was shown to have a molecular weight of 12.054 Kd (by mass spectrometry) and the following N-terminal sequence:

(i) Asp-Pro-Ala-Ser-Ala-Pro-Asp-Val-Pro-Thr-Ala-Ala-Gln-Gln-Thr-Ser-Leu-Leu-Asn-Asp-Leu-Ala-Asp-Pro-Asp-Val-Ser-Phe-Ala-Asp (SEQ ID NO: 62).

This polypeptide was shown to induce proliferation and IFN- $\gamma$  production in PBMC preparations using the assays described above.

Additional soluble antigens were isolated from *M. tuberculosis* culture filtrate as follows. *M. tuberculosis* culture filtrate was prepared as described above. Following dialysis against Bis-Tris propane buffer, at pH 5.5, fractionation was performed using anion exchange chromatography on a Poros QE column 4.6 x 100 mm (Perseptive Biosystems) equilibrated in Bis-Tris propane buffer pH 5.5. Polypeptides were eluted with a linear 0-1.5 M NaCl gradient in the above buffer system at a flow rate of 10 ml/min. The column eluent was monitored at a wavelength of 214 nm.

The fractions eluting from the ion exchange column were pooled and subjected to reverse phase chromatography using a Poros R2 column  $4.6 \times 100$  mm (Perseptive Biosystems). Polypeptides were eluted from the column with a linear gradient from 0-100% acetonitrile (0.1% TFA) at a flow rate of 5 ml/min. The eluent was monitored at 214 nm.

25 Fractions containing the eluted polypeptides were lyophilized and resuspended

10

The fraction with biological activity was separated into one major peak plus other smaller components. Western blot of this peak onto PVDF membrane revealed three major bands of molecular weights 14 Kd, 20 Kd and 26 Kd. These polypeptides were determined to have the following N-terminal sequences, respectively:

- 5 (j) Xaa-Asp-Ser-Glu-Lys-Ser-Ala-Thr-Ile-Lys-Val-Thr-Asp-Ala-Ser; (SEO ID NO: 129)
  - (k) Ala-Gly-Asp-Thr-Xaa-Ile-Tyr-Ile-Val-Gly-Asn-Leu-Thr-Ala-Asp; (SEQ ID NO: 130) and
  - (I) Ala-Pro-Glu-Ser-Gly-Ala-Gly-Leu-Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID NO: 131), wherein Xaa may be any amino acid.

Using the assays described above, these polypeptides were shown to induce proliferation and IFN-γ production in PBMC preparations. Figs. 1A and B show the results of such assays using PBMC preparations from a first and a second donor, respectively.

above were obtained by screening a *M. tuberculosis* genomic library using <sup>32</sup>P end labeled degenerate oligonucleotides corresponding to the N-terminal sequence and containing *M. tuberculosis* codon bias. The screen performed using a probe corresponding to antigen (a) above identified a clone having the sequence provided in SEQ ID NO: 96. The polypeptide encoded by SEQ ID NO: 96 is provided in SEQ ID NO: 97. The screen performed using a probe corresponding to antigen (g) above identified a clone having the sequence provided in SEQ ID NO: 52. The polypeptide encoded by SEQ ID NO: 52 is provided in SEQ ID NO: 53. The screen performed using a probe corresponding to antigen (d) above identified a clone having the sequence provided in SEQ ID NO: 53. The screen performed using a probe corresponding to antigen (d) above identified a clone having the sequence provided in SEQ ID NO: 24, and the screen performed with a probe corresponding to antigen (e) identified a clone having the sequence provided in SEQ ID NO: 25.

professional grandes. Note on Silvino influence embologies to the amino desilving teneds for a

15

The amino acid sequence for antigen (i) was found to be homologous to a sequence from *M. leprae*. The full length *M. leprae* sequence was amplified from genomic DNA using the sequence obtained from GENBANK. This sequence was then used to screen an *M. tuberculosis* library and a full length copy of the *M. tuberculosis* homologue was obtained (SEQ ID NO: 94).

The amino acid sequence for antigen (j) was found to be homologous to a known *M. tuberculosis* protein translated from a DNA sequence. To the best of the inventors' knowledge, this protein has not been previously shown to possess T-cell stimulatory activity. The amino acid sequence for antigen (k) was found to be related to a sequence from *M. leprae*.

In the proliferation and IFN- $\gamma$  assays described above, using three PPD positive donors, the results for representative antigens provided above are presented in Table 1:

<u>IABLE 1</u>

<u>RESULTS OF PBMC PROLIFERATION AND IFN-y ASSAYS</u>

Sequence	Proliferation	IFN-γ
(a)	+	
 (c)	311	+++
(d)	++	++
(g)	+++	+++
 (h)	+++	+++

In Table 1, responses that gave a stimulation index (SI) of between 2 and 4

The second of th

5

25

indicate that these antigens are capable of inducing proliferation and/or interferon-y production.

#### **EXAMPLE 2**

#### USE OF PATIENT SERA TO ISOLATE M. TUBERCULOSIS ANTIGENS

This example illustrates the isolation of antigens from *M. tuberculosis* lysate by screening with serum from *M. tuberculosis*-infected individuals.

Dessicated *M. tuberculosis* H37Ra (Difco Laboratories) was added to a 2% NP40 solution, and alternately homogenized and sonicated three times. The resulting suspension was centrifuged at 13,000 rpm in microfuge tubes and the supernatant put through a 0.2 micron syringe filter. The filtrate was bound to Macro Prep DEAE beads (BioRad, Hercules, CA). The beads were extensively washed with 20 mM Tris pH 7.5 and bound proteins eluted with 1M NaCl. The NaCl elute was dialyzed overnight against 10 mM Tris, pH 7.5. Dialyzed solution was treated with DNase and RNase at 0.05 mg/ml for 30 min. at room temperature and then with α-D-mannosidase, 0.5 U/mg at pH 4.5 for 3-4 hours at room temperature. After returning to pH 7.5, the material was fractionated via FPLC over a Bio Scale-Q-20 column (BioRad). Fractions were combined into nine pools, concentrated in a Centriprep 10 (Amicon, Beverley, MA) and screened by Western blot for scrological activity using a serum pool from *M. inherculosis*-infected patients which was not immunoreactive with other antigens of the present invention

The most reactive fraction was run in SDS-PAGE and transferred to PVDE. A band at approximately 85 Kd was cut out yielding the sequence:

(m) Xaa-Tyr-lle-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-lle-Val-Pro-Gly-Lys-Ile-Asn Val-His-Leu-Val: (SEQ ID NO: 132), wherein Xaa may be any

The first contents that model of continuous standard in this desire was

degenerate oligonucleotides corresponding to the N-terminal sequence of SEQ ID NO:137. A clone was identified having the DNA sequence provided in SEQ ID NO: 198. This sequence was found to encode the amino acid sequence provided in SEQ ID NO: 199. Comparison of these sequences with those in the genebank revealed some similarity to sequences previously identified in *M. tuberculosis* and *M. bovis*.

## EXAMPLE 3

#### PREPARATION OF DNA SEQUENCES ENCODING M. TUBERCULOSIS ANTIGENS

This example illustrates the preparation of DNA sequences encoding *M. tuberculosis* antigens by screening a *M. tuberculosis* expression library with sera obtained from patients infected with *M. tuberculosis*, or with anti-sera raised against *M. tuberculosis* antigens.

# 15 A. <u>Preparation of M. Tuberculosis Soluble Antigens using Rabbit Anti-sera</u> <u>Raised against M. Tuberculosis Supernatant</u>

Genomic DNA was isolated from the *M. tuberculosis* strain H37Ra. The DNA was randomly sheared and used to construct an expression library using the Lambda ZAP expression system (Stratagene, La Jolla, CA). Rabbit anti-sera was generated against secretory proteins of the *M. tuberculosis* strains H37Ra, H37Rv and Erdman by immunizing a rabbit with concentrated supernatant of the *M. tuberculosis* cultures. Specifically, the rabbit was first immunized subcutaneously with 200 µg of protein antigen in a total volume of 2 ml containing 100 µg muramyl dipeptide (Calbiochem, La Jolla, CA) and 1 ml of incomplete Freund's adjuvant. Four weeks later the rabbit was boosted subcutaneously with 100 µg antigen in incomplete Freund's adjuvant. Finally, the rabbit was immunized intravenously

estre in the content of the content of the property of the content is an entire with a section of the content o

Thirty two clones were purified. Of these, 25 represent sequences that have not been previously identified in *M. tuberculosis*. Proteins were induced by IPTG and purified by gel clution, as described in Skeiky et al., *J. Exp. Med. 181*:1527-1537, 1995. Representative partial sequences of DNA molecules identified in this screen are provided in SEQ ID NOS: 1-25. The corresponding predicted amino acid sequences are shown in SEQ ID NOS: 64-38.

On comparison of these sequences with known sequences in the gene bank using the databases described above, it was found that the clones referred to hereinafter as TbRA2A, TbRA16, TbRA18, and TbRA29 (SEQ ID NOS: 77, 69, 71, 76) show some homology to sequences previously identified in *Mycobacterium leprae* but not in *M. tuberculosis*. TbRA11, TbRA26, TbRA28 and TbDPEP (SEQ ID NOS: 66, 74, 75, 53) have been previously identified in *M. tuberculosis*. No significant homologies were found to TbRA1, TbRA3, TbRA4. TbRA9, TbRA10, TbRA13, TbRA17, TbRA19, TbRA29, TbRA32, TbRA36 and the overlapping clones TbRA35 and TbRA12 (SEQ ID NOS: 64, 78, 82, 83, 65, 68, 76, 72, 76, 79, 81, 80, 67, respectively). The clone TbRa24 is overlapping with clone TbRa29.

# B. <u>Use of Sera from Patients having Pulmonary or Pleural Tuberculosis to</u> <u>Identify DNA Sequences Encoding Merculosis Antigens</u>

20 The genomic DNA library described above, and an additional H37Rv library, were screened using pools of sera obtained from patients with active tuberculosis. To prepare the H37Rv library, *M tuberculosis* strain H37Rv genomic DNA was isolated, subjected to partial Sau3A digestion and used to construct an expression library using the Lambda Zap expression system (Stratagene, La Jolla, Ca). Three different pools of sera, each containing sera obtained from three individuals with active pulmonary or pleural disease, were used in

rise bette for All Poyloiders, and the Color of the Color

lacked increased reactivity with the recombinant 38 kD *M. tuberculosis* H37Ra phosphate-binding protein.

All pools were pre-adsorbed with *E. coli* lysate and used to screen the H37Ra and H37Rv expression libraries, as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones deduced.

Thirty two clones were purified. Of these, 31 represented sequences that had not been previously identified in human *M. tuberculosis*. Representative sequences of the DNA molecules identified are provided in SEQ ID NOS:: 26-51 and 100. Of these, TbH-8-2 (SEQ. ID NO. 100) is a partial clone of TbH-8, and TbH-4 (SEQ. ID NO. 43) and TbH-4-FWD (SEQ. ID NO. 44) are non-contiguous sequences from the same clone. Amino acid sequences for the antigens hereinafter identified as Tb38-1, TbH-4, TbH-8, TbH-9, and TbH-12 are shown in SEQ ID NOS.: 89-93. Comparison of these sequences with known sequences in the gene bank using the databases identified above revealed no significant homologies to TbH-4, TbH-8, TbH-9 and TbM-3, although weak homologies were found to TbH-9. TbH-12 was found to be homologous to a 34 kD antigenic protein previously identified in *M. paratuberculosis* (Acc. No. S28515). Tb38-1 was found to be located 34 base pairs upstream of the open reading frame for the antigen ESA1-6 previously identified in *M. bovis* (Acc. No. U34848) and in *M. tuberculosis* (Sorensen et al., *Infec. Immun* 63:1710-1717, 1995).

15

Probes derived from 1b38-1 and 1bH-9, both isolated from an H37Ra library, were used to identify clones in an H37Rv library. Tb38-1 hybridized to Tb38-1F2, Tb38-1F3, Tb38-1F5 and fb38-1F6 (SEQ, ID NOS: 107, 108, 111, 113, and 114). (SEQ ID NOS: 107 and 108 are non-contiguous sequences from clone Tb38-1F2.) Two open reading frames

the restriction of the superconduction of HERRS from the HERRS II SEQ. ID NO. 75-b, which

- '

ID NO. 105) is a partial clone of TbH-8. The deduced amino acid sequences for these three clones are presented in SEQ ID NOS: 102, 104 and 106.

Further screening of the *M. tuberculosis* genomic DNA library, as described above, resulted in the recovery of ten additional reactive clones, representing seven different genes. One of these genes was identified as the 38 Kd antigen discussed above, one was determined to be identical to the 14Kd alpha crystallin heat shock protein previously shown to be present in *M. tuberculosis*, and a third was determined to be identical to the antigen TbH-8 described above. The determined DNA sequences for the remaining five clones (hereinafter referred to as TbH-29, TbH-30, TbH-32 and TbH-33) are provided in SEQ ID NO: 133-136, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NO: 137-140, respectively. The DNA and amino acid sequences for these antigens were compared with those in the gene bank as described above. No homologies were found to the 5' end of TbH-29 (which contains the reactive open reading frame), although the 3' end of TbH-29 was found to be identical to the *M. tuberculosis* cosmid Y227. TbH-32 and TbH-33 were found to be identical to the previously identified *M. tuberculosis* insertion element IS6110 and to the *M. tuberculosis* cosmid Y50, respectively. No significant homologies to TbH-30 were found.

Positive phagemid from this additional screening were used to infect *E. coli* XL-1 Blue MRF, as described in Sambrook et al., *supra*—Induction of recombinant protein was accomplished by the addition of IPTG—Induced and uninduced lysates were run in duplicate on SDS-PAGF and transferred to nitrocellulose filters—Filters were reacted with human *M. tuberculosis* sera (1:200 dilution) reactive with TbH and a rabbit sera (1:200 or 1:250 dilution) reactive with the N-terminal 4 Kd portion of lacZ. Sera incubations were performed for 2 hours at room temperature. Bound antibody was detected by addition of <sup>12</sup>I-labeled Protein A and subsequent exposure to film for variable times ranging from 16 hours

#### TABLE 2

5	Antigen	Human M. tb <u>Sera</u>	Anti-lacZ <u>Sera</u>
	тьн-20	45 Kd	45 Kd
	тын 30	No reactivity	29 Kd
	ТЬН-32	12 Kd	12 Kd
	TbH-33	16 Kd	16 Kd

10

Positive reaction of the recombinant human M. tuberculosis antigens with both the human M. tuberculosis sera and anti-lacZ sera indicate that reactivity of the human M. tuberculosis sera is directed towards the fusion protein. Antigens reactive with the anti-lacZ sera but not with the human M. tuberculosis sera may be the result of the human M. tuberculosis sera recognizing conformational epitopes, or the antigen-antibody binding kinetics may be such that the 2 hour sera exposure in the immunoblot is not sufficient.

Studies were undertaken to determine whether the antigens TbH-9 and Tb38-1 represent cellular proteins or are secreted into *M. tuberculosis* culture media. In the first study, rabbit sera were raised against A) secretory proteins of *M. tuberculosis*, B) the known secretory recombinant *M. tuberculosis* antigen 85b, C) recombinant Tb38-1 and D) recombinant TbH 9, using protocols substantially as described in Example 3A. Total *M. tuberculosis* lysate, concentrated supernatant of *M. tuberculosis* cultures and the recombinant antigens 85b. TbH-9 and Tb38-1 were resolved on denaturing gels, immobilized on nitrocellulose membranes and duplicate blots were probed using the rabbit sera described above.

The results of this analysis using control serging mol by indication and in the control serging mol by indication and indicati

Fig. 7. The second constraint of the second co

25

residues and would therefore be expected to migrate with a mobility approximately 1 kD larger that the native protein. In Figure 2D, recombinant TbH-9 is lacking approximately 10 kD of the full-length 42 kD antigen, hence the significant difference in the size of the immunoreactive native TbH-9 antigen in the lysate lane (indicated by an arrow). These results demonstrate that Tb38-1 and TbH-9 are intracellular antigens and are not actively secreted by *M. tuberculosis*.

The finding that TbH-9 is an intracellular antigen was confirmed by determining the reactivity of TbH-9-specific human T cell clones to recombinant TbH-9, secretory *M. tuberculosis* proteins and PPD. A TbH-9-specific T cell clone (designated 131TbH-9) was generated from PBMC of a healthy PPD-positive donor. The proliferative response of 131TbH-9 to secretory proteins, recombinant TbH-9 and a control *M. tuberculosis* antigen, TbRa11, was determined by measuring uptake of tritiated thymidine, as described in Example 1. As shown in Figure 3A, the clone 131TbH-9 responds specifically to TbH-9, showing that TbH-9 is not a significant component of *M. tuberculosis* secretory proteins. Figure 3B shows the production of IFN-γ by a second TbH-9-specific T cell clone (designated PPD 800-10) prepared from PBMC from a healthy PPD-positive donor, following stimulation of the T cell clone with secretory proteins, PPD or recombinant TbH-9. These results further confirm that TbH-9 is not secreted by *M. tuberculosis*.

# 20 C USE OF SERA FROM PATIENTS HAVING EXTRAPPT MONARY TUBERCITIOSIS TO IDENTIFY DNA SEQUENCES ENCODING M. 19 BETT 1085 ANTIGENS

Genomic DNA was isolated from *M tuberculosis* Frdman strain, randomly sheared and used to construct an expression library employing the Lambda ZAP expression system (Stratagene, La Jolla, CA). The resulting library was screened using pools of sera obtained from individuals with extrapulmonary tuberculosis. In the arithmetic of the contraction of the co

on the first of the first that a street connectation adversed to the

<sup>50</sup> March N. C. C. N. P. and N.P. S. white contributions only similarly to known a finite of the

5

153, respectively, with the 5' and 3' DNA sequences for XP32 being provided in SEQ ID NOS: 154 and 155, respectively. The predicted amino acid sequence for XP14 is provided in SEQ ID NO: 156. The reverse complement of XP14 was found to encode the amino acid sequence provided in SEQ ID NO: 157.

Comparison of the sequences for the remaining 14 clones (hereinafter referred to as XP1-XP6, XP17-XP19, XP22, XP25, XP27, XP30 and XP36) with those in the genebank as described above, revealed no homologies with the exception of the 3' ends of XP2 and XP6 which were found to bear some homology to known M. tuberculosis cosmids. The DNA sequences for XP27 and XP36 are shown in SEQ ID NOS: 158 and 159, respectively, with the 5' sequences for XP4, XP5, XP17 and XP30 being shown in SEO ID NOS: 160-163, respectively, and the 5' and 3' sequences for XP2, XP3, XP6, XP18, XP19, XP22 and XP25 being shown in SEQ ID NOS: 164 and 165; 166 and 167, 168 and 169; 170 and 171; 172 and 173; 174 and 175; and 176 and 177, respectively. XP1 was found to overlap with the DNA sequences for TbH4, disclosed above. The full-length DNA sequence for TbH4-XP1 is plovided in SEQ ID NO: 178. This DNA sequence was found to contain an open reading frame encoding the amino acid sequence shown in SEQ ID NO: 179. The reverse complement of TbH4-XP1 was found to contain an open reading frame encoding the amino acid sequence shown in SEQ ID NO: 180. The DNA sequence for XP36 was found to contain two open reading frames encoding the amino acid sequence shown in SEQ ID NOS: 181 and 182, with the reverse complement containing an open reading frame encoding the amino acid sequence shown in SEQ ID NO. 183

Recombinant XP1 protein was prepared as described above in Example 3B, with a metal ion affinity chromatography column being employed for purification. Recombinant XP1 was found to stimulate cell proliferation and IFN- $\gamma$  production in T cells isolated from an M tuberculosis immune donors.

More than the first of the first of the second state of the second state of the second second

serological activity with a serum pool from *M. tuberculosis*-infected patients which showed little or no immunoreactivity with other antigens of the present invention. Rabbit anti-sera was generated against the most reactive fraction using the method described in Example 3A. The anti-sera was used to screen an *M. tuberculosis* Erdman strain genomic DNA expression library prepared as described above. Bacteriophage plaques expressing immunoreactive antigens were purified. Phagemid from the plaques was rescued and the nucleotide sequences of the *M. tuberculosis* clones determined.

Ten different clones were purified. Of these, one was found to be TbRa35, described above, and one was found to be the previously identified *M. tuberculosis* antigen, HSP60. Of the remaining eight clones, six (hereinafter referred to as RDIF2, RDIF5, RDIF8, RDIF10, RDIF11 and RDIF12) were found to bear some similarity to previously identified *M. tuberculosis* sequences. The determined DNA sequences for RDIF2, RDIF5, RDIF8, RDIF10 and RDIF11 are provided in SEQ ID NOS: 184-188, respectively, with the corresponding predicted amino acid sequences being provided in SEQ ID NOS: 189-193, respectively. The 5° and 3° DNA sequences for RDIF12 are provided in SEQ ID NOS: 194 and 195, respectively. No significant homologies were found to the antigen RDIF-7. The determined DNA and predicted amino acid sequences for RDIF7 are provided in SEQ ID NOS: 196 and 197, respectively. One additional clone, referred to as RDIF6 was isolated, however, this was found to be identical to RDIF5.

Recombinant RDIF6, RDIF8, RDIF10 and RDIF11 were prepared as described above. These antigens were found to stimulate cell proliteration and IFN-y production in T cells isolated from *M\_tuber culosis*-immune donors.

20

25

1 XAMPLE 4

And Market and the straining to the state of their dimensional parties protein and their graphs (PDD) in these

WO 98/16645 PCT/US97/18214

PPD was prepared as published with some modification (Seibert, F. et al., Tuberculin purified protein derivative. Preparation and analyses of a large quantity for standard. The American Review of Tuberculosis 44:9-25, 1941). *M. tuberculosis* Rv strain was grown for 6 weeks in synthetic medium in roller bottles at 37°C. Bottles containing the bacterial growth were then heated to 100°C in water vapor for 3 hours. Cultures were sterile filtered using a 0.22 μ filter and the liquid phase was concentrated 20 times using a 3 kD cutoff membrane. Proteins were precipitated once with 50% ammonium sulfate solution and eight times with 25% ammonium sulfate solution. The resulting proteins (PPD) were fractionated by reverse phase liquid chromatography (RP-HPLC) using a C18 column (7.8 x 300 mM; Waters, Milford, MA) in a Biocad HPLC system (Perseptive Biosystems, Framingham, MA). Fractions were eluted from the column with a linear gradient from 0-100% buffer (0.1% TFA in acetonitrile). The flow rate was 10 ml/minute and eluent was monitored at 214 nm and 280 nm.

Six fractions were collected, dried, suspended in PBS and tested individually in *M. tuberculosis*-infected guinea pigs for induction of delayed type hypersensitivity (DTH) reaction. One fraction was found to induce a strong DTH reaction and was subsequently fractionated further by RP-HPLC on a microbore Vydac C18 column (Cat. No. 218TP5115) in a Perkin Flmer/Applied Biosystems Division Model 172 HPLC. Fractions were eluted with a linear gradient from 5-100% buffer (0.05% TFA in acctonitrile) with a flow rate of 80 ml/minute. Fluent was monitored at 215 mm. Fight fractions were collected and tested for induction of DTH in *M. tuberculosis*-infected guinea pigs. One fraction was found to induce strong DTH of about 16 mm induration. The other fractions did not induce detectable DTH. The positive fraction was submitted to SDS-PAGE gel electrophoresis and found to contain a single protein band of approximately 12 kD molecular weight.

25 This polypeptide, herein after referred to as DPPD, was sequenced from the

and the state of the state of the source of the state of

#### **EXAMPLE 5**

# SYNTHESIS OF SYNTHETIC POLYPEPTIDES

Polypeptides may be synthesized on a Millipore 9050 peptide synthesizer using FMOC chemistry with HPTU (O-Benzotriazole N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to clute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray mass spectrometry and by amino acid analysis.

This procedure was used to synthesize a TbM-1 peptide that contains one and a half repeats of a TbM-1 sequence. The TbM-1 peptide has the sequence GCGDRSGGNLDQIRLRRDRSGGNL (SLQ ID NO: 63).

20

15

5

## EXAMPLE 6

## USE OF REPRESENTATIVE ANTIGENS FOR SERODIAGNOSIS OF TUBERCULOSIS

25 This Example illustrates the diagnostic properties of several representative

i.e. the pare contents were than removed and the wells were blocked for a

times with PBS/0.1% Tween 20<sup>TM</sup>. 50 μL sera, diluted 1:100 in PBS/0.1% Tween 20<sup>TM</sup>/0.1% BSA, was then added to each well and incubated for 30 minutes at room temperature. The plates were then washed again five times with PBS/0.1% Tween 20<sup>TM</sup>.

The enzyme conjugate (horseradish peroxidase - Protein A, Zymed, San Francisco, CA) was then diluted 1:10,000 in PBS/0.1% Tween 20<sup>τM</sup>/0.1% BSA, and 50 μL of the diluted conjugate was added to each well and incubated for 30 minutes at room temperature. Following incubation, the wells were washed five times with PBS/0.1% Tween 20<sup>τM</sup>. 100 μL of tetramethylbenzidine peroxidase (TMB) substrate (Kirkegaard and Perry Laboratories, Gaithersburg, MD) was added, undiluted, and incubated for about 15 minutes. The reaction was stopped with the addition of 100 μL of 1 N H<sub>2</sub>SO<sub>4</sub> to each well, and the plates were read at 450 nm.

Figure 4 shows the ELISA reactivity of two recombinant antigens isolated using method A in Example 3 (TbRa3 and TbRa9) with sera from *M. tuberculosis* positive and negative patients. The reactivity of these antigens is compared to that of bacterial lysate isolated from *M. tuberculosis* strain H37Ra (Difco, Detroit, MI). In both cases, the recombinant antigens differentiated positive from negative sera. Based on cut-off values obtained from receiver-operator curves, TbRa3 detected 56 out of 87 positive sera, and TbRa9 detected 111 out of 165 positive sera.

Figure 5 illustrates the FHSA reactivity of representative antigens isolated using method B of Example 3. The reactivity of the recombinant antigens TbH4, TbH12. Tb38-1 and the peptide TbM 1 (as described in Example 4) is compared to that of the 38 kD antigen described by Andersen and Hansen, *Infect Immun*, 57:2481-2488, 1989. Again, all of the polypeptides tested differentiated positive from negative sera. Based on cut-off values obtained from receiver-operator curves. TbH4 detected 67 out of 126 positive sera, TbH12 detected 50 out of 125 positive sera, 38-1 detected 61 out of 101 positive sera and the 1bM 1 and 1 14 and 157.

the copies (86 pt) in the compared of the contract and compared

5

to the reactivity of *M. tuberculosis* lysate and the 38 kD antigen. The results are presented in Table 3, below:

TABLE 3

REACTIVITY OF ANTIGENS WITH SERA FROM M. TUBERCULOSIS PATIENTS

	Acid Fast		ELISA Values					
Patient	Sputum	Lysate	38kD	TbRa9	TbH12	ТЪН4	TbRa3	
Tb01B93I-2	++++	1.853	0.634	0.998	1.022	1.030	1.314	
Tb01B93I-19	++++	2.657	2.322	0.608	0.837	1.857	2.335	
Tb01B93I-8	++-	2.703	0.527	0.492	0.281	0.501	2.002	
Tb01B93I-10	1-+-+	1.665	1.301	0.685	0.216	0.448	0.458	
Tb01B93I-11	+++	2.817	0.697	0.509	0.301	0.173	2.608	
Tb01B93I-15	+++	1.28	0.283	0.808	0.218	1.537	0.811	
Tb01B93I-16	+ • • •	2.908	<b>≯</b> 3	0.899	0.441	0.593	1.080	
Tb01B93I-25	+-++	0.395	0.131	0.335	0.211	0.107	0.948	
Tb01B93I-87	1	2.653	2.432	2.282	0.977	1.221	0.857	
Tb01B93I-89		1.912	2.370	2.436	0.876	0.520	0.952	
Tb01B94I-108	+••	1.639	0.341	0.797	0.368	0.654	0.798	
Tb01B94I-201	+- + +	1.721	0.419	0,661	0.137	0.064	0.692	
Tb01B93I-88		1,939	1.269	2.519	1.381	0.214	0.530	
Thotpost.or	ļ		!		· ·	!	ļ	

			x + x	111	, qu.	1
fiedBodi (2.4 )	· · · · · · · · · · · · · · · · · · ·	er Con	0.251	, c <sub>1</sub> ,	· Foon	m '5

	Acid Fast		ELISA Values					
Patient	Sputum	Lysate	38kD	TbRa9	TbH12	ТЬН4	TbRa3	
Ть01В931-9	+	2.649	0.278	0.210	0.140	0.181	1.586	
Tb01B93I-14	+	>3	1.538	0.282	0.291	0.549	2.880	
Tb01B93I-21	+	2.645	0.739	2.499	0.783	0.536	1.770	
Tb01B93I-22	+	0.714	0.451	2.082	0.285	0.269	1.159	
Tb01B93I-31	-+	0.956	0.490	1.019	0.812	0.176	1.293	
Tb01B931-32	-	2.261	0.786	0.668	0.273	0.535	0.405	
Tb01B93I-52	_	0.658	0.114	0.434	0.330	0.273	1.140	
Tb01B93I-99		2.118	0.584	1.62	0.119	0.977	0.729	
Tb01B94I-130	-	1.349	0.224	0.86	0.282	0.383	2.146	
Tb01B94I-131		0.685	0.324	1.173	0.059	0.118	1.431	
AT4-0070	Normal	0.072	0.043	0.092	0.071	0.040	0.039	
AT4-0105	Normal	0.397	0.121	0.118	0.103	0.078	0.390	
3/15/94-1	Normal	0.227	0.064	0.098	0.026	0.001	0.228	
4 15 93-2	Normal	0.114	0.240	0.071	0.034	0.041	0.264	
5 26 94-4	Normal	0.089	0.259	0.096	0.046	0.008	0.053	
5/26/94-3	Normal	0.139	0.093	0.085	0.019	0.067	0,01	

Based on cut-off values obtained from receiver-operator curves. TbRa3 detected 23 out of 27 positive sera. TbRa9 detected 22 out a 22 out

The reactivity of the recombinant antigen TbRa11 with sera from M. tuberculosis patients shown to be negative for the 38 kD antigen, as well as with sera from PPD positive and normal donors, was determined by ELISA as described above. The results are shown in Figure 6 which indicates that TbRa11, while being negative with sera from PPD positive and normal donors, detected sera that were negative with the 38 kD antigen. Of the thirteen 38 kD negative sera tested, nine were positive with TbRa11, indicating that this antigen may be reacting with a sub-group of 38 kD antigen negative sera. In contrast, in a group of 38 kD positive sera where TbRa11 was reactive, the mean OD 450 for TbRa11 was lower than that for the 38 kD antigen. The data indicate an inverse relationship between the presence of TbRa11 activity and 38 kD positivity.

10

15

25

The antigen TbRa2A was tested in an indirect ELISA using initially 50 µl of serum at 1:100 dilution for 30 minutes at room temperature followed by washing in PBS Tween and incubating for 30 minutes with biotinylated Protein A (Zymed, San Francisco, CA) at a 1:10,000 dilution. Following washing, 50 µl of streptavidin-horseradish peroxidase (Zymed) at 1:10,000 dilution was added and the mixture incubated for 30 minutes. After washing, the assay was developed with TMB substrate as described above. The reactivity of TbRa2A with sera from *M. tuberculosis* patients and normal donors in shown in Table 4. The mean value for reactivity of TbRa2A with sera from *M. tuberculosis* patients was 0.444 with a standard deviation of 0.309. The mean for reactivity with sera from normal donors was 0.109 with a standard deviation of 0.029. Testing of 38 kD negative sera (Figure 7) also indicated that the 1bRa2A antigen was capable of detecting sera in this category.

TABLE 4

REACTIVITY OF IBRAZA WITH SERVEROM M. TO BUR TO COSE PARIENTS AND FROM NORMAL DONORS.

158		. B	· Star	
1588		113	1, 1	1
	٠			- 1

Ть91	TB	0.393
Tb92	TB	0.401
Ть93	TB	0.232
Ть94	ТВ	0.333
Ть95	TB	0.435
Tb96	TB	0.284
Tb97	TB	0.320
Ть99	ТВ	0.328
Tb100	ТВ	0.817
Tb101	TB	0.607
Tb102	TB	0.191
Tb103	ТВ	0.228
Tb107	ТВ	0.324
Tb109	ТВ	1.572
Tb112	TB	0.338
DL4-0176	Normal	0.036
AT4-0043	Normal	0.126
AT4-0044	Normal	0.130
AT4-0052	Normal	0.135
ΑΤ4-0053	Normal	0.133
AT4-0062	Normal	0.128
AT4-0070	Normal	0.088
AT4-0091	Normal	0.108
Α Γ4-0100	Normal	0.106
AT4-0105	Normal	0.108
AT4-0109	Normal	0.105

The reactivity of the recombinant antigen (g) (SEQ ID NO: 60) with sera from *M tuberculosis* patients and normal donors was determined by ELISA as described above. Figure 8 shows the results of the titration of antigen (g) with four *M tuberculosis* positive sera that were all reactive with the 38 kD antigen and with four donor sera. All four positive sera were reactive with antigen (g).

The reactivity of the recombinant antigen TbH-29 (SEQ ID NO: 137) with

Silver of the second of LISA represents inversing markets of the second o

donors and with a pool of sera from *M. tuberculosis* patients. The mean OD 450 was demonstrated to be higher with sera from *M. tuberculosis* patients than from normal donors, with the mean OD 450 being significantly higher in the indirect ELISA than in the direct ELISA. Figure 11 is a titration curve for the reactivity of recombinant TbH-33 with sera from *M. tuberculosis* patients and from normal donors showing an increase in OD 450 with increasing concentration of antigen.

The reactivity of the recombinant antigens RDIF6, RDIF8 and RDIF10 (SEQ ID NOS: 184-187, respectively) with sera from *M. tuberculosis* patients and normal donors was determined by ELISA as described above. RDIF6 detected 6 out of 32 *M. tuberculosis* sera and 0 out of 15 normal sera; RDIF8 detected 14 out of 32 *M. tuberculosis* sera and 0 out of 15 normal sera; and RDIF10 detected 4 out of 27 *M. tuberculosis* sera and 1 out of 15 normal sera. In addition, RDIF10 was found to detect 0 out of 5 sera from PPD-positive donors.

#### 15

#### EXAMPLE 7

#### PREPARATION AND CHARACTERIZATION OF M. TUBERCULOSIS FUSION PROTEINS

A fusion protein containing TbRa3, the 38 kD antigen and Tb38-1 was prepared as follows.

20 Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR in order to facilitate their fusion and the subsequent expression of the fusion protein TbRa3-38 kD-Tb38-1. TbRa3, 38 kD and Tb38-1 DNA was used to perform PCR using the primers PDM-64 and PDM-65 (SEQ ID NO: 141 and 142), PDM-57 and PDM-58 (SEQ ID NO: 143 and 144), and PDM-69 and PDM-60 (SEQ ID NO: 145-146), respectively. In each case, the DNA amplification was performed using 10 μl 10X Pfu buffer, 2 μl 10 mM dNTPs, 2 μl each of the PCR primers at 13 M and 143.

Sold of the Color of George P. Color of the English Life for English 1884.00 cm dental properties of the Color of the Colo

68°C for 15 sec and 72°C for 3 min, and finally by 72°C for 4 min. For Tb38-1 denaturation at 94°C for 2 min was followed by 10 cycles of 96°C for 15 sec, 68°C for 15 sec and 72°C for 1.5 min, 30 cycles of 96°C for 15 sec, 64°C for 15 sec and 72°C for 1.5, and finally by 72°C for 4 min.

The TbRa3 PCR fragment was digested with Ndcl and EcoRI and cloned directly into pT7^L2 II. 1 vector using Ndcl and EcoRI sites. The 38 kD PCR fragment was digested with Sse8387l, treated with T4 DNA polymerase to make blunt ends and then digested with EcoRI for direct cloning into the pT7^L2Ra3-1 vector which was digested with Stul and EcoRI. The 38-1 PCR fragment was digested with Eco47III and EcoRI and directly subcloned into pT7^L2Ra3/38kD-17 digested with the same enzymes. The whole fusion was then transferred to pET28b using NdcI and EcoRI sites. The fusion construct was confirmed by DNA sequencing.

5

10

15

20

The expression construct was transformed to BLR pLys S *E. coli* (Novagen, Madison, WI) and grown overnight in LB broth with kanamycin (30 μg/ml) and chloramphenicol (34 μg/ml). This culture (12 ml) was used to inoculate 500 ml 2XYT with the same antibiotics and the culture was induced with IPTG at an OD560 of 0.44 to a final concentration of 1.2 mM. Four hours post-induction, the bacteria were harvested and sonicated in 20 mM. Tris (8.0), 100 mM NaCl, 0.1% DOC, 20 μg/ml Leupeptin, 20 mM. PMSF followed by centrifugation at 26,000 X g. The resulting pellet was resuspended in 8 M urea, 20 mM. Tris (8.0), 100 mM. NaCl and bound to Pro-bond nickel resin (Invitrogen, Carlsbad, CA). The column was washed several times with the above buffer then eluted with an imidazole gradient (50 mM, 100 mM, 500 mM imidazole was added to 8 M urea, 20 mM. Tris (8.0), 100 mM NaCl). The cluates containing the protein of interest were then dialzyed against 10 mM. Tris (8.0).

The DNA and amino acid sequences for the resulting fusion protein

5

10

15

20

procedure to that described above. The DNA sequence for the TbH9-Tb38-1 fusion protein is provided in SEQ ID NO: 151.

A fusion protein containing TbRa3, the antigen 38kD, Tb38-1 and DPEP was prepared as follows.

Each of the DNA constructs TbRa3, 38 kD and Tb38-1 were modified by PCR and cloned into vectors essentially as described above, with the primers PDM-69 (SEQ ID NO:145 and PDM-83 (SEQ ID NO: 200) being used for amplification of the Tb38-1A fragment. Tb38-1A differs from Tb38-1 by a Dral site at the 3' end of the coding region that keeps the final amino acid intact while creating a blunt restriction site that is in frame. The TbRa3/38kD/Tb38-1A fusion was then transferred to pET28b using Ndel and EcoR1 sites.

DPEP DNA was used to perform PCR using the primers PDM-84 and PDM-85 (SEQ ID NO: 201 and 202, respectively) and 1 μl DNA at 50 ng/μl. Denaturation at 94 °C was performed for 2 min, followed by 10 cycles of 96 °C for 15 sec, 68 °C for 15 sec and 72 °C for 1.5 min; 30 cycles of 96 °C for 15 sec, 64 °C for 15 sec and 72 °C for 1.5 min; and finally by 72 °C for 4 min. The DPEP PCR fragment was digested with EcoRI and Eco72I and clones directly into the pET28Ra3/38kD/38-1A construct which was digested with DraI and EcoRI. The fusion construct was confirmed to be correct by DNA sequencing. Recombinant protein was prepared as described above. The DNA and amino acid sequences for the resulting fusion protein (hereinafter referred to as TbF-2) are provided in SEQ ID NO: 203 and 204, respectively.

#### EXAMPLE 8

USE OF <u>M. Tuberculosis Eusion Proteins for</u> Sprodiagnosis of Tuberculosis

25

The effectiveness of the fusion protein TbRa3-38 kD-Tb38-1, prepared as described above, in the serodiagnosis of tuberculosis infection was examined by ELISA.

The LEIS V protocol was as described above in Example 6, with the fusion

10

the three antigens individually or in combination. Such a panel enabled the dissection of the serological reactivity of the fusion protein to determine if all three epitopes functioned with the fusion protein. As shown in Table 5, all four sera that reacted with TbRa3 only were detectable with the fusion protein. Three sera that reacted only with Tb38-1 were also detectable, as were two sear that reacted with 38 kD alone. The remaining 15 sera were all positive with the fusion protein based on a cut-off in the assay of mean negatives +3 standard deviations. This data demonstrates the functional activity of all three epitopes in the fusion protein.

Table 5

<u>Reactivity of Tri-Peptide Fusion Protein with Sera from M. Tuberculosis Patients</u>

Serum ID	Status	ELISA	and/or West	Fusion	Fusion	
		Reactivity	with Individ	ual proteins	recombinant	Recombinant
		38kd	Tb38-1	TbRa3	OD 450	Status
01B93I-40	TB	-	_	4-	0.413	• .+
01B93I-41	ТВ		+	+	0.392	+
01B93I-29	ТВ	-+	-	+	2.217	+
01B93I-109	TB	+	+	+	0.522	<del>+</del> .
01B93I-132	TB	+	+	ł	0.937	+
5004	ТВ	1	ł	::	1.098	
15004	TB	-+-	+	-1	2.077	+
39004	TB	-	+	•	1.675	+
68004	ТВ	•	1		2.388	
99004	ТВ			:	0.607	. 3010 4-0-
107004	TB	-		:	0.667	,
92004	ТВ	·			1.070	
97004	TB		;   -	1	1.152	•
118004	TB			ı	2.694	,
173004	ТВ		;		3.258	+
175004	TB			,	2.514	•
271001	113	*	t	•		

289004	ТВ	-	_	+	0.848	+
308004	TB	-	+	-	3.338	+
314004	TB	_	+	-	1.362	+
317004	ТВ	+	-	-	0.763	+
312004	TB	-	-	- →	1.079	+
D176	PPD	-		-	0.145	•
D162	PPD	-	-	-	0.073	
D161	PPD	-		-	0.097	
D27	PPD	-	-	-	0.082	-
A6-124	NORMAL	-		-	0.053	-
A6-125	NORMAL	-			0.087	_
A6-126	NORMAL	-	-	-	0.346	+ .
A6-127	NORMAL	- :	-	-	0.064	-
A6-128	NORMAL	<u>-</u>	-	-	0.034	-
A6-129	NORMAL	-	-	-	0.037	-
A6-130	NORMAL	-	-	-	0.057	-
A6-131	NORMAL	-	-	-	0.054	-
A6-132	NORMAL	-	-		0.022	-
A6-133	NORMAL	-	-		0.147	-
Δ6-134	NORMAL	-	<u>-</u>	-	0.101	-
A6-135	NORMAL		<u>-</u>		0.066	-
Δ6-136	NORMAL	-	-		0.054	-
A6-137	NORMAL		-	-	0.065	-
Δ6-138	NORMAL				0.041	-
A6-139	NORMAL	-			0.103	
A6-140	NORMAL	-	-	-	0.212	-
.\6-141	NORMAL		-		0.056	-
.16-142	NORMAL			-	0.051	

The reactivity of the fusion protein Tb1-2 with sera from *M\_tuberculosis*-infected patients was examined by FLISA using the protocol described above. The results of these studies (Lable to demonstrate that all the national time to account and the following studies).

 $\label{thm:table 6} \mbox{Reactivity of TbF-2 Fusion Protein with TB and Normal Sera}$ 

Serum ID	Status	TbF OD450	Status	TbF-2 OD450	Status	ELISA Reactivity			<del></del>
· · · · · · · · · · · · · · · · · · ·	<del>                                     </del>	1	· · · · · ·		<del> </del> -	38 kD	TbRa3	Tb38-1	DPEP
B931-40	TB	0.57	+	0.321	it	-	+	-	+
B931-41	TB	0.601	+	0.396	+	+	+	+	-
B931-109	TB	0.494	4	0.404	-4		+	±	†
B931-132	TB	1.502	+	1.292	+	1	+	+	<u>+</u>
5004	TB	1.806	+	1.666	-+	±	+	+	-
15004	TB	2.862	+	2.468	+	+	+	+	-
39004	TB	2.443	+	1.722		4	+	+ -	i -
68004	TB	2.871	+	2.575	+	+	+	+	<del> </del> -
99004	TB	0.691	+	0.971	+	†-·	<u>.</u>	+	
107004	TB	0.875	+	0.732	+	1 -	±.	+	<del>  -</del>
92004	TB	1.632	ŧ	1 394	-4	1 4	<u>+</u> :	±	-
97004	ТВ	1.491	+	1 979	-+	+	<u>±</u>	-	+
118004	TB	3.182	+	3.045	i	-+	±	-	-
173004	TB	3.644	+	3.578	+	-+	+	+	-
175004	ТВ	3.332	+	2.916	+	1	+	-	-
274004	TB	3.696	+	3.716	+	-	1	-	+
276004	TB	3.243	+	2.56	+	-	-	+	-
282004	TB	1.249	+	1.234	+	-+	-	† <u>-</u>	-
289004	TB	1.373	+	1.17	i	-	+	-	-
308004	TB	3.708	•1	3.355	+	-	-	+	-
314004	ТВ	1.663		1.399	±	-		4	-
317004	TB	1.163	+	0.92	+	+	-	-	-
312004	TB	1.709	+	1.453	+	-	+	-	-
380004	ТВ	0.238	-	0.461		-	+	-	
451004	TB	0.18	-	02		-	-	-	+
478004	TB	0.188	-	0.469	i	-	-	-	,
410004	TB	0.384	+	2.392	•	<u>.</u> +	-	-	
411004	TB	0.306		0.874		-	i .	-	
421004	TB	0.357		1.456		-		-	
528004	TB	0.047		0.196		1	-	-	· · ·
.16-87	Normal	0.094		0.063		1 -	-	-	
16-88	Normal	0.214		0 19		-	-		-
A6-89	Normal	0.248	- 1	0.125		-		-	
A6-90	Normal	0.179	-	0.206			0 . 000	- 1	-
A6-91	Normal	0.135	-	0.151	· ×		-	-1	
A6-92		0.064		0.097			-		
A6 93	Normal	10.072		0.098			<del> </del>	· · · · ·	
		0 072		0.064	·				
A6 95	· Sorma	· (1)		0.150		<del>-</del>	1 1		0

5

One of skill in the art will appreciate that the order of the individual antigens within the fusion protein may be changed and that comparable activity would be expected provided each of the epitopes is still functionally available. In addition, truncated forms of the proteins containing active epitopes may be used in the construction of fusion proteins.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

#### SEQUENCE LISTING

#### (1) GENERAL INFORMATION:

- (i) APPLICANTS: Reed, Steven G. Skeiky, Yasir A.W. Dillon, Davin C. Campos-Neto, Antonia Houghton, Raymond Vedvick, Thomas S. Twardzik, Daniel R. Lodes, Michael J.
- (ii) TITLE OF INVENTION: COMPOUNDS AND METHODS FOR DIAGNOSIS OF TUPERCULOCIC
- (111) NUMBER OF SEQUENCES: 209
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: SEED and BERRY LLP
  - (B) STREET: 6300 Columbia Center, 201 Fifth Avenue
  - (C) CITY: Seattle
  - (D) STATE: Washington
  - (E) COUNTRY: USA
  - (F) ZIF: 98104-7092
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (T) OPERATING SYSTEM: PC-DOSTMS-DOS
  - (b) SOFTWARE: Patientin Release #1.0, Versial #1.30
- (VI) CURPENT APPLICATION DATA:
  - PA APELICATION NUMBER:
  - TE FILLES MATE: SI-CT-10-
  - (C CLASSIFICATION:
- WALL AND PMIT AGENT THE EMALIGNE
  - " MAME: Mass, Lavis ".
  - PORED TO BANCON MUMBERS (F.)

  - Strong and the settle of the American American Section 1997
     Strong and American Section 1997
  - 2. Thus 1. The second second

#### (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

60	GAAGAACACA	CAAACGAACG	AATCGACGGG	CCAAACGCAC	GTAGTTTGAA	CGAGGCACCG
1.20	AATCGGCGCC	COGCGGCTGC	GGTCTGACCG	CATCGCCGCA	TGGTGAAATC	ACCATGAAGA
180	GATGCAGCCG	TCGTATACCA	GGCGGCCCGG	CATCATGGCT	GTGTGACTTC	GCTGCGGCCG
240	CCCGACCGCC	CCCCTGACGT	CCGGCATCCG	GCCGTTGGAC	GCGCGCCACT	GTCGTCTTCG
300	TGCGAACAAG	ACSTGTCGTT	GCCGATCCCA	CAACAGCCTC	CCAGUCTGCT	GOCCAGTIGA
360	CCACAAGCTG	GCATCGCCGA	ACCGAGGCGC	ÇATCGGGGGC	TCGAGGGGG	GGCAGTCTGG
420	CATCCAGGCG	GCUTGACGAA	CTGTCGTTCA	GGATCTGCCG	CCBABCACGG	AAGAA33CCG
490	CTCGTCGCCS	GTÉCGAAGGT	TCCGTCTCGG	CGCCGACGTT	GTTCGGCCAC	GCGGCCGCCĞ
540	CGCATCGGCG	TGCTGTCACG	GGCGGCTGGA	UGTGAATCAA	ACCTCACGTT	GTCACCCAGA
600	CCGCTGTTCA	CGGNTTCAGC	ATTGGGGGGC	AGGNAACTG	TGCAGGCCGC	ATGGAGTTGC
660	GCACGGTGCG	CGCGCGTGTA	GTCGAACACT	AUGUGTOCAT	CCGCCTGGTG	SCTACGCCCC
2.75	GTGGTGNCTC	TOGAGATAG	GAAAGCCGTC	ACCGCCCGGT	GGNCGCACGC	CTNTCCCCAC
7 est,		ONATOA	CALASMA SIA	шиговисти	AMCACCCÓCD	CHUACCACING

TO REPORMATION FOR SECTION NOTE:

CARLEST CHARACTERISTICS:

(A) LEMGIE: '(a) base pairs

(B) TYLE: medicing of the strain of the strai

STANDS OF THE SECTION OF SECTION

The things of the obtained to see the probability of the probability of  $\chi_{\rm poly} = \chi_{\rm poly} = \chi_{\rm poly}$ 

WO 98/16645 PCT/US97/18214

60

TTTCTCGACG	ACSTGACOST	GAGCCGTCGC	CATGCTGAAT	TCCCCTTGGA	AAACAACGAA	300
TTCAATGTCG	TCGATGTCGG	GAGTOTCAAO	GGCACCTACC	TCAACCGCGA	GCCCGTGGAT	360
TOGGOGGTGC	TGGCGAACGG	CGACGAGGTC	CAGATCGGCA	AGCTCCGGTT	GGTGTTCTTG	420
ACCGGACCCA	AGCAAGGCGA	GGATGACGGG	AGTACCGGGG	CCCCGTGAGC	GCACCCGATA	480
GCCCCGCGCT	GGCCGGGATG	TCGATCGGGG	CGGTCCTCCG	ACCTGCTACG	ACGGGATTTT	540
CCCTGATGTC	CACCATCTCC	AAGATTOGAT	TCTTGGGAGG	CTTGAGGGTC	NGGGTGACCC	600
agraegeaea	CTCATTCNGG	GGTNTCGGCN	SCTTTCACCC	CNTACCNACT	GCCNCCIGGN	660
TTGCNAATTC	NTTCTTCMAT	GCCCNNAAAG	GGACCHTTAN	CTTGCCGCIN	GAAANGGTNA	72C
TCUNGGGCCC	NTCCTNGAAN	CCCCNTCCCC	CT			7 <b>5</b> 2
(2) INFORMA	TION FOR SE	Q ID NO:3:				• *

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 813 base pairs

(B) TYPF: nucleic acid

(C) STRAMDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE LEGGRIPTION: SEQ 10 No: 1:

## PCT/US97/18214

61

GAGCAACGCA GACCGGGACA ACWGCTATCG ATAGCCGCCN AATGCCGGCT TGGAACCCNG 780

TGAAATTATC ACAACTTCGC AGTCACNAAA NAA						
(2) INFORMATION FOR SEQ ID NO:4:						
(i) SEQUENCE CHARACTERISTICS:  (A) DENGTH: 447 base pairs  (B) TYFE: nucleic acid  (C) OTFANDEDNESS: single  (D) TOFGLOGY: linear						
•						
(M1) SEQUENCE DESCRIPTION: DEC 10 NO.4:						
CGCTATGAAT ACGCCUGCGI CTGATAACTT CCAGCTGTTV CAGGGTGGGC AGGGATTCGC	€0					
CATICCEATE GGGCAGGGA TGGCGATGGC COGCGAGATC CGATEGGGTG GGGGGTCAGC	12:0					
CACCGTTCAT ATCGCCCCTA CCCCCCTTCCT CGCCTTGGGT GTTGTCCACA ACAACGGCAA	180					
CGGGGCAGGA GTCCAATGCG TGGTTGGGGAG OFTTGGGGGG GCAAGTCTCG GCATCTCCAC	240					
CGGCGACGTG ATCACCGCGG TCGAEGGGCC TCCGATCAAC TCGGCCACCC CCATGCCGGA	300					
CONSTITATO COCCATORATO DE GALORA EL CATOTOGA AACEGORAA OCAAGETOGO	36,0					
CGGCAGGCET ACAGGGAAGE THA TATTE E GSAGEGACOC OCE FOITSAT STOGTGGYGG	420					
ATAPATUT - POOLET AATTOGA	4.4.					
DE THE RESTOR FOR DRY IN NOTE.						
F. TED TE MEARA TEER ENTITY :  A. LEM STEER OF EACH :  BEREE OF LEAST :  CONTAINED TO EACH :  CONTAINED TO EAC						

dan també e gara

WO 98/16645 PCT/US97/18214

62

CCGGCGACGG	NGAGCGCCGG	AATGGCGCGA	GTCACCIACCT	CONCACTUAT	GCCCACNCTG	240
ATCCAATCAA	CCTGNATTCG	GNCTGNGGGN	CCATTTGACA	ATCGAGGTAG	TGAGCGCAAA	300
TGAATGATGG	AAAACGGGNG	GNGACGTCCG	NTGTTCTGGT	GGTGNTAGGT	GNCTGNCTGG	360
NGTNGNGGNT	ATCAGGATGT	TCTTCGNCGA	AANCTGATGN	CGAGGAACAG	GGTGTNCCCG	420
иидиисснаи	GGNGTCCNAN	CCCNNNNTCC	TCGNCGANAT	CANANAGNOS	NTTGATGNGA	480
NAAAAGGGTG	GANCAGINNN	AANTNGNGGN	CCHAANAANC	риисииаиии	NNAGNTNGNT	540
CHATTATANG	ОТИМИМИМА	nngnngnnun	NNN AANCNN	AAAGUUUUTU	NNGGNTTNTT	600
TAAH						604

#### 12) INFORMATION FOR SEQ ID NO: 0:

- (:) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 633 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

### (x:) DEQUENCE DESCRIPTION: SEC IP NO:6:

WO 98/16645 PCT/US97/18214

63

(A) LENGTH: 1362 base pairs(B) TYPE: nucleic acid(C) STRANDEDNESS: single(D) TOPOLOGY: linear

#### (xi) SLQUENCE DESCRIPTION: SEQ ID NO:7:

CGACGACGAC	GACGCCGCAG	AGCGGGGGGCG	AACGGCGATC	GACGCGGCCC	TGGCCAGAGT	60
CGGUACCACC	CAGGAGGGAS	TOGAATCATO	AAATTTGTCA	ACCATATTGA	GCCCGTCGCG	120
CCCCGCCGAG	condendate	GOTTEC IGAÇ	STOTATOCCO	Masadada i	GSAGTTCGGC	180
cqueraceca	AGCCGCTCGC	CATGOTHOO	CCCCACGAGG	GACTICTCAC	CCCCGGCTGG	240
GC BACGTTOC	GCGAGACACI	aumaamaaac	CASSTGCCSC	GTGGCCGCAA	GGAAGCCGTC	300
GCCGCCGCCG	rogoggodag	CCTGCGCTGC	CHCTGGTGCG	TCGA::GCACA	CACCACCATG	360
CTGTACGCGG	CAGGCCAAAC	CGACACCGCC	GCGGCGATCT	TGGCCGGCAC	AGCACCTGCC	420
accestaacc	CGAACGCGCC	STATGTGGCG	TSGGCGGCAG	GAACHGGGAH	ACCGGCGGGA	480
ACACCGGCAC	CGTTCGGCCC	SGATGTCGCC	COATAADCOB	TGGGGAGGG	GGTGCAATT	540
GACTECATCG	CVCCCCLCCL	CCTGCTGCTG	AAADDABBTD	CCTTCCTGCC	GGGGGGCCG	600
ergreger toMVC.	AGCTCATGGG	· Y 7 37 3, Y 73 1 1	GHACTHOTHE	minderedelet/We	997 KOKOOG	660
GAGDATCOGC	CHROCCOCTC	"Arrest Mar	alladMacata.	WARCETT.	HACHAT CTG	00.
GCATGGGCAN	CAMOGTOCIA	$G:= ATA \oplus G$	Valential Area	. Transfer Jan	COMCONCETTO	78.1
$\exists A^{\prime}  \forall C \cap C \in C \cap$	CFACCLF	$C_{i,i,j} V G G Q V_{i,i,j}$	TT-MATT	1	. १ स्टब्स्टिस्टिस्टिस	r 4 1
1 - 7 - 1	A. DAN	H.T. A 76. "	Y	56 11.7	tat wat it	•
0.00		80				
	a (100)			fa fa tara	. 11	
••• • • • •			* 1			
	W 74 4	**********		N. (1)		* * 1
7						

WO 98/16645

#### PCT/US97/18214

64

GGACCGGACG GTCACCGGGG GTCACCCTGC GCGCCCAAGG AA	1362
(2) INFORMATION FOR SEQ ID NO:8:	

#### (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1458 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

	(xi) Sl	EQUENCE DESC	CFIPTION: S	EQ ID NO:8:			•
GCG/	AC BACCC	CGATATGCCG	CG IACCGTAN	CGAAAGCCG1	CGCCGACGCA	CTCGGGCGCG	60
GTAI	Caprac	CCTTGAGGAC	ATTCAGGACT	GCGTGGAGGC	CHGGCTGGGG	GAAGCCCGTC	120
TEGA	тембет	GGCCCGTGTT	TACATCATCT	ACCGGCAGCG	GCGCGCCGAG	CTGCGGACGG	180
CTAA	GGCCTT	GCTCGGCGTG	CGGGACGAGT	TAAAGÖTGAG	CTTGGCGGC	GTGACGGTAC	240
TGCG	CGAGCG	CTATUTSCTS	CACGACGAGC	AGGGCGGGCC	GGCCGAGTCG	ACCGGCGAGC	300
TGAT	'GGACCG	ATCGGCGCGC	TGTGTC%CGG	CUGCCGAGGA	CCAGTATGAG	CCGGGCTCGT	360
CGAG	GCGGTG	GGCC JAGCCG	TTCGCCACGC	TATTACGCAA	CCTGGAATT	ETGCCGAATT	420
aggr	CAUGTT	GATGAACTO!	W. W. X. W.	THEATTHE	Car regeritan	TITGITCTGC	48 Ĝ
TEAT	TGAGGA	AK STE SOTT	ELEMA MALE	e AdibiTekse	hehadeet e	CAGCTGCAGC	5 <b>5</b> A
G(3C)	TGGAGG	$\mathrm{GGO}(N) \mathrm{GGOM}$	TATOMATTOM	PARTITIONS	K : $And (Angeria)$	ATCHIGTOR	F
CCLIV	"At "IGG	Contraction of	ActivityActive		A telepith strain	Wiwilder	tib i
• • •		11 1 7, 11 11 1	*		. 19-17	91 77 FT 1	
		DZD 11 \$1-1111		11.2	27 + 1 + 1 4	A mark	
	W. 1			:		Al. vi	1
7. 17.			that the second	7. 127 · 17 · 4	· · · Ar · · ·	- 2013 - 20 <b>3</b> 0 (10)	+1
T !		Villa Amilia	· I · · · · · · · · · · · · · · · · ·	* * * * * * * * * * * * * * * * * * * *		ης .σ.σ. <b>σ.</b> ηξης:	100

WO 98/16645 PCT/US97/18214

65

deedcoedsc	CACCCGCAAG	ATCGGGGTGG	GAGTCATGAG	TTTGGCGGAA	TGCTTGCCG	1260
CACTGGGTAT	TCCGTACGAC	AGTGAAGAAG	CCGTGCGGTT	AGCCACCCGG	CTCATGCGTC	1320
GCATACAGCA	GGCGGCGCAC	ACGGCATCGC	GGAGGCTGGC	CGAAGAGCGG	GGCGCATTCC	1380
CGGCGTTCAC	CGATAGCCGG	TTCGCGCGGT	CGGGCCCGAG	GCGCAACGCA	CAGGTCACCT	1440
CCGTCGCTCC	GACGGGCA					1458

#### (2) INFORMATION FOR SEQ ID NO:9:

i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 862 base pairs
- (B) TYPE: nucleic acid (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (x1) SEQUENCE DESCRIPTION: SEQ 10 NO:9:

ACGGTGTAAT	PGTGCTGGAT	CTGGAACUGU	STRUCCION	ACCTACCGAG	ATCTACTGGC	60
GGCGCAGGGG	SCTGSCCCTG	GSCATCGCGG	TOSTCOTAGT	CGGĞATCGCG	STEGGCATCE	120
TOATCOCCTT	PETCGACAGC	Ascuccians	COMMISSION	CAGCCCCAT	AAGCCGGCCT	180
CONCONTACAG	(ATCUGAGE	:: 13.C. XI 78.	CONTRACTOR CO	$\mathbb{C}(\mathbf{M}\mathbf{GCCCCCC})$	GGGCAAACCG	240
AASGTAA/NC	managgan	, w.C., w.V. ?:	A. AWW.s	лимамамат.	Madra Mark.	<b>4</b> ∶‱. •
COCCOGTOCA	Partin	TO INTO MADE	$h(X) \cap W(X) \cap X$	man and 1970)	1 74A, 1,117 (2),	Sitt
COSTONANCO	TTT WAYAA		symple styres in	×,	EASTTONEON	<b>Q.</b>
	where !	* * *		0.0		* **
1 · · · · · · · · · · · · · · · · · · ·		·		10 m ×		1
ulli A A	* * * * *	*11		* 52, * 52, *	. The state of	• •
; :p; • · · (; )	- 727	or see . H			. There is M	£ 1
					2. To	

			20.515.6	# fs (1)		2				,
ż	 / /	INT	·URP	T/A I .	L UZIV	PUR	SEC	. 1	2017	

(i)	SEQUE	NCE	CHA	:ACTE	ŀΚΙ	STI	ICS:	
	170.3	TEME	- נויחי	800	h.		r. a i	٠.

(A) LENGTH: 622 base pairs(B) TYPE: nucleic acid(C) STRANDEDNESS: single(D) TOPOLOGY: linear

#### (x1) SEQUENCE DESCRIPTION: SEQ ID NO:10:

TTEATCAGCA COGGCAAGGO GTCACATGCC TCCCTGGGTG TGCAGGTGAC CAATGACAAA 60 GADACCOUGS GUUCCAAGAT CGTCGAAGIA UTUGCCGGTG GTGCTGCCGC GAACGGTGGA 170 STERCEAR DE COSTOSTES. MONTAGOSEN GAMBACCEC CUATCARGAS ESCOGACICA 180 . TTSGTTGCCG CCGTGCGGTC CAAAGCGCC | GSCGCCACGG TGCCGCTAAC CTTTCAGGAT 240 CCCTCGGGCG GIAGCGGCAC ACTGCAAGTC ACCCTCGGCA AGGCGGAGCA GTGATGAAGG 300 TOGOCGOGON GIGITONANG CIOGGATATA COGTGGONOC CATGGANONG CGTGOGGAGT 360 TGGTGGTTGB GCCGGCACTT OFCCTCGTCS TTGACGATCG CACGGCGCAC GGCGATGAAG 420 ALMACAGOUR RECONTESTS ACCORDING TO LARGUSASTO LOCOTITOTI SICOROGODO 480 TOSTGGGGGT STOGGT DAGT GAGTGGGAGA TO GAAATGG STEGAAGATA WGGTGATGG 540 SCOOGGEOGA COTOCEOGES VEGETOSSER SAPOSONOT SAPERCE LE SAFOTACO.  $\Theta \cup \{1$ COSAA ICCAC - CONGACATÍ CT 15.

#### CONTRACTION FOR REPORTED MODIFIES

No. 1984 to CHARA THE 1. 11 to 1.

1. TO NOTH: 1. The control of t

GCCTACGTGC	GATCGTGCCC	GGGCTACACG	TTGGACTACA	ACGCCAACGC	STCCGGTGCC	240
GGGGTGACCC	AGTTTCTCAA	CAACGAAACC	GATTTCGCCG	GCTCGGATGT	CCCGTTGAAT	300
CCGTCGACCG	GTCAACCTGA	CCGGTCGGCG	GAGCGGTGCG	GTTCCCCGGC	ATGGGACCTG	360
CCGACGGTGT	TCGGCCCGAT	CGCGATCACC	TACAATATCA	AGGGCGTGAG	CACGCTGAAT	420
CTTGACGGAC	CCACTACCGC	CAAGATTTTC	AACGGCACCA	TCACCSTSTG	GAATGATCCA	480
CAGATCCAAC	STOMADITOC	UGGCÁCCGAC	CPGCCGCCAA	CACCGATTAG	CGTTATCTTC	540
CGCAGCGACA	AGTCCGGTAC	GTCGGACĀAC	TTCCAGAAAT	ACCTOGACGG	TGTATCCAAC	600
sgggggffggs	<b>SCAAAGGCGC</b>	CAGCGAAACG	TTCAGCGGGG	GOGIOGGOUI	CGGCGCCAGC	660
GGGAACAACG	GAACGTCGGC	CCTACTGUAG	ACCACCGACG	GRYCGAYCAC	• CTACAACGAS	720
TOGTOGTTTG	CGGTGGGTAA	GCAGTTGAAC	ATGGCCCAGA	TCATCACGTC	GGCGGGTCCG	780
GATCCAGTGG	CBATCACCAC	CGAGTCGGTC	GGTAAGACAA	TOGCOGGGGC	CAAGATCAT	840
GGACAAGGCÀ	ACGACCTCCT	ATTGGACACG	TOGTOSTICT	ACAGACCCAC	ccaccercă.	900
TCTTACCCGA	TCGTGCTGGC	GACCTATGAG	ATOGTCTGCT	CGAAATACCC	GGATGČGACĠ	960
ACCGGTACTG	CGGTAAGGGC	GITTATCCAA	OCCGCCATTG	GTCCAGGCCA	AGAMGUCCTG	1020
GAMMATAM :	ROTOCATTO	GTTGUUGAAA	TUSTTICANS	"AAAATT""	a madamara	1086
AATG TAT IT	CTTÜA SITAG	TAAGGAAAT	TARRAMENTAL	$\mathcal{C}(jA_+^m,iC_1)\cap\mathcal{C}_1^m$	T. COCKOGTA	1140
ga steroryka	manna ny	T WELLEVARY	and the second s	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0 X 2 0 0 0 0 0 X	. 1717

#### Ch information for SEP to Noil i

P. MANGET OF WEAR 1994 IN THE WEAR AND A SERVICE OF THE ANALYSIS OF THE ANALYS

GGTTSCTCCA AGCOGTGGCC GCCGACGGCC GCATCCACAC CACGTTCAAC CAGACGATCG	240
COGOGACOGO COGGOTETOS TOGACOGAAS COAACCIGOA GAACATOCOG ATCOGCACOG	300
ACGCGGGCCG GCGGATCCGC GACGCGTTCG TGGTCGGGGA CGGTTACGCC GAGTTGATGA	360
CGGCCGACTA CAGCCAGATC GAGATGCCGA TCATGGGGCA CCTGTCCGGG GACGAGGGCC	420
TCATOGASSC STTCAACACC GGSGAGGACC 1GTATTCGTT CGTCGCCGTCC CGGGTGTTCG	480
GT 3T SOCIAT CGACGAGGIC ACCGGGGGAGT TGCGGGGGCCG GGTCAAGGCG ATGTCCTACG	540
GG-TEGTEFA CGG-TTGAGC GCCTACGGCC TGTCGCACCA GTTGAAAATC TCCACCGAGG	600
COTODATORO TODOTODOS DESTINATORS CONTRATORO CACTACOTOS	660
DEGEORGRAST TRANSCORRECT CHICARGUAGE RETAINANCETE GAGGGTGCTG GGCCGTCGCT	720
GOVARCIPACO CUARCITUCACIA AUCAGURACO UTCAAUTOCU GGAGGGCCCO GAGGGGGCCI	730
CGCTGAAQGC GCCCATCCAS GGCAGOGCGG CCGACATCAT CAAGGT3GCC ATGATCCA3G	840
TOGACAAGGO GOTCAACGAG GOACAGOTGG COTOGOGCAT GOTGCTGCAG GTCCACGACG	900
ASCITATI CHARATCOCI COCCOTOARO POARVEGOT CARGOCCITE GIGOGGANCA	960
ASATGGGCGS COCTTACCCG CTUGACCTCC CUCTGCAGGT GTCGGTGGGC TACGGCCGCA	1020
GOTGBOARGO GURCARRAN TOAGTBOOMA PRINTECATRIT PROCESCIMA TOGGGGGATT	1080
TTTCGGCCTT MGTT WEST TCGGCGCAA) GOGACCGA CTTTGTCCAG CCTGTACCCGT	1145
COASTRACTOR PORCA	1.00

THE INFORMATION I'M REPORT NOTES

DEASIENTE - HARACTERICTICE A LENGTH: "The complex E TOTAL FOR COMPLEX TO THATTHE MESTER OF

and switch of an expension of the first of

GGCC	CCTCGGA	TGCCGATTCC	GCACCTTCCC	GGTGGGACGG	CTĞĞCCCCCC	GAGCACGGAC	24
ATCC	GAGAACT	CTCGGGGTTC	GGCUAACGTT	ATCTCAGTGG	AATCTCAGTC	CACGCGCGCA	301
ACCI	PAGTTGT	GCAGTTACTG	TTGAAAGCCA	CACCCATGCC	AGTCCACGCA	TGGCCAAGTT	360
GGCC	CCGAGTA	GTGGGCCTAG	TACAGGAAGA	GCAACCTAGC	GACATGACGA	ATCACCCACG	420
GTAI	TCGCCA	CCGCCGCAGC	AGCCGGGAAC	CCCAGGTTAT	GCTCAGGGGC	AGCAGCAAAC	480
GTAC	CAGCCAG	CAGTTCGACT	GGCGTTACCC	ACCGTCCCCG	COCCCCCAGO	CAACCCAGTA	54(
CCST	CAACCC	TALĞAGGEGT	TGSGTGGTAC	CCBGCCGGGT	CIBAIA 1000G	GCGTGATICC	600
GAJC	ATGACG	COCOTATE	SGATGSTTCG	CCAACGCCCT	CTTGCAGKA	TGTTGGCCAT	660
3550	(TOCTO	ACGATAGCGG	TGGTGTCCGC	CGGCATCCGC	GGGGGGGGCG	CATCCCTGGT	720
SGG()	TTČAAC	OGSCCACTOG	COGN ENDOMG	CGGCGGCCCA	Grocere na	GCGCGGGGCC	780
A ACC	ATCCCC	GCAGCAAACA	TGCCGCCGGG	GTCGGTCGAA	CAGGT 3GEUG	CCAAGGTGGT	840
gong	AGTOTO	GTCATCTTGC	AAAGGGATGT	GGGCCGCCAG	TCGGAGGA33	GUTECGGCAT	900
CATT	CIGICI	GCCCAGGGGC	TGATCTTGAC	CAACAACCAC	CTGATCGCFG	CGGCCGCCAA	960
Geerr	COCCTG	GGGAGTCCGC	CGCCGAAAAC	GACGGTAACC	TTCTCTGACG	GGCGGACCGC	1020
WHOO	TTĆAĆG	GTGGTGGSGC	CTGACCCCAC	CAGTGATATC	GCCGTCGTCC	GTGTTCAGGG	1080
	"''' (' /')''	ADC DUPARTED	TCTCCCTGGA	dimensional factors	BACCTGAGG 1	TCGGTCAGCC	114.
30.7%	TIVE I	ATEC COTCO	CHOTO IGNIT	9940098411	037740,077043	GCATICATCAC	1.100
. : : T	ru naan	Mar CWardi	COACOACT FO		Asia in Madawa ina	To Trail Talests	$1 \times \omega$
ngerë:	WITCAG	VCCeMiterace	CHATCAACHC	SWITAACTC	Garage Contraction of the Contra	TGGTGAACAT	1323
ian '	T Titl	T	I-MARIN SE	AT WELL WIT	.L .::	A. M. Morrish	
	7. A.	; * * *; *	*		+ · // · =	Maria Mai	::
	-7 7	11-8	140				
VAT	a AAA	:W.W;	er mangar	101 17We 1A	$1 - \epsilon > 1 + \epsilon > \epsilon$		17.
W.		·* 1 · · · · · · · · · · · · ·	:1 'A.i	74.1 77WG 11.1	with William	en allemana.	****

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1058 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

#### (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTCCACCGCG (TGGCCCCC	C1CTAGAACT	ACTGGATUCC	COGGGCTGCA	GGAATTEGGC	60
ACGAGGAICC GACGICGCAG	GTTGTCGAAC	ccaccaçada	C.; NACTNTCÇ	GTCCATCCCT	120
ABONOGGOGA TOGGOBAGOGE	COGAATOOCG	CUASTGAGGA	GGCGGGCAAT	TTUGUGGGGGC	180
COGGCGACGG CGAGCGCCGG	AATOGEGEGA	GTGAGGAGGC	GUGCAGTÇAT	GCCCAGCGTG	240
ATCCAATCAA CCTGCATTCe	GCCTGCGGGC	CCATTTGACA	AYCGAGGTAG	TGAGC GCAAA	300
TGAATGAIGG AAAACGGGCG	GTGACGTCCG	CTGTTCTGGP	GGTGCTAGGT	GCCTGCCTGG	360
CGTTGTGGCT ATCAGGATG1	TCTTCGCCGA	AACCTGATGC	GGAGGAACAG	GGTGTTCCCG	420
TRAGOCCGAC GROSTCCGAC	00000000000000000000000000000000000000	TOGOOGAGAO	CAGGCAGTCG	CTTGATGCGA	480
TAAAAGGSTT GAGCAGCGTG	CACGTAGEG ;	TO JOHACAA .	COGGAAAGTC	GACAGCTTGC	440
TGUGTATTAC CAGTOCCGAT	interiAción, el	· G FICKNICC	STRUCKUGGEA	AAGGGCGTAT	600
BACBAGBABB AWATERANG	.::7: T	**************************************	A FEMANAT	25 19 19 19 15 A	(16,1
AACTGTTCGA ACGACTGGA C	Walland J	13771701.00	A TOTHAKT	To Accordance	790
TOGATICTON TWY 1 GARRE	ATGTA ETFET		W. GWA. ST.	www.ccawa	··· •) , ;
ing was to wrome our	***************************************	Consider the	· Art Ive	1/1 -	7 4
	• :		•	* 1	
	*. ""			ers i l'és	• •
mining managaway	· · · · · · · · · · · · · · · · · · ·	a 1 1 5 A 11			1 4
milmowa albumi a	***********				

### (D) TOPOLOGY: linear

	(xi) 5	SEQUENCE	DESC	CRIPTION: S	EQ ID NO:15	:		
GAAT	TCGGC	CGAGAG	GTGA	TCGACATCAT	CGGGACCAGC	CCCACATCCT	GGGAACAGGC	60
GGCG	GÇGGAC	GCGGTC	CAGC	GGGC 3CGGGA	TAGCGTCGAT	GACATCCGCC	TOSCTOSGT	120
CATT	GAGNAG	GACATG	GCCG	TGGATAGCGC	CGGCAAGATC	ACCTACCGCA	TCAAGCTCGA	180
AGTG	TOSTT	: AAGATG	AGGC	ÉGGCGCAACU	GCCCTAGCAC	GGGCCGCCGA	GCAAGACGCA	240
λаћт	qqcåce	GTTIGO	GGTT	GATTCGTÖCH	ATTITGTGTC	TGCTCGCCGA	GGCCTACCAG	3,00
1700	ĞGDÜCA	. gg1.cca	Jama .	CTGCGTATI	CAGGCGTGCA	TUGCGATTOU	SHEGGERCACG	360
CCGG.	AGTTAA	TGCTTC	GCGT	CGACCCGAA	TGGGCGATCC	GCCGGNGAGG	TGATCGATGA	420
CCGT	GGCTAG	CCCGTCC	GATG	CCCGAGTTG	CCGAGGAAAC	GTGCTGCCAG	UCCUGTAGGA	480
rd, G	TCCGTA	. GGCGGCC	3076	CTGACCGGCT	CTGGCTGCGC	CCTCAGTGCG	GCCAGCGAGC	540
3G								542
12)	INFORM	ATION E	OR SE	Q 10 NO:16:				

(:) SEQUENCE CHARACTERCUTICS:

"A) LENGTH: H' base parry

(B) TYPE: nucleuc agid

TO STRANDEDNESS, cincle So TOP MOSE: Linear

". is, and does the it, in the ens regarde with an element of the result of the region of NORTH AND A CONTRACT OF A SERVICE AND A SERVICE OF A SERVICE OF A SERVICE AND A SERVIC COMPANIES CONTRACTOR OF THAT OF ANY ANALYSIS OF THE SECOND OF THE SECOND

WO 98/16645

GTTTGCCGCC	AATATTCJGG	GSGCACCGCC	AGACCCGCCG	GGGCCACCAT	TGCCGCCGGG	480
CACCGAAACA	ACAGCCCAAC	GOTGOOGCCG	GCCCCGCCGT	TTGCCGCCAT	CACCGGCCAT	540
TCACCGCCAG	CACCGCCGTT	AATGTTTATG	AACCCGGTAC	CGCCAGCGCG	GCCCCTATTG	600
CCGGGCGCCG	GAGNGCGTGC	CCGCCGGCGC	CGCCAACGCC	CAAAAGCCCG	GGGTTGCCAC	660
aggadaaggaa	GGACDUACCO	GTCCCGCCGA	TCCCCCCGTT	GCCGCCGGTG	CCGCCGCCAT	720
TGGTGCTGCT	ÇAAGJOGTTA	GCGCCGGTTC	CGCSGCTTCC	GCCGGTGCCG	CCNTGGCCGC	780
ceracceecc	STTGTUSTAT	Adddaccato	marragecene	GTTGCCSCCA	TTGCCGCCAT	840
TOCCGCCGTT	SCCGCCATTER	CAGCCGTTCC	carcaccacc	GCCGGNTTGG	CCGCGGGGGG	900
caccaacaac	CGC					913

### (2) INFORMATION FOR SEQ ID NO:10:

# (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1872 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

### CREE DEQUELATE DESCRIPTION: SEC 10 Sec 1/:

GACATTGAAC	GGGTTGATCC	AGTTCGATGC	CGCAATCCAG	G CCCGGTGATI	, ceasaceaca	720
CGTCGTCAAC	GGCCTAGGAC	AGGTGGTCGG	TATGAACACG	GCCGCGTCCG	ATAACTTCCA	780
GCTGTCCCAG	GGTGGGCAGG	GATTCGCCAT	TCCGATCGGG	CAGGCGATGG	CGATCGCGGG	840
CCAAATCCGA	TCGGGTGGGG	GGTCACCCAC	CGTTCATATC	GGGCCTACCG	CCTTCCTCGG	900
CTTGGGTGTT	GTCGACAACA	ACGGCAACG	CCCACGAGTC	CAACGCGTCG	TCGGAAGCCC	960
TCCGGCGGCA	AGTCTCGGCA	TOTOCACOGG	CGACGTGATC	ACCGCGGTCG	ACGGCGCTCC	1020
GATCAACTCG	GCCACCGCGA	TGGCGGACGC	GCTTAACGGG	CATÉATCCCG	GTGACGTCAT	1080
CT./GGTGAAC	TGGCAAACCA	AUTOGGGGGG	CACCOCCTACA	ARTROBANDOO	CATTGGCCGA	1140
GGSACCCCCC	GCCTGATTTC	TOGCOGATAC	CACCTOGCOGG	CCGGCCNATT	GGATTGGCGC	1200
CADCOGTGAT	TGCGGGGGA	GCCCCCGAGT	recertates	GIECECGIGE	CATTGTGGAA	1260
GCAATGAACG	AGGCAGAACA	CAGCGTTGAG	CACCCTCCCG	TGCAGGGCAG	TTACGTOGAA	1320
GGCGGTGTGG	TOGAGOATOO	GGATGCCAAG	GACTTCGGCA	Gågadagatia	COTICOOGCO	1380
GATCCGACCT	GGTTTAAGCA	CGCCGTUTTC	TACGAGGTUC	TGGTCUGGGC	GTTCTTCGAC	1440
GCCAGCGCGG	ACGGTTCCGN	CGATCTGCGT	GGACTCATCG	ATCCCCTCFA	CTACCTGCAG	1500
TGGCTTGGCA	TOGACTOCAT	CIGITGCCSC	CGTTCCTACS	actëaecse i	GCCCGACGGC	1560
GGTTAGGACA	TTCGCGACTT	CTACAAĞƏTƏ	CTGCCCGAAT	TCGGCACCGT	CEACGATTTC	1620
GTCGCCCTGG		mayarı başır (7)	a d'Ambarra	INATUACCOA	FOLGGEGATG	, <del>(</del> ,8%)
<b>Vy</b> acvovaca.	CGĞAGTCUDA	over Technic	CAGUAGTOU	od at attation a	AGACCOACTC	***
TACGGTGACT	ATTACGETT	dN: cNcV	APTATION.	WWW WWW		; W.
ideath Weit	11 1/1/4 - 10 - 1	.,	*** *	i saanii ya	7	, et i
						100

<sup>.</sup> TENTEN E SHARWITER COMPANI A. ISBN 182 I Post base part. B. TYPE: C. Pert and i TORING WITE STA

CTTCG DCGAA	ACCTGATGCG	GAGGAACAGG	GTGTTCCCGT	GAGCCCGACG	GCGTCCGACC	60
CCGCGCTCCT	CGCCGAGATC	AGGCAGTCGC	TTGATGCGAC	AAAAGGGTTG	ACCAGCGTGC	120
ACGTAGCGGT	CCGAACAACC	GGGAAAGTCG	ACAGCTTGCT	GGGTATTACC	AGTGCCGATG	180
TCGACGTCCG	GGCCAATCCC	CTCGCGGCAA	AGGGCGTATG	CACCTACAAC	GACGACCAGG	240
GTGTCCCGTT	TCGGGTACAA	GGCGACAACA	TCTCGGTGAA	ACTGTTCGAC	JACTGGAGCA	300
ATCTC-GCTC	GATTTCTGAA	CTGTCAACIT	CACGCGTGCT	CGATCCTGCC	GCTGGGGTGA	360
CGCAGCTHCT	GTCCCGTGTC	ACGAACCTCC	AAGCGCAAGG	TACCGAAGTG	ATAGA EGGAA	4.20
TTTCG CONC	CAMAATCACO	GGÇACCAT. C	CristioAG IIC	TSTCAAGATG	TTGATCCTG	4 8 ()
GCGCCAARAG	TGCANGGTGG	OCGACCGIGI	GGATTGCCCA	GGACGGCTCG	CACCACCTOS	5:0
TONGA KÖGAG	CATCGACCTC	CONTCCOGST	CGATTCAGCT	CACGCAGTCG	AAATG SAACG	61. C
AADCCOTCAA	CGTCGAGTAG	GOOGAAGTTG	CGTCGACGCG	TTRITTOGAAA	JECCCTTETE	650
AACGGTGTCA	ACGGCACCCG	AAAACTGACC	CCCTGACGGC	ATCTGAAAAT	TGACCCCCTA	720
FACCGHIGGIG	TTGGTGGTTA	TTCTTCGGTG	CTTCCGCCTC	GTGGGACGCG	GCCGAGGTCG	780
GGTCTTTGA	GUCGGTAGCI	GTCGCCTTTC	AGGGCGACGA	CTTCAGCATG	GTGGACGAGG	8:0
GGTCSATCA	TGGCGGCAGC	AACGACGTCG	TCGCCGCCGA	DOBOTODAAA	CCACCGGCCG	900
AAGOON TTAT	TÖÖACÜTGAY	CATCAAGCTG	CCCCCCTCAT	ACCGGGAGSA	CACCAGCTGG	96()
TOSM MALTE	TOHULANTIN		PARTOTARO	CONCTROUTO	AAC MCCAGG	* 17 m/s
Varrigatage	-1000MAN1011	GGTGA ITTOL	m magamod	000 000FG	STGAGGGTCG	1080
A. Www. arkin.	CTATTONTO	umma viiti ji	3 1 150 1 2 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1		(4) 5 (1)(A('A)(')	1140
)	Witte V	t talan	2:	01743	· AAMA	1
4. 1 1 1 H 1.1 1		an Alba	-	*	1 1 1011	• • • •
iran in	yw iadu i	Maria Section	11 + 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	vi 11 "	in styl Vacet	1 2,50
· . · · · · · · · · · · · · · ·	Alexandra	$\gamma,  \cdot^{(2)} t_{i}  \cdot  \cdot  \gamma^{-n}  ,$		· / / · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	; · · · · ·
	ATELLITE					1441
			* -			0.4

(C)	STRANDEDNESS: single
(D)	TCFOLOGY: linear

# (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GAATTCGGCA	CGAGCCGGCG	ATAGCTTCTG	3GCCGC3GCC	GACCAGATGG	CTCGAGGGTT	60
DGTGCTCGGG	GCCAC /GCCG	GGCG PACCAC	JC PGAC JĞGT	GAGGGCCTGC	AACACGCCGA	120
COGTCACTOG	TTOAT BOTGG	ACGCCACCAA	CCCGGCOGTG	GTTGCCTACG	ACCOGGCCTT	180
DGCCTAGGAA	. ATOGGDIACA	TOGNOGAAAG	CGUACTOGCC	AGGATGTGCG	GGGAGAACCC	240
G GAGAA JATO	TTOTTOTACA	TUACUGTUTA	CAACGAUCCG	IACGTGCAGC	CGCCGGAGCC	300
GGAGAACTTC	GATCCCGAGG	GOGTGCTGGG	GGGTATCTAC	CGNTATCACG	CGGCCACCGA	360
GUAAGGCACC	AACAAGGNGC	AGATICTGGC	CTCCGGGGTA	GCCATGCCCG	CGGCGCTGCG	420
GGCAGCACAG	ATGCTGGCCG	CUGA-TIGGUA	ng redecace	GACGTGTGGT	CGGTGACCAG	480
TTGGGGGGGAG	CTAAACCGCG	ACGGGGTGGT	CATCGAGACC	GAGAAGCTCC	GOCACCCGA	540
TCGGCCGGCG	GGCGTGCCCT	ACGTGACGAG	AGEGETGGAG	AATGCTCGGG	GCCCGGTGAT	600
CGCGCTGTCG	GACTGGATGC	GCGCGGTCCC	CGAGCAGATC	CGACCGTSGG	TGCCGGGCAC	€60
ATACCTCACC	##GGGCANCG	ACGGUTTCGG	TTTTCCGAC	ACTCGGCCCC	cossicoros	720
TTACTICAAC	$V_{\alpha,\beta}((Y_{b},G_{a},\beta))$	AATTCCAG Ĥ	TRITOGERIT	TTGGGAGEG	GTT SHCCGGG	780
ICGAMAGGTG	AAIRIQARO	CATTUGGTG	0.3020.000.000	hadeaccara.	ARTIA COST	84r
7.11 1 to 1 1/2 to	7.11 1 7 7 7 7 7	711223	MAGIT			

# . DE PMATERIE E VE. 1 Notion

THE PROPERTY OF A PROPERTY OF A STREET FOR THE PROPERTY OF A STREET OF A STREE

CAGATTCATA	ACGAATTCAC	AGCGGCACAA	CAATATGTCG	CGATCGCGGT	TTATTTCGAC	120
AGCGAAGACC	TGCCGCAGTT	GGCGAAGCAT	TTTTACAGCC	AAGCGGTCGA	GGAACGAAAC	180
CATGCAATGA	TGCTCGTGCA	ACACCTGCTC	GACCGCGACC	TTCGTGTCGA	AATTCCCGGC	240
GTAGACACGG	TGCGAAACCA	GTTCGACAGA	CCCCGCGAGG	CACTGGCGCT	GGCGCTCGAT	300
CAGGAACGÇA	CAGTCACCGA	CCAGGTCGGT	CGGCTGACAG	CGGTGGCCCG	CGACGAGGGC	360
GATTTCCTCG	GCGAGCAGTT	CATGCAGTGG	TTCTTGCACG	AACAGATCGA	AGAGGTGGCC	420
TT:GATG 3CAA	CCCTGGTGCG	GGTTGCCGAT	cecccceccc	CCAACCTGTT	CGAGCTAGAG	480
AMOTT COLUG	CACGTGAAGT	SGATGTGCCG	CCGGCOGCAT	CAGGCGCCCC	GCACGITGCC	540
003GG 20G 10	TOTAGATOON	rgg-ggggat	CAGCGAGTGG	TCCCUTTCGC	cogcoper or	600 600
TODAGDUAGG	COTIGGIGG	GCCGGGGTGG	TGAGTACCAA	TCCAGGCCAC	CCCGACCTCC	660
CGCNAAAAGT	DGATGTCCTC	STACTCATOG	ACSTTCCAGG	AGTACACCGC	CCGGCCCTGA	720
GCTGCCGAGC	GGTCAACGAG	TTGCGGATAT	TOCTTTAACG	CAGGCAGTGA	GGGTCCCACG	780
GCGGTTGGCC	CGACCGCCGT	GCCGCACTG	CTGGTCAGGT	ATCGGGGGGGT	CTTCHOGAGO	840
AACAACGTCG	GCAGGAGGGG	TGGAGCCCGC	CGGATCCGCA	GACCGGGGGG	GGGAAAACGA	900
CATCAACACC	GCACGGGATG	GATCTGCGGA	GGGGGGTGCG	GGAATACTGA	ACCEGTTAG	960
GAGCGCCAGC	AGTTGTTTTT	CCACCASCGA	AGRITTTOG	COTTAIN CORRESPONDE	GGCENTTAAG	1020
T						1 (1) (-)

# (2) INFORMATION FOR SEQ ID NO. 21:

- TO SEQUENCE CHARACTERISTICS:
  - A DENGLE TIPE OF ALT SP CYPE INTO A CONTROL OF A CONTROL

多数,是完美的是《新沙斯·拉拉·拉斯·克克·拉斯·克耳·

WO 98/16	645
----------	-----

#### PCT/US97/18214

77

TECGCCCCTG ANGTCCEGAC ÉGCCCCCCAG TGGACCAGNC TGETCAACAG NOTCGNEGAT	240							
CCCAACGTGT CGTTTGNGAA CAAGGGNAGT CTGGTCGAGG GNGGNATCGG NGGNANCGAG	300							
GGNGNGNATC GNCGANCACA A	321							
(2) INFORMATION FOR SEQ ID NO:22:								
(i) SEQUENCE CHARACTERISTICS:								
(A) LENGTH: 373 base pairs								
(ii) minorii. 575 case paris								

(x)) SEQUENCE DESCRIPTION: SEQ 10 NO:22:

(B) TYPE: nucleic acid(C) STRANDEDNESS: single(D) TOFGLOGY: linear

TOTTATURET TOURGITISC BAIRGETTIT GERNGUNGER GETTAACCC CTCBGCCAGC 60

CGATCSACGG GEOGRASAC GTCBACTCIS ATACTCRICG CGCBCTGSAG CTCBAGGIGC 120

CCTCGGTGGT GNACCGGDAA GUCGTGAAGG AGCBGTTGNA GACDGRGATC AAGGGGATTG 180

ACGCGATGAC GIGGATCRIC CGCAGGCAGC GCCBAGGACC CGCAAGACCG 240

GCAAAAAACCG CIGTCTGTGT CGGACACCAT CCTCAAACCA GCCGGAAGAA CTGGGAGTCC 300

GGTUGATCCC AAGAAGCA G TOCGCTTGTS TATA GTTGG CIATCRIGGTA AGAAGGGGAA 360

CCTTAACATCRI CCG

# A. & INFORMATION FOR SECTION NO.23:

\*.) DEQUENCE CHARACTERLITTCH:

(A) LENGTH: (%) hase prix
b Till: A. 1. article
1 (18ANATMENT): Length
5 (1 Till: Linear

examplement for the first of the property

GUGTGGAGGT TTTUGTCAUC GUUAGUCGTG GNAAGTGGGA CAUGUTGUGU GUCATNGNGT	300
TTGACGACGA NCCATATCGG NGATTCCCNC ACATNCGAAG TTCCGANGGA GA	352
(2) INFORMATION FOR SEQ ID NO:24:	
(i) SEQUENCE CHARACTERISTICS:  (A) LENGTH: 7.6 base pairs  (B) TYPE: nucleic acid	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

(C) STRANDEDNESS: single (D) TOPOLOGY: linear

GAAATJUGCG	TTCATTCCGT	TOGACUAGOG	GCTGGCGATA	ATCGACGAAG	TGATCAAGCC	60
GCGGTT030G	GEGETCATGS	GTCACAGCGA	GTAATCAGCA	AGTTOTOTGG	TATATCGCAC	120
CTAGCOTOCA	GTTGCTTGCC	AGATOGOTTT	CGTACCGICA	TOGCATGTAC	COGTTCGCGT	187
SCOGCACCOT	CATGUTGGC	GCGFGCATCC	r #GCCACGGG	TOTOGOGGGT	CTCGGGGTCG	240
SCGCGCACTO	CGCAGCCCAA	AC /3CGCCGG	TGCCCGACTA	CTACTGGTGC	ccggggcagc	300
CTTTCGACCC	CGCATGGGGG	CCCAACTGÓG	ATCCCTACAC	CTCCCATCAC	GACTTCCACC	360
FICGACALICUA	$\cup \emptyset \emptyset \emptyset \cup \emptyset ( \emptyset \emptyset ) \emptyset \cup \emptyset \emptyset \emptyset $	MANA MODELL	AUTACUCUGG	ACHIATORITA	GAAGGTCCCG	420
LOCTTOACJA				1.3777777 1.477	TATAGCGCI	48)
$(\zeta((\mathbf{L}_{i}^{*})\mathbf{M}(\zeta_{i})):$	GCCGCATCAG	22474445	TATAAACCCG	SHOUTGENE.	CJSWAAGCTA	540
NATE OF STREET	Translavior		,; 'AT	Act to the second of	ACT ACTO	to the
AAAAC (C) - CC	A 11 777 17 11			gaverna.	ing words	
$(f_{i}, f_{i})_{i}$	4 - 50 - 50	16,471.1 1 4 1	W. II. II.		rolling	
A**-***I+-						

THE DANGER OF STREET

TO THE THE RESERVE OF THE STATE OF THE STATE

(xi)	SEQUENCE	DESCRIE	TION: S	EQ ID NO:25	:				
CGCGACGAC	G ACGAACG	TCG GGC	CCACCAC	CGCCTATGCG	TTGATGCAGG	CGACCGGGAT	60		
GGTCGCCGA	C CATATCO	AAG CAT	GCTGGGT	GCCCACTGAG	CGACCTTTTG	ACCAGCCGGG	120		
CTGCCCGAT	G GCGGCCC	GGT GAA	GTCATTG	CGCCGGGGCT	TGTGCACCTG	ATGAACCCGA	180		
ATAGGGAACA	A ATAGGGG	EGT GAT	TTGGCAG	TTCAATGTCG	GREATGGCTG	GAAATCCAAT	240		
GGCGGGGCA1	T SCTEGGE	GCC JAC	CAGGCTC	GCGCAGGCGG	GUUAGCGCGA	ATCTGGAGGG	300		
AGCACTCAAT	r -agoggog.	ATS AAG	CCCCGGA	CCGGCGACGG	TOCTTTGGAA	GCAACTAAGG	360		
AGGGGGGGG	G CATTGTG.	ATU CGA	GTACCAC	TTGAGGGTGG	CGGTCGCCTG	GTCGTCGAGC	420		
TGACACCCGF	a CGAAGCC	add GCA	CTGGG. G	ACGAACTCAA	ACCCCTTACT	AGCTAAGACC	480		
AGCCCAACGG	GGAATGG"	rcg gcg:	PTACGCG	CACACCTTCC	GGTAGATGTC	CAGTGTCTGC	540		
FCGGCGATGT	TATGCCCAC	GGA GAAG	CTCTTGG	ATACAGCGCT			580		
(2) INFORM	MATION FOR	R SEQ 11	) NO:26:						
(i) SEQUENCE CHARACTERISTICS:  (A) LENGTH. 160 base pairs  (B) TYPF: nucleic acid  (C) STRANDEDNESS: single  (D) TOPOLOGY: linear									

(x)) SEQUENCE DESCRIPTION: PEG 10 Notice:	
AAC RIARRIC Et 1999 RICCO (1, 1 FTG, Report) (1, 1	+ -*
TATION OF OUTSIDE THE HOUSE OF THE HOUSE OF THE CONTROL OF THE CON	
ACAMA A TANA AMAMA	
L THE RMATTON FOR ALL II NOT .	
) SEQUENCE CHAMACTERRICTIVE:  A FENCTH: Contact that  FOR THE CONTROL OF THE	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:	
GACACCGATA CGATGGTGAT GTACGCCAAC GTTGTCGACA CGC	TCGAGGC GTTCACGATC 60
CAGCGCACAC CCGACGGCGT GACCATCGGC GATGCGGCCC CGT	TCGCGGA GGCGGCTGCC 120
AAGGCGATGG GAATCGACAA GCTGCGGGTA ATTCATACCG GAA	TGGACCC CGTCGTCGCT 180
GAACGCGAAC AGTGGGACGA CGGCAACAAC ACGTTGGCGT TGG	CGCCCGG TGTCGTTGTC 240
GCCTACGAGO GCAAGGTACA GACCAAGGCC CO	272
(2) INFORMATION FOR SEQ ID NO:28:	

# (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 317 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (1) TOPOLOGY: linear

### (xi) SEQUENCE DESCRIPTION: SEQ 10 NO:28:

GCAGCCGGTG	GTTCTCGGAC	TATCTGCGCA	COGTGACGCA	GCGCGACGTG	CGCGAGCTGA	60
AGCGGATCGA	GCAGACGGAT	CGCCTGCCGC	GOTTCATGCG	CTACCTGGCC	CCTATCACCG	120
CGTAGGAGCT	GAACGTGGCC	GAAGCGGCGC	GGTCATCGS	G FTCGACGUS	GEGACGATOC	180
GTTCGBATCT	TOCKTARTIC	GAGA GGTGT	ATCTGCTACA	Treatist	SICT-GTCG	.1444
GGAATGTĞAG	CGCGAAGATC	AAGAAGCCGT	CAAAGATCCA	COTCCTCCAC	AGTGGCTTCG	300
CGCCCTGGTT	Argraga					: ;

INFOMMENT FRANCE, IN NO. 10

- AND MANUAL TRANSPORTED BY A STREET OF THE STREET STREET STREET STREET.
- $\mathcal{E} = T^{*}H^{+}H^{*}_{*}H^{*}_{*}H^{*}_{*} + H^{*}_{*}H^{*}_{*} + H^$

WO 98/16645	PCT/US97/18214

GCAGCCCCC ACCACCTCCC CGCTGGCCAL CATGCTGATG ACCACGTCGG CCTCGGCCAC	120
CGCTTCGGGC GCGCTACGAA ACACCGCGGAC ACCGTGCGCG GCGGCGCGG ACGCCGCCGT	180
GG	182
(2) INFORMATION FOR SEQ ID NO:30:	
(i) SEQUENCE CHARACTERISTICS:  (A) LENGTH: 308 base pairs  (B) TYPE: nucleic acid  (C) STRANDEDNESS: single  (D) TOPOLOGY: linear	
(xi) SEQUENCE DESCRIPTION: SEQ 10 NO:30:	
TEDECEAAD TITESTOADE ASSTORTED ASSTORTED ASSTORED TO TO TO THE FAADDO TAKE	·5.)
GGGGGTTCAC GAGGGGAAGA CAUGCCTGTC CGAGCTGCTC CGGCTGGTCT ACGGCGGGGA.	120
GAGGTTGAGA TTGCCCCCCC CGGCCGACCCC CTAGCAAAGC TTGTGCCGCT GCATCCTCAT	180
BAGACTOGGO OGTTARGOAT TGAUCATRGO UTGTACCIOG TGCCCGACGA TTTGGACGCT	240
CGTTGTCAG ACCACGTGCT CGAACGTTTT CACCCGTGAA GCGCTACCTC ATCGACACCC	3(10
ACCTPYTGG	308
(Z) INFORMATION FOR SEQ ID NO.31:  (A) LENGTH: 267 base pairs  (B) TYPE: nucleic a inf  (C) STRANDEDNESS: sincle  (10) TO COLOGY: Linear	
w. The Hill to the special of the High House Hills	
TOWN HAR THE STANDARD RESIDENCE OF THE PROPERTY OF THE PROPERT	
THE PROPERTY OF THE PROPERTY O	
order myerr. Kirolin erreg ergenne eine e	

WO 98/16645 PCT/US97/18214

82

# (2) INFORMATION FOR SEQ ID NO: 32:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1539 base pairs

(B) TYPE: nucleic acid

(C) STPANDEDNESS: single

(D) TOPOLOGY: linear

# (xi; SEQUENCE DESCRIPTION: SEQ ID NO:32:

CTCGTGCCGA	AAGAATGTGA	GGGGACACGA	TGAGCAATCA	CACCTACCGA	GTGATCGAGA	60
TOSTÓGGGAC	CTCGCCCCAC	SGCGTCGACG	CGGCAATCUA	JGGCGCTCTG	GCTCGAGCTG	120
C.J. AGADPAT	geatracat fa	SATTGGTTCG	AAGTACAGTC	AATTOGAGGO	CACCTGGTCG	180
ACGGAGGGGT	CGCGCACTTC	CAGGTGACTA	TGAAAGTCGG	CTTCCGCTGG	AGGATTCCTG	240
AACCTTCAAG	CGCGGCCGAT	AACTGAGGTÖ	CATCATTAAG	CGACTTTTCC	AĞAACATCCT	30¢
GACGCCCTCG	AAACGCGGTT	CARCCGACGS	TGGCTCCGCC	GAGGCGCTCC	CTCCAAAATC	360
CCTGCGAGAA	TTCGTCGGGG	GCGCCTACAA	GUANGTEGGT	CCTGAATTCG	TEGGGTATET	420
GETOGACOTG	TOTGGGCTGC	AGCCGGACGA	MaCGGTGCT'	GACGTCGGCT	GAGGATAGGG	480
SCEGATEGEG	TTGCCGCTCA	CCGGCTATCT	GNATAGCGAG	GGACGCTACG	CCGGCTTCGA	5.10
TATCTCGCAS	AAAGCCATCG	derragradudA	UBACCACATO	ACTROCOCCE	LOCCOAACTT	600
DENGTTOGAG	GTCTTCBAÇA	T - 2/14 /11 = 1 +	FT HACAA	COMMON IN	AATA/ CAGTC	оыс
ACTAGACTTT	CGCTTTCCAT	ATTTGGATGC	- HORTEGMI	GPGGTGTTT	1TACCTCGGT	720
dad William	Variation, see	mananan o.	* ***	N. 14747		· . <u></u>
$\mathcal{A}_{N}:\cdots::$	5. 13.7.1	:-":"}::	** ** * * * *	· · · · · · · · · · · · · · · · · · ·	1	A(.*
	41.4 1. 4.4 I	5p 1.14 **		di.	es o significan	••
* 1/1 / 1/1/1 / 1/1/1 *	· : : : · · · ::::::::::::::::::::::::	NA CHINA TO			- FWL CTUA	ă, -
THE WAR COTT	:	para saan sa	ATC WITA	: :	1.00000000	1 %.
	. 0	0.00			• •	

WO 98/16645 PCT/US97/18214

83

AGCCTGCGTG	ATCGGTCATC	ACCAACGGTG	ACAGCAGCCG	GTTGTGCACC	AGCGCGAACG	1320
CCACCCGGT	CTCCGGGTCT	STOCAGOOGA	TOGAGOOGCC	CAAGCCCACA	TGACCAAACC	1380
CCGGCATCAC	GTTGCCGATC	GGCATACCGT	GATAGCCAAG	ATGAAAATTT	AAGGGCACCA	1440
ATAGATTTCG	ATCCGGCAGA	ACTTGCCGTC	GGTTGCGGGT	CAGGCCCGTG	ACCAGCTCCC	1500
GCGACAAGAA	CCGTATGCCG	TCGATCTCGC	CTCGTGCCG			1539

# (2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 851 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (b) TOPOLOGY. Linear

# (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

CTGCAGGGTG GCGTGGATGA GCGTCACCGC GGGGCAGGCC GAGCTGACCG CCGCCCAGGT 60 COGGGTTGCT GCGCCGGCCT ACGAGACGCC GTATCCCCTG ACGGTGCCCC CGCCGGTGAT 120 CGCCGAJAAC CGTCCTGAAC TGATGATTCT GATAGCGACC AACCTTTGG GGCAAAACAC 180 CCCGGCBATC GCCCCGAACG AGGCCGAATA CGGCGAGATG TGCCCCCAAG ACGCCGCCGC 1.40 GATGTTTSGG TAGGOGGGG GGACGTGGAG GEGGAG TOGG AGGTTGTTSGG GGTTGFAGGA GOOGENSONG ATOMORADES CENTROGOST ESTOSANIAS AS SERVICES IS TO TOMOGRAGO 30,0 CTONIACACI OPPOSED NA LO ACETEAT MAA MATET E COMA ESCENTIGAAACAGTT 4000 April 1 . A 4 With the theath of Albertains The Third the second of the se and the state of t 

WO 98/16645	PCT/US97/18214
-------------	----------------

0	4
O	+

(2)	1.05%	ORMAT	ION	FÓR	SEQ	ID	NO:	34:	:
	(i)	SEQUI	ENCE	СНА	ARACT	ERI	STI	ICS:	
		(A)	LEN	GTH:	254	ba	se	pai	r.s

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

### (x1) SEQUENCE DESCRIPTION: SEC ID NO:34:

GATCGATCGG GCGGAAATTT GGACCAGATT CGCCTCCGGC GATAACCCAA TCAATCGAAC 60
CTAGATTAT TCCGTCCAGG GGCCGGAGTA ATGGCTCGCA GGACGGAAC CTTACTGCTG 120
CGGGCACCTG TCGTAGGTCC TUGATAUGGC GGAAGGGGTC GACATTTTCC ACUGACACCC 180
CCATUCAAAC GTTCGAGGGC CACTCCAGCT TGTGAGCGAG GCGAGGCAGT CGCAGGCTGC 240
GCTTGGTCAA GATC 254

### (2) INFORMATION FOR SEQ ID NO:35:

- (I) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1227 base pairs
  - (B) TYPE: nucleic acid
  - (G) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

### RED COMPUTENCE OFFICE LETTON: CECCLO DE 35:

ANTONIA DE ARATTERIO DE MARCELLA DE RECENTRA DE LA COMPANSIONE DEL COMPANSIONE DEL COMPANSIONE DE LA COMPANSIONE DEL COMPA

CGCTGCTCAG	CTTGGCCAAG	GCCTGATCGG	AGCGUTTGTC	GCGCACGCCG	TOSTOGATAC	600
CGCACAGCGC	ATTGCGAACG	ATGGTGTCCA	CATCGCGGTT	CICCAGCGCG	TTGAGGTATC	66O
CCTGAATCGC	GGTTTTGGCC	GGTCCCTCCG	AGAATGTGCC	TGCCGTGTTG	GCTCCGTTGG	720
TGCCGACCCC	GTATATGATC	GCCGCCGTCA	TAGCCGACAC	CAGCGCGAGG	GCTACCACAA	780
TGCCGATCAG	CAGCCGCTTG	TGCCGTCGCT	TCGGGTAGGA	CACCTGCCGC	GGCACGCCGG	840
CATATGCGCC	ango agawac	HCCGCGTCGT	CTGACGGTCC	CGGGGGGAAG	GCCGGTTCGG	9()
CCGCCCCCAG	91.00 k3 1886	ragteca336	CTTGGGGTTC	GTGGGATGAG	GGCTCGGGGGT	960
ACGGGGGCGG	TOCGTTRATC	radhananan	COTTCOCCGA	GT3G3GACCG	GGCATTGTGG	1020
TTCTCCTAGG	OTGGTG ACG	GACCAGCTG	CTAGGGCGAC	AACCGCCCGT	CGCGTCAGCC	1080
GGCAGCATČG	GCAATCAGGT	GAGCTCCCTA	GGCAGGCTAG	ÇGCAACAGÖT	GCCGTCAGCT	1110
CTCAACGCGA	aggagraga	ngaggagaaa	ATAATGTTGA	AAGACTAGGC	AACCTTAGGA	1200
ACGAAGGACG	GAGATTITGT	GACGATO				1227
(2) INFORMA	ATION FOR SE	10 ID No:36:				

#### / International Part Cont. To the Pro-

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 181 base pairs
  - (B) TYPF: nucleic acid
  - (C: STRANDEDNESS: Single
  - (D: TOPOLOGY linear

### N THOURN'S PROPERTIES. IN MICH.

\* The state of the

THE EMPLIE DIE BOTH A THOUGHT IN THE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:							
GCGGTGTCGG CGGATCCGGC GGGTGGTTGA ACGGCAACGG	CGGTGTCGGC	GGCCGGGGCG	60				
GCGACGGCGT CTTTGCCGGT GCCGGCGGCC AGGGCGGCCT	CGGTGGGCAG	GGCGGCAATG	120				
GCGGCGGCTC CACCOGCGGC AACGGCCGTC TTGGCGGCGC	GGGCGGTGGC	GGAGGCAACG	180				
CCCCGGAUGG CGGCTTCGGT GGCAACGGCG GTAAGGGTGG	CCAGGGCGGN	ATTGGCGGCG	240				
GCACTCAGAG GGCGACCGGC CTCGGRGGTG ACCGCGGTGA	CCGCGGTGAC		290				
(2) INFORMATION FOR SEQ ID NO:38:							
(i) SEQUENCE CHARACTERISTICS:  (A) LENGTH: 34 base pairs  (B) TYPE: nucleic acid  (C) STEVANDEDNESS: Single  (D) TOPOLOGY: linear							
(xx) SEQUENCE DESCRIPTION: SEQ 10 No. 36:							
GATCOAGTGG CATGGNGOOT CTCAGTGGAA CCAT			34				
(2) INFORMATION FOR SEC 10 NO.39:							
(i) SEQUENCE CHARACHERISTICT:  (A) LEN TH: 150 Fase parish  (10) TYPE: number would  (C) STRANDENERS: number  (C) I I I VV: 1 steem							

AND THE CONTROL OF A CONTROL OF A STORY CLARACTER AND A CONTROL OF A C

March CHM World Title March, Indiane

(A) DERGIN: IN DASP PAIRS  (B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
(xi) SEQUENCE DESCRIPTION: SEQ 15 NO:45:	
(XI) SEQUENCE DESCRIPTION. SEQ 15 NO. 40.	
ATGGCCTTCA CGGGGCCG GGGACCGGGC AGCCCGGHGG GGCCGGGGGG TGG	53
(2) INFORMATION FOR SEQ ID NO:41:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 132 base pairs	•
(B) TYPE: nucleic acid	
(C) STRANDEDMESS: single	
(D) TOPOLOGY: linear	
	•
(xi) SEQUENCE DESCRIPTION: SEQ ID MO:41:	
GATCCACCGC OGGTGCAGAC GSTGCCCGGG GCCAAACCS: SAMCAGCCCC GGCAACGGCG	60
GCACCGGCGG CAACGGCGCG AACGCTACCG "CGTUGGNGE GCUCGGCGGG GCCGGCGGCA	120
AGGGTGGCAA CG	4.34
Middle Oth WW (1)	1 31
(.) INFORMATION FOR JED 10 No:42:	
::: SEQUENCE CHARACTERISTICS:	
(A) DEMOTH: Pl Pase page	
(B) TYFF: namlei marid	
THE ADMINISTRATION OF THE STATE	
and the decade	
enson segrando se se a parton sos portes en entre en	
Was entropy to the Additional and Additional Additional and Additional Additi	,
to the first the second of the	

(i) S	EQUENCE	$\subseteq HARACI$	ERISTI	LCS:
-------	---------	--------------------	--------	------

- (A) LENGTH: 702 base pairs
- (B) TYPE: nucleic acid
- (C) STEANDEDNESS: single
- (D) TOPOLOGY: linear

### (x1) SEQUENCE DESCRIPTION: SED ID ND:43:

CGGCACGAGG	ATCGGTACCC	CGCGGCATCG	GCAGCTGCCG	ATTCGCCGGG	TTTCCCCACC	60
CGAGGAAAGC	CGCTACCAGA	TGGCGCTGCC	GAAGTAGGGÇ	GATCCGTTCG	CGATGJCGGC	120
ATGAAGĞŞGC	GGCATCAAAT	TAGTGCAGCA	ACCUITCAGI	TTAGCGACGA	TAATGGTTAT	186
AGCACTAARG	AUGATGATC	HATATGAÇGE	AUTOGOAGAG	CGTGACGGTG	GATCAGCAA;	240
AGATTTTGAA	CAGGGCCAAC	GAGGTGGAGC	CCCCGATGGC	GGACCCACCG	ACTGATGTCC	300
CCATCACACC	STGSGAACTS	ACGGNGGNTA	AAAACGCCGC	CCAACAGNTG	GTNTTGTCCG	360
CCGACAACAT	GCGCCAATAC	CIGGCGGCCG	GTGCCAAAGA	GCGGCAGCGT	CTGGCGACCT	420
CGCIGCGCAA	COCCCCAAC	GNOTATOGUS	ASSTTOATOA	GGAGGGTGCG	ACCCCCCTCC	•480
ACAACGACGA	CGAAGGAACT	GTGCAGGCAN	ÃALEGGCC(s.)	GGCCTTCGGA	GGGGACAGTT	540
CGGCCGAACT	AMCCGATACG	CCGAGGGTCG	CAMARIGACIE	#PAACCCAVAC	TTCATGGATC	600
TOAAAGAAGG	GGCAAGG <b>A</b> AG	OTO WANCOU	STOAKMAAG :	CONATROCTA	(COTACTED)	6.6.0
PROMOTER :	CARCE ADAAC	V.,C. J.CW.,;	THE WAY STEEL	f **.		i ().

# FAR INFORMATION FOR SEQ ME NO:44:

CILIUMNO THAFACTERING OF SCHOOL HELL WORLD FOR E TYPER HELLE TO ALL TO ALCOHOL TO ALL TO ALCOHOL STREET

WO 98/16645

277

CTGGGG	CGGCG	GTGGCATGGG	AATGCCGATG	GGTGCCGCCC	ATCAGGGACA	AGGGGGGGCC	240
AAGTCC	CAAGG	GTTCTCAGCA	GGAAGACGAG	GUGCTCTACA	CCGAGGATCC	TCGTGCCG	298
(2) IN	IFORMA	ATION FOR SE	EQ ID NO:45:	<b>:</b>			

### (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1058 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) DEQUENCE DESCRIPTION: SEQ ID NO:45:

COSCACGAGS ATOGAATOGO GTOGOCGGGA GCACAGCGTO GCACTGCACO ASTGGAGGAG 60 SCATSACCIA CTUGCOGGGI AACCOCGGAT ACCOGGAAGO GCAGCOCGGA GGCTCCTACG 120 GAGGOGTUAC ACCOTCOTTO GOUCACGOOG ATGAGGGTGO GAGCAAGCTA COGATGTADO 180 TGAACATCGC GGTGGCAGTG CTCGGTCTGG CTCCGTACTT CGCCAGCTTC GGCCCAATGT 240 TCACCCTCAG TACCGAACTC GGGGGGGGTG ATGGCGCAGT GTCCGGTGAC ACTGGCCTGC 300 CGGTOGGGT GGGTCTGCTG GATGCGGTG TTGCCGGGGT GGTTCTGGTG CCTAAGGCCA 360 AGAGGCATGT GACGGTAGTT AUGGTGCTCG COSTACTCGG COTATTTCTC AUGGTCTCGG 420 CHAMBITAA CAAGGOAS O TOTAS TORA CONNITIONS NATIONOS NO TOROCOS 4 40 ACALEGREET TWANDERS IN A SECURITIES IN SIZE TOTAL GET CLAUSE CERTICITATEA. 540 CCG MCCCGC GC WCCGCCC AACTTCGATC TOTATCGACA STACGGC C TACLECAGT was a grant of the control of the grant Pri Albini III. I sa na Albini III. Ilan biya na Amban banna na banna an banna na bangangangan × 1 Control with a war was a control of the control of

(2)	INFORMATION	FOR SEC	(1)	NO:46:
1 6. 1	T TAT CALADACATE T CALL	I UN CILI	3 6/	1.0

(1)	SEOUENCE	: CHARACTERIST.	ICS:

(A) LENGTH: 327 base pairs

(B) TYPE: nucleic acid(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

#### (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CGGCACGAGA	GACCGATGCC	GOTACOCTOS	CGCAGGAGGC	AGGTAATTTC	GAGCGGATOT	60
COGGOGACCT	GAAAACCCAG	ATCGATCAGG	TGGAGTCGAC	GGCAGGTTCG	TTGCAGGGCC	120
AGTGGCGCGG	C0C3020666	ACIGORGO	$V(G,\mathcal{A},\mathcal{A},\mathcal{A},\mathcal{A},\mathcal{A})$	CATOCOCT FO	TAAGAAGCAG	180
CCAATAAGCA	GAAGCAGGAA	CTCGAUGAGA	TCTCGACGAA	TATTOGTCAG	GCCGGCGTCC	240
AATACTCGAG	GGCGGACGAG	GASCAGCAGC	AGGUGCTOTO	GTCGCAAATG	GGCTTCTGAC	300
CCCTAATAC	GAAAAGAAAC	GGAGCAA				327

### (2) INFORMATION FOR SEC ID NO:47:

- (i, SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 170 base pairs
  - (B) TYPE: nucleus acid
  - (C) STRANDEDNESS: single
  - (b) TOPOLOGY: linear

### THE REPORT OF THE PROPERTY OF THE PROPERTY.

TORREST THAT HAS A STORY HAVE A STORY OF STORY O

# THE REAL REPORT OF MALE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:	
GATCCGGCGG CACGGGGGGT GCCGGCGGCA GCACCGCTGG CGCTGGCGGC AACGGCGGGG	60
CCGGGGGTGG CGGCGGAACC GGTGGGTTGC TCTTCGGCAA CGGCGGTGCC GGCGGGCACG	120
GGGCCGT	127
(2) INFORMATION FOR SECTION 0:49:	
(1) SEQUENCE CHARACTERISTICS:  (A) LENGTH: 81 base pairs  (B) MYPE: micleic acid  (C) STRANDEDNESS: single  (D) TOPOLOGY: linear	
(x)) : EQUENCE DESCRIPTION: SEQ ID NO:49:	
AAGGRODDA ADRONDOND TORNDONAR BENAANDONG BOGADDEDA GAADDDODG	67
CGROGGOTOC GGCOTCAACG G	81
(2) INFORMATION FOR DEC ID NO:SU:	,
THE SECUENCE CHARACTERISTICS:  (A) LENGTH: 149 base pairs  ES TYPE: nacion to acid  () CTRAMPROMESU: Simile  () Tomology: I	
AL AR ALLES STEED TO ARTHUR STEED A SALAR TO A SALAR DESIGNATION OF	
TATE OF THE PROPERTY OF THE PR	

### (D) TOPOLOGY: linear

(xi)	S	EQUENCE DES	CRIPTION: S	EQ ID NO:51	:		
CGGCACGA	AGA	TCACACCTAC	CGAGTGATCG	AGATCGTCGG	GACCTCGCCC	GACGGTGTCG	60
ACGCGGNI	TAI	CCAGGGCGGT	CTGGCCCGAG	CTGCGCAGAC	CATGCGCGCG	CTGGACTGGT	120
TOGAAGTA	λCΛ	GTCAATTCGA	7GF-ACCTGG	TURATOTACT	GGTCGCGCAC	TTCCAGGTGA	180
CTATGAAA	AGT	ČGG TTTCCGC	STGGAGGATT	CCTGAACGTT	CAAGCGCGGC	CGATAACTGA	240
COTGCATO	AT	TAAGCGACTT	TTCCAGAACA	TCCTGWCGCL	CTCGAAACGC	GGTTCAGCCG	300
ACGGTGGE	TC	CGCCGAGGCC	CTGCCTCCAA	AATCCCTGCG	AUAATTOĞT.	GGCGA	3:, 15
(2) INFC	RMA	ATION FOR SI	EQ ID NO:52:	:			

# (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 999 base pairs (B) TYPE: nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: linear

# (x1) DEQUEETE DESCRIPTION: TR. 10 MO: 5.::

AT PARTY I STANDARD A CONTRACTOR AND A STANDARD F. C. CHARTSON A THE KIAT CHOROGER COMERCY ACCIDED A CONTROLL THE TIME IN THE TRANSPORT OF THE BATTER THE TRANSPORT OF THE BATTER. and the graph of the second of Anneas when his edge of the comment of the

GCCAACGGGG	TGTCTGGAAG	CGCGTCGTAT	TACGAAGTCA	AGTTCAGCGA	TCCGAGTAAG	720
CCBAACGGCC	AGATCTGGAC	GGGCGTAATC	GGCTCGCCCG	CGGCGAACGC	ACCGGACGCC	780
GGGCCCCCTC	AGCGCTGGTT	TCTGGTATGG	CTCGGGACCG	CCAACAACCC	GGTGGACAAG	840
GGCGCGGCCA	AGGCCCTGGC	CGAATCGATC	CGSCCTTTGG	TCCCCCCCCCC	GCCGGCGCCCG	900
GCACCGGCTC	CTGCAGAGCC	CGCTCUGGCG	CCGGCGCCGG	CCGGGGAAGT	GGCICCTACC	960
CCGACGACAC	CCACACCGCA	GCGGACCITA	CCGGCCTGA			999

- (2) INFORMATION FOR SEQ ID NO:53:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 332 amino acids

(xi) SEQUENCE DESCRIPTION: SEQ 15 No:53:

- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

Eet 1	His	His	His	His G	Hi	His	Met	Him	GIn I	Val	Asp	Fro	Asn	Leu 15	Thr
At a					Len										
All a	or r	1: 35	Val.	Th:	Va	1. + ±1	V ±1 1′+	1:	Ala	Ölir.		85.81: 115	Α	wil	j÷; :
Gla	Pro Cri	EIA	F1	150	V-01	11-	" lt	71.	A a	41 <b>.</b>	Temp e an	ė.	19 c.	=101	Thir
	ž Žiu i		•	î :	Fe File	*:	.7 1	11.7			Î1	i e		•	
1, 1	٠.				•				*						÷ .
		,						: .							

94

145					150					155	i				160
Gly	Gln	Pro	Pro	Pro 165	Val	Ala	Asn	Asp	Thr 170		He	Val	Leu	Gl y 175	-
Leu	Asp	Gln	Lys 180	Leu	Tyr	Ala	Ser	Ala 185		Λla	Thr	Asp	Ser 190		Ala
Ala	Ala	Arg 195	Leu	Gly	Ser	Asp	Met 200	Gly	Glu	Phe	Tyr	Met. 205		Tyr	Pro
Gly	Thr 210	Ar/j	He	Asn	GIn	61u 215	Thr	Val	Ser	Leu	Asp 220	Ala	Asn	G∶y	Val
Ser 225	Gly	Ser	Ala	Ser	Tyr 230	Ty:	Glu	Val	Lys	Phe 235	Ser	Asp	Pro	Ser	Lys 240
Fro	Asn	Gly	Gln	11e 245	Trj-	The	Gly	Va <sub>1</sub> :	11e 250	ĠĴγ	Ser	Pro	Ala	Ala 255	Asn
Ala	Pro	Asp	Ala 260	Gly	Pro	Orf	Gln	Arg 265	Ttp	Phe	Val	Val	Tip 270	Leu	Gly
Thr	Ala	Asn 275	Asn	Pro	Val	Asp	Lys 280	Gly	А а	Дlа	Lyn	Ala 285	Lèu	Ala	Glu
Ser	11e 290	Arq	Pro	Leu	Va]	Ala 295	Pro	Pro	Pro	Ala	Pro 300	Ala	Pro	Ala	Pro
A.a 305	Clu	Pr→	ři i i	Pro	Ala 310	Pro	$E_{1}$ , $\epsilon$	Pro	Ala	G17 315	Gla	Va.;	Alа	Pro	Thr 320
Fro	The	Thi	Hio	The	PΣ÷	GH.	Wa.		Lera Çiri	Pro	Aira				
. MF.W	MATI	C 11 F	· 13	EQ: .	D 145	:54:									
J		LEN TYL	THA NYTH: Firma ANTE	 r r	arib a Si	;									

... Which is a Market of Branch

- (2) INFORMATION FOR SEQ ID NO:55:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 15 amono acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS:
    - (D) TOPOLOGY: linear
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Ala Val Glu Ser Gly Met Leu Ala Leu Gly Thr Pro Ala Pro Ser 1 5 15

- (2) INFORMATION FOR SEQ ID NO:56:
  - (1) SEQUENCE CHARACTERISTICS:
    - (A) LENSTH: 19 amano acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS:
    - (D) TOPOLOGY: Linear
  - (xx) SEQUENCE (ESCRIPTIVE: SEC TO BU:56:

Ala Ala Med Tyo ProcArd The Gly Aspectly For Lem Glu Ala Ala Lys 1 6 16

do sily Arr

- (C) INFORMATION FOR SP. IN BUILDING
  - THE TRANSPORT OF THE PROPERTY OF THE PROPERTY

Ü

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 14 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear
- (xi) SECSENCE DESCRIPTION: SEV 10 NO:54:

Asp Ile Sly Ser Slu Ser Thr Sla Asp Gln Gln X'ma Ala Val 177

- (2) INFORMATION FOR SEC 10 NO:59:
  - .ii SE,MEMER MARASTERINII.SE
    - (A' LENGTH: 1) amino acids
    - de TYPE: amine acid
    - ( STRANDEDHESS:
    - (F" TOPOLOGY: Linear
  - (xil sequence description: see in worse:

Alle du un der ib der im der in New Le Walter

- . AF IMATI NO FEET, IN A FEET
  - . 1 Mary 1890 CF 1874 with the Lift 1990 F "Will EFN 2018 10" on the light con-

    - H' TYPE: amin'ami TOPANERINETTE

- (1) SEQUENCE CHARACTERISTICS:
  - #A) LENGTH: 15 amino acid.;
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ IN NO.61:
- Also by The Tyr March 4.5 to bet by. By The Am. The gry T  $\frac{1}{16}$
- (D) INFORMATION FOR DEC 140 NO:62:
  - i Broughte Characterius; Si
    - A, LERSTH: 30 amin a fids
    - .B: TYPE: amino acid
    - OC CIRAMDEDNESS:
    - it, ToPOLOGY: linear
  - .x: \* DENUENCE DESCRIPTION: UN. 1. NO.62:
  - Amplific Algorithm Acp Units The Algorithm Algorithm 3m + 15 = 15
  - termines Aut. Aug. Les. A. a. A. p. Cr. (Aug. March Frontier Charles)
- See INFORMATION FOR STRUCK IN NOTE OF
  - TO STUTENIE CHARACTERISTICS.

    A CONTINE CASSIST ASSESSED TO A SESSED TO A SESS

#### (LF INFORMATION FOR NEQ ID NO:64:

- (3) BEQUENTE CHAPACTERISTICS:
  - (A) LENGTH: 187 amine acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

#### exil reposm r benegition: Ds. 10 mars:

- Aun für Tur Met Lya Met Val Lya Der Fle Ata A. a Coy Leu für Ata
- Also Also Also Bly Also Also Also Also Vy Va. The Second Met. Also  $\mathcal{A}^{*}$
- GIV GIV Fro Val Val Tyr GIN Ref Gin Fr. Val Var Pho GIV Ala Pro  $\frac{1}{100}$
- Lem Er. Lem Asp Er. Als der A.a Er. Asp Vil Pro The Ala Afri Gli6%
- Less that it is best best four iver less Ala Arp from Aut. Val iver like Ala  $\frac{\pi^2}{4\pi}$
- Automorphistic Den Value Automorphist for the The Contact Act Art
- The Arabay But My Deatly\* problem Araba (Guiller Co., weg. Dec. Problem 11)
- en Angel Anton Comment of the Salah and the control of the

in service and the service of the se

- (G) STRANDEDDEDG: Bingir
- (D: TOPOLOGY: Linear

# (xi) SEQUENCE DESCRIPTION: SEQ 1D NO:65:

- Asp Glu Val Thr Val Glu Thr Thr Jer Val Phe Ang Ala Asp Phe Leu
- Cer 315 Let Acp Ala Fr. A. a Tim Ala Siy Thr. Co Ser Ala Val Ser
- Gby Val. Glu Gby ten for the Gby den. Als led fer Val. Asl bys Arg. 35  $$40^{\circ}$$
- oly Fr. Ash Ali Sly Jor Arr the Lod Len Asp. J.n. Ari L.e Chr. Ser
- Ald Gly Ard His Pro Asp Ser Asp 11e Pho Leu Aspo Asp Val Thr Val
- ver And Ard His Ala Sid the And Lou Sid Ash Jens Sid Phe Ash Val
- Val A.p. Val Sly  $\mathbb R^n$  less A.a. Sly St., Tyr Val Asn Arg Glu Fro Val 1.1 ↔
- and then All that he has a Arma May Appendix that the early byes been
- and Levi Ca. The Levi limit ring for the ring of the ring of Augustian ( ) of the ring  $(1000\,{\rm Aug})$
- 111
- to twee to the control of the contro

- Gly Pro Asp Ard Ser Ala Ser Lea Ser Lea Val Arg His Ang Arg Sin 2:,
- Gln Arg Asp Ala Lou Cys Lou Ser Ser Thr Gln 11e Ser Arg Gln Ser 35 40
- Ash Leu Pro Pro Ala Ala Gly Gly Alo Ala Ash Tyr Ser Arg Arg Ash
- the Asp Val Ard the Lys Sie End Met Led Val Thr Ala Val Val Led 65 65
- lend Tys Cys for Ely Tal Ala Inc Ala Ala Pro Eye Inc Tyr Cys silu 85 95
- CHe Lou Lys Sly Thr App Thr Sly Sha Alla Con Gli Lee Gli Met Sei 100 100 100 110
- Asprileo Ala .ve A nothe Admilipo Section to the Typ Typ to susp 119 125
- Cln Lys Sor Low Win Am. Tyr Low Ala Win The Ang Amp Lym Pho Leu 139
- Bor Ala A's One for der Des Des Des Dos Ala Fre Dyr Glu ben Asp
- Fig. The sec As a increase that it will be a first and  $\exp(\pi N a^{\frac{1}{2}} G \ln a)$
- As a field for the first transfer of the Astronomy Styles for the state of the field of the state of the stat
- The Cycle of the second of the specific terms of

WELL SEQUENCE DEPORTETION: DEC IN NOTE ':

The Ala Ala Ser Asi Asia Phe Sin Led ...: Oin Siy Siy ein Siy Phe 1 55

Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg Ser 20 25 30

Gly Gly Ser Pro Thr Val His He Gly Pro Thr Ala Phe Leu Gly

Value of the Alaskov Alaskov for lensely lie for the Gry Aup  $v_{a1}$  and  $v_{b1}$ 

The The All. Wal App Glv Als is . I've Ash Ber Als inc Ais Met Als  $\pi^{\pm}$ 

Amp Ala Jed Aan edy His die die dig Amp Vo. 120 vor Val Amm Trp. 190 – 110

Sty Pro Fre Ala 130

INFORMACIONER WORL, MODERNE

. H. OED GO HARA TERM, IT IS

7- 149147H: 10 one for and

THE EVILLATION AND A

TO PARTICIONE PERMI

A \* Pri I\* Yr ...m ar

entre entito for the region of entry and one particular

\* \*

Ger Ala G.y Mot Ala Arq Val Arg Arg Fri Maa Val Mot Fro X a Val 65 -70 -70 -75 -80

lle Gln Ser Thr Xaa Ile Arq Xaa Xaa Sly Pro Phe Asp Asn Arg Gly 85 90 95

Ser Glu Arg Lys 100

- (2) INFORMATION FOR SEQ ID NO:69:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 163 amin. a sign
    - .P) TYPE: arane alid
    - (C) STRANDEDNESS: single
    - (C) TOPOLOGY: Timear

1812 JEQUENIE DEGUKIPTION: DEG 17 NO:64:

Met Thr Asp Asp Ile Iou hou He Asp Thr Asp Glu Asg Val Asg Thr 1 10 15

led Thr Leu Ash Ard Pro Tim Ser Ard Ash Ala Lea Ser Ala Ala Leu 10 40

The App Mal Mal life section of a set of the Mal area of Alase  $\gamma$ 

THE THE CAR THE TOWN OF THE PARTY WAS CARDED A PARTY.

And the difference disease with a second of the second of

# (2) INFORMATION P P SEQ IN NOTION

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 344 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

### LXIA CEQUENCE PERFEIPTION: CFC IN NO. W.:

Met Lys Phr Val Asa Br. Leessurr - Sa. Ali Pro Ard Arg Ala Gly 1 5 15

Sky Ala Va. A a Glu Va. Tye Ala Shu Ala And Ang The Photology Ang  $\mathbb{Z}^2$ 

Den Pro-Old Fro Den Ala Met Led Ser Fro App 51 a 57y hed Den Thr 30 4

Also Gly Trp Also The low Arm the Text to a been Valide vide Valide Pro-

Are Gly Are Lyr Don Alm Vel A. r Ale Ale Vel Ale Ale Per Leu Arg where  $\mathcal{F}_{ij}$  and  $\mathcal{F}_{ij}$  and  $\mathcal{F}_{ij}$ 

The Top by Tal Ap A a New Hermon Medical Terr A a Alic Copy 20

of the Adams of Adams Adams to the African without a Ref. Africa Adams of the  ${\cal A}$ 

TV AN TERROR AND IT IN THE TERROR AND ALL DESIGNATION OF THE THE TERROR AND ALL DESIGNATIONS OF THE TERROR AND ALL DESIGN

tik ordan geria araban yanta baran baran baran ba

	211	:				V15									
A1a 225	Ala	. Log	ı Şiver	H2 si	H15	Les	$\Delta m_I^2$	Thr	Ála	17. 235		Less	ı Pro	iro	Pro 240
Thr	Arg	Gln	Väl	Val 245	Arq	Arg	Val	Val	Gly 250	Sor	Trp	His	Gly	Glu 255	Pro
Met	Pro	Mot	Ser 250	Ser	Arg	Ттр	Thr	Asn 26f	Ğlu	His	Thr	Ala	01u 270	Leu	Pro
Ala	Ал	: :::/::	Н1р	Ala	1*2	Thr	Ard Ar	Ista	Āļa	1.04	Len	Thr JRN	Gly	Leu	Ala
ine	H: 1 2 W	73 : 11	· .	7:11	A")	Ara 	Ange	Val	Ala	Ala	Ā1.5 300	Arq	Ser	Leu	Læn
Asp.	74.1	Altq	Z. L.i	AT i	liera Vito	V-4.1	11.7	Ælu		Ala 31	Tr;	5.1	Ala	Fr.e-	Thi 321
À. a	Ż. i	/v+ 1	M 1	11e 325	üly	VER	lip	i y -	130 330	41.	řs i	A. i.a	Tilu:	61 <i>y</i> 335	Glr.
Val.	CC:	Arq	-11:: 140	Agn	Pro	Chr	ely								
MEOR	MAT1	71 F	Par C	E. :	0 11-1	:71:									
T ;	751 [+1	TYI XII	GTH: A: a AM E	RACT 4P5 BID EMELY	enel alti	ne a i	11 (1,3								
	-)(-		113					* :	:						
1			· ; ·	r											
					A 1 .									ı	

Ars.	1 -311	i Are	i Tyr	n Len	i læti	Htt	- Авр	· (11)	i 41.1 9.1	r 1.	/ At	1 1 2	: Al	a -5: 95	1 Se
Thi	Gly	/ Gli	i Leu 100	n Met	Asp	Ātq	; Ser	Λ1,		г Суз	i Val	. Ala	1 Ala 11(		i GJ
Asr	Glr:	Tyr 115	Glu	Pro	Gly	Ser	Ser 120		, Arg	Trp	λla	G1t		Pho	Al.
Thir	Leu 130		ĀΙĢ	Aon	Leu	GIu 145		Lou	Fro	Aon	Ser 140		Thr	Len	. Mot
Asn 145	Ee:	:117	Thr	Aup	Len 150	Зιγ	Tæn	beu	Ara	G1 y 155		Pho	Ver!	Len	Pro-
110	*1. i	žir)	2001	1 + 1	;;;,	Be :	150	Env	Ala I''n	Ti.r	Letti	нігу		Ala 136	
. "1	ii	Ψ'.1.	A+ 4 190	Ā	$C, \gamma$	чгу	4. V	ihr iff	з. у	Syr	Ä.a	Effe	1'-1 2 +1	H:::	Len
At-a	lig,i	Ala 195	32 y	V-(1)	Aru		80a 200	. in 1	îî:	ŧΩγ		71.1 2.34.	Ala	.;	Пγ
Ptx	7i 710	1.63	Ere	Leni	Ar i	e. Pre	Pyr	h. Ţ	P*+* 1	Ali	Ata De	sily	· · i .	Val	Sor
M++ 2.15-	TI y	r i j	ist (	Āt:	$m_{\alpha}^{i}$	r:* ,	Ž\$ 1 1	Ty		Alu 23	74.	L. S.	řá je	71.	Ser F10
h	1.1	Živo \$		14. 14.	/s.:;	Εħ	. 13	1.1.	Λ. :	. 5	es 4 (	) . "ī		1:	2"001
	1.	: .	E .	Ens	Avrit.	. Y	٠.		17			. :		1.0	. ** .
21.1	<i>r</i> *	. il	-1 1	A2 1	≠s.°f.	1.	1011	.1		· · · · · ·		÷ . • .	e 1	e.: :	11.2
					*== 1 [										
					1 .										
				***	*,				!	;	,	e :			* .

310 Asp Val the Asp Val Ser And Tyr Pro Pho Pro Glu Leu Gly Glu Ala 390 395 400 Ala Arg Ala Thr Arg Lys Ile Gly Leu Gly Val Met Gly Leu Ala Glu 410 Leu Leu Ala Ala Leu Gly Ilo Fro Tyr Asp Sez Glo Glu Ala Val Arq 42C 425 430 Leu Ala Thr Ard Lea Met Arit Ard lie Gin Gin Ala Ala His Thr Ala 440 Ser Ard Ard Lea Alassian Chicara By Alasike Pro Alashe The Asp Ser Ar r Phe Alia Ar r Ser Sty Free Ard Ar r Ard Alia Shi Wall The Sec 4-15, Va. Ala Pre Tin 317 485 (2) INFORMATION FOR JEG IN NOTICE . CE JE,MENCE CHAPACTERISCIENT MA IFNGTE: 262 amin - anima (B) TYPE: amino acid " STRANDEDNESS: Finale T Films: Imeas M. JANUELL BASIL WALLS & LANCE sky Vak 10. Var leefar helefar helefar op 180, de on jûn oar ele

or the Article Court of Court

- The fro The Ala Ala Val out it fro bro Val bed bye Glu Gly Asp 110 and 120 fro Asp See The bed Ala Val bys Gly Led The Ash Ala Pro 115 fro Tyr Val Gly Asp Gln Pro Lys Phe The Met Val Val The Ash 130 fro 135 from 135 from 140 from 140 from 145 from 150 fro
- TO Val The Tee Ala Va, Ter Try Tee y Met y Ser Ala Fro Arq 196 - 206 - 705
- One Dry Les Gry Arg Pro-Ali The oly Fro Sly The Two Ann Les Val 210 215 225
- Value of the Lea May Ash Lea Art I is Lev. in Value of the Lea Ash 225  $^{\circ}$  . The Lea Ash 245  $^{\circ}$
- A a thirt character (SIA) of the object of any only.

INFORMATION FOR MET TO MET TO

D. TEMPORE CONTROL OF A DEMONSTRATE CONTROL OF

0 0 0 0

Val Arg Ser Lys Ala Pro Gly Ala Thr Val Ala Leu Thr Phe Gln Asp 65 70 75 80

Pro Ser Gly Gly Ser Arg The Val Gln Val Thr Leu Gly Lys Ala Glu

Glin

CONTRACTOR FROM IN NATIO

.Im GEQUEE W HARACTERISTICS:

TAL LENGTH: 604 amin' acids

Til TYLL: aftin and

(T) STRANDRIMEN : 015(10)

My 754 Dier: Linear

(RO) PRODERY DESCRIPTION OF THE HARRY WITH

Hily Ale Ale Val ter bounds . Ale Ale High the Lem Val Lem Inc Ale

The symbol of the Armster Companies for Armster Leville See

A CARLON CONTRACTOR OF THE CON

The grade and the Armed Committee of the first two Armada Committee of the first Armada Committee of the first transfer of transfer of the first transfer of transfer of

The second of th

14;					711					:::					160
G1:	ı 11 <i>+</i>	: Glr	ι <b>Λ</b> Σ.	Leu 165	Asn	Ser	Ğİy	Thi	Азј 170		ı Pr	· Ira	77.1	17:	o Ile
Ser	Val	. Ile	Phe 180	Arg	Ser	Asp	Lys	Ser 185	Gly	Thr	Ser	Asp	Asr 190		Gln
Lys	Tyr	195	Asp	Gly	Val	Ser	Asn 200	Gly	Ala	Тгр	317	Lys 205		Ala	Se:
ol t	Thr 210	Hie	Ser	аlу	Gly	Va! 215	11 v	Val	ijγ	Ala	179 1720	Cly	Asn	Asn	Gly
7nr 225	Herr	Als	1-m	Len	riin 1 Vo	Thir	77:1	A W	el y	301 235	110	liii	Түз	Asn	(41) (14)
Trp	Jer	11)	Àl:	7.4 f	шу	• • • •	·ii	Le a	A. f. Tria	Mer	Fig. 4	Шn		116 265	Thr
÷ +, 1,	$\Lambda \mathbb{N}_{2}$	1.5	1 : 4 2 (c1)	Ass)*	Pre	7,1	λla	11.	lttr	äni	OF.1	Der	Va i 270	Sly	Lys
Phir	Him	Ala 275	Gly	Ala	Lys	[ ] e <sup>§</sup>	Met Zed	GTy.	G15	Gly		Asp 285	Leu	Va l	Leu
Aup	Thr 290	Ser	Sier	11.75	m ç. •	Arj Pak	ŗ•	** <del>!</del> !!1	H:.		G1 ; 300;	Peri	TVI	Fr.	116.
701. - 14.	l.eri	A* i	11.3	5 ya	7. ; 3. ()	:	T.,	Ve	χ'· :	T 777.	1;;•		is. I	÷vi	That
* + +	415	·· } •	A 4	7-1 . 	šr:	4.7 :	l farra	Y- :		ni i	Νi :	I I	· i .	i '	iΞγ
liti		, is.	10040	A q	<b>:</b> .	***	* 17		.)(*	i t	i ·	1:	Ly.		]*t.
					÷		,								
		٠.		٠.											
		1 - 13-	1 1	х і. •••.	." 1										

1	: Alï	1 A	i žvi :	( * ( . ). ( .	' Ard	I day	. '.' .	l Ar	ι Δ:- ()	a Im	1119	, Hir	: Al.	15	i Asp
Gln	Thr	His	Gln 20	Asp	Arq	Let	н Наз	5 Hii: 25	Gly	y Cys	Arq	j Arq	Λ1a 30	Ala	Val
Val	Val	Arg 35	Gln	Asp	Arg	Ala	Se1	Val	Ser	Ala	Thr	Ser 45	Ala	Arg	Pro
Pro	Arg 50	Arg	H15	ĥτο	Ala	Gin gr	Gly	His	Arq	Ara	Arg 60	Vai	Ala	Pro	Ser
∰iy oʻ	Çîly	Arq	Ara	ñ: ı	F1 = 9	Hin	1 = 1	Hii ·	His	Val 75	G; r.	Free	Asp	Asp	A: q 86
Δrq	Asp	Ara	· · ·	Ala na	Terr	I ::	Ži sj		Thr	Gin	:: '	Ant	Gla	His Wi	Pro
Asp	Pro	His	Arit 130	At 1	ody.	9:	ř. Ta	71.* [ ** ] ** ]	Pr	· . '}'	At 4	. 1.	Arq Ilo	a. Y	Arg
Olγ	Ard	1 eu 175	Atsi	Arg	Vai	Int <sup>2</sup> F	M4 1.**	317	Ar q	Lesa	Cln	120	Ац	Ar j	Asp
Ala	Asp 1 (0	His	61.7	Ali	1:	. 41	At 1	*: ;	Ard	il y	1 m 140	Hili	Агд	ЯΞУ	Val
*in 145	Eis	Ara	OTV.	7.9	)	V (6)	Fire	: '	hi i	Ara	Verl	i i i ·	7. 7	V (1	Arq 161
· *, •	Ala	Hin	A7 1	lo i	17		At 4	A; i	· . !	A. i	Ā. s	Pro	. ;	üln 14.	ЗТу
Å (			213-1 14	Ast a	÷Ī.,		A	* .		Ar i	i en i			e e la composição	
7x 1 =	P 1.		ž ži	:	P	Ar ;		*. •	Ŧ.:	Ar i	÷ i	inspire	1 v /	G 1 '	Vil
								٠:			te :	* * ; -	X* :		
					:							1			:
	•		٠,	:		٠					)				. 1

290 (9)

Asm Arg Pro Arg Arg. 305

## (2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 580 amino acids
  - (B) TYPE: amine acid
  - (C) STRANDEDNESS: Single
  - .D: TOLOLOGY: lines:

wit depresses the pressions out in Moston:

ter Ala Vil til Cys Des Aum bly in: Thr bly Ard His Ard His Gly

And Cys And Val And Ala Ser Sty Tri And Ser Ser Ser Ann And Trip Cys 25 30

For the time also Argo types we Also the Light The Fro the She Also Also 45  $$40^{\circ}$$ 

ver the best WhatAri Ari its The Cyc Tips Den tri Alia Mal Gry Cyc Gen

And the Arm of the tro Wat Act with Der Alastes of Walls Ard Inc.

Dear Administrative Service of the Vall Vall Architecture of the Leader Committee of the Co

en de la companya de la co

elly let.	II. II.	Thy Mail 1	The Pre The Me	* The Property Pro-	Hiv Met
	ر خود ،		," (r ()	$P\Theta^{*}$	

- Val Arg Gln Arg fre Arg Alg Siy Met Leu Ala Ile Giy Ala Val Thr 210 215 220
- lle Ala Val Val Ser Ala Gly Ile Gly Gly Ala Ala Ala Ala Ser Leu Val 225 - 230 - 240
- Uly Phe Ash Ard Ala Pro Ala Gly Pro Ser Gly Gly Pro Val Ala Ala 245 250 255
- Ger Ala Ala Err For The Err Ala Ala Aon Met Pro Prz Gly Ser Valuet  $\frac{1}{2}$  with  $\frac{1}{2}$
- Gim film Val Alla Ata Sys to that about the last val Val Mee Lem Josephin (27) [40]
- Asp Levi Sy Ara Sun Ser Shi o. Shy ser G.y The file her her Ala Sh
- Gia Cly bea the few Thr Asn Asn Him Val file Ala Ala Ala Ala Lye 305  $$\tau_{\rm CS}$$
- From the Lemma y for the Lie Lie Lie The The Wall Int the Sec As: 326
- (C) V At 1 The Als Fr. Ph. The Vil. Cal. Cy A + Alg Fr. The Dec App. (44)
- where e = 1 is Val. And Val. i.e. any two there is you exhibit through the green e
- In a structure of the character was an expectation of the first state of the  $A_{1}$  ,  $A_{2}$  ,  $A_{3}$
- the state of the significant of the sale and 
486 .**;** (; . Val Sin Val Thr Ash Asp 1ys Asp The Pro Siy Ala Ly. 118 Val Glu 500 505 510 Val Val Ala Gly Gly Ala Ala Ala Ash Ala Gly Val Pro Lys Gly Val 520 Val Val Thr Lys Val Asp Asp Arg Pro Ile Asn Ser Ala Asp Ala Leu 530 5 4 5 Was Ala Ala Val Arq Ser bys Ala Fro Cly Ala The Wal Ala ben Thr 1.15 The Gin Asp Ire Ser Gly Cly Ser Asi Thr Vil Gin Vil The Lea Gly 1,717 5 (°) las Ala Ghi Blin ENEMPHATION FOR SEC 10 NO:77: : GEQUENCE CHARACTERISTICS: (A) LENGTH: 234 amin acrids (B) TYPE: amino acid C. STRAMBERMENCE SELECTION TOPENOW: Jane 8 AL CALIENTS PER WIFTI BY SECOND WHILE Met Aust Syn (Min om Star Ava Mar Mar Alberta de Sarras e Lear Carlotte) olympic was en Alexand In Section (1995), the member pyrige . 1. of the first of the congruence of the energy of the appropriate the con-

- Val Oir Gly Asp Ach Fre Ser Val Lya Len Ehe Asp Asp Trp Ser Ash 115 129 129
- Leu Gly Ser Ile Ser Glu Lou Ser Thr Ser Arg Val Leu Asp Pro Ala 130 135 140
- Ala Gly Val Thr Gln Leu Leu Ser Gly Val Thr Asn Leu Gln Ala Gln 145 150 155 160
- By The Glu Val He Asp Gly He for the the bys He the Gly The 175
- Lie Fre Ala Ger Cer Val Lyo H t Lea Amp Fre Gly Ala Lyo Ger Ala 190 - 190
- And the Ala car Va. Try lie Ala Cin Amp Gly Ser His His Leu Values  $\mathbb{R}^{n}$
- At a Ala for the Arg Low Sty Ser , yeller the Sin is . The Sin Ber B10  $$\rm PH$$
- lys Trp Asn The Pro Val Ash Val Asp 225
- TO INFORMATION FOR DRO IN NOTIFIE
  - ET DESTRICE CHARACTERISTING:
    - (A) LENGTH: + c aming a left
    - "P' TYPE: amin's reid
    - SCY STRAMS EDMENC: Sale: --
  - REPORT OF THE FREE PARTY OF THE PARTY.

  - \*

  - The second of th

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 69 among acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- OMIC SEQUENCE DESCRIPTION: SEG ID NO.74:
- Vai fro tro Ala Fro I: Jon Pro Fr: Lew Fro Fro Ser Fro IIe Ser I to 15
- Typ Ala Ser Er. Fro Ser Ev. Co. Let Dro Typ Ale Bro Pro Wal Ala 20 .do .do .do .do .do
- Fro G.y Fry Fry Motors Pro Det. Asporter Superior Fro Algebra Pro Pro 3%
- Loss Pr + Tyr Ser Thr ProcEre Cly Ala Pres Let Fre Pro Pro Pro Pro 50 60

der bro bro bed from

- TAL PURCHMATION FOR SEPTION 11 HT: 45:
  - THE PURNISH CHARACTURE OF COR.
    - As DENOTH: Street Court
    - "Ex TYER: andrewers in
    - CONTRACTION OF THE
    - In This Could be a see as

types at the second of the sec

Market Carlot Control of the Carlot C

化油油 医二甲磺胺二甲二甲甲甲基甲基甲基甲基甲基甲基

- Gly The Val The Asp Er Ash Giv Val Mai Leu Thr Ash Ash His Val 85
- Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Va! Gly Ser Gly Gln
  100 105 110
- Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala 115 120 125
- Val Dou Cln Len Ard Gly Ara Gly Cly Len Pro Ser Ara Ara 110 Gly 130 140
- Ny My vah Ala Val Gly Glo Fra Val Van Ala Met Gly Ass Ser Gly 145 150 150 160
- by Tin U, Ti, the Ervary Ala Val Free Try Arq Va. Val Ala Leu  $10^{\circ}$
- Giv 3In The Val Gle Ala for Asy for Loc. The Giv A,  $\tau$  Gle sile the 180  $^{\circ}$  190
- leu Aon Gly Les lie Gin ine Asp Ala Aia He Gin Pro Gly App Ser 200 - 205
- Ala Ala Ser Asp Ann Phet Clin Leu Dez Ger Sty Wey elin Sty Phet Ala 275 - 237 - 238 - 248
- The Barotte of Ada Ala Mathanesia, Ala Marata, din Anglie Anglie ang Tin
- The second and a limit will be a second of the second of t
- THE WAR MALE AND ADD AND THE WAR DESCRIPTION OF A PARTY OF THE WAR DESCRIPTION OF THE PARTY OF T

- And the first section of the first section of the s

- (2) INFORMATION FOR BEQ ID NO.81:
  - (1) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 205 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - x. SEQUENCE DESCRIPTION: SEQ IN NO.81:
  - For Pro Lys Pro Asp ALs Gin din din din 5.9 Ms. if (7). For two limits  $\frac{1}{2}$

  - The type Gly Lee The Ster Well His Wel Ale Well to The Let U.y Lye 36
  - Val Asm for Lem 1et GLy 11e Thr Lem A. (Asy Val Asm V.) Arm Ala 50 -
  - and the Lea Ala Ala type by very eye by the type Arr. Arr. Lea the stry are
  - Min. Fire Pho Ard Milesth ofly Ag. Ach. Look for Min. Lyu. Discher App. 80
  - with the the Appellant styling to the transfer to the Theorem Angles Angles Weight for the transfer to the tr

2. 4. 5.	1117 1	1	CHARAG	1 1 1 1 1 1	 
	. I I .	12.191 2.	21/1/2015	1 15 15	 

- (A) LENSTH 286 aman arran
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

#### (xi) OROJENOR DESCRIPTION: DEC ID NO:92:

- Thy Ampuder The Trip Alia Alia Ampudin Met Alia Ara Giy Phe Vaji
- Let Giv Ala The Ala Siv Art The The Den The Giv Glo Gly Let Gln  $_{\rm c}$  . The Giv Glo Gly Let Gln  $_{\rm c}$
- Him Ala Amp sily Him Sor Less Less Less Amp Ala Thr Ash Pro Ala Vaf 35 40 40 45
- Val Ala Tyr Asp Pre Ala He Ala Tyr GD. The Gly Tyr The Kaa GDu 50
- der Gly Les Ala Ard Met elys Gly Con Arm Fre Glo Arm The The Phe of
- Type the The Val Type Ach all the Type Val Alm Fro From Glu  $\frac{1}{26}$
- Arm The App Free God Sty Vi. Let. Gly 60y 15. Two As a Tyr Him Ala
- At a Thr. Fig. 415 Ar a Thr. A m 5 5 3 a a vi 5 1.8 de r A a der 1.9 Van 115 - 126 - 128 - 138 - 138 - 138 - 138
- Also Met des Also also as part we have the Met to be a constant of the second of the
- in the control of the

\* \*

.730 1. Ala Gla Ser Win Val Gly And Gly Pha Gly And Gly Ing Post Gly And 245 250 Arg Val Asn Ile Asp Pro Phe Gly Ala Gly Arg Gly Pro Pro Ala Gln 260 265 Leu Pr. Gly Phe Amp Glu Gly Cly Gly Leb And Pro Kaa Lys .7 f. i ACT INFORMATION FOR DECOME NOTE OF . DEPENDENCHARACTERISTICS SAP IEDOTE: 10% aren's aring (P) TYLE: amin. a rd pri Sipandeporovi single 📍 of) Tofall Stranders SWIT SPONGERS FOR BIFTING OF THE BUTTERS The Type the Bir Ale Length our Art Chille Win Arm "He Pro The Ala Ali Gib Tib Tive Val Ana II- Mia Val Lyr Ib Tip Ser Gli Asy Deposits of an including the Annexistation of the Annex of Ministry Annex In the first of the first of the Area of t

And And Per Ly And the His And And div dry Ard Iss. Let

#### (2) INFORMATION FOR SEQ ID NO:84:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 107 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (10 TOPOLOGY: linear

# with sequence becomes retroit of, if notweet

- And All Arg. Within a Life Arm that the More Park Web Valueyor for the  $\Gamma$
- Maj The See Ile Met Ala Cly Sty He Maj Maj Tye In Met StrePre 45 45
- Val Val His Say Ase in Terrir Lan Asp Fro Marcher Ala Fro Made to the Control of - We first the atomic duming the Xee box box box Art. You for Xee Arg. we see
- We will be supplied to the supplier to the first the supplier to the supplier  $\mathcal{X}_{ij}$

ing natural density of the second of the sec

PCT/US97/18214

231 Ang Ang Ala Lea oliv Les Sin Ala Pr. Ser Val Val Xas Ang Sin Gly 35 40 45 Val Lys Glu Pro Leu Xaa Thr Gly Ile Lys Ala Ile Asp Ala Met Thr 50 55 Pro the Gly Ar: Gly Gla Ar: Gla Lea He the Gly Asp Ard Lys Thr olly lys Acm And And Less by And Thir last the Der Ann Glin And Glin 3.3 Lea Gry Var Ard Trp ity Pro Ard Ser Ard Cys Ald Cys Val Tyx 125 Add Gry Har Ara Ala Ara Ara Cly Thr Tyr Hi. Ara Ara 116 INFORMATION FOR SECTION NOTES: -. SEQUENCE CHARACTERISTICS: (A) LENGTE: 11 amind accor 180 TYPE: amin acid of STRANCHPHESS: ofhice dh formbosy: Lucar SEC. REMOVE CENTRED TO USE USE TO SHARE TV - g ABB CALMER NORTH TO CONTRACT OF SERVICE CONTRACT.

And the Con Kin div.

#### (2) INFORMATION FOR SEQ ID NO:87:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 103 amino acids
  - (B) TYPE: amino acid
  - (C) STRAMBEDNESS: single
  - Hit Topology: Limear
- MI, SELUENCE LEGISTRATE OF THE NEEDS
- Mer Tyr Ard Inc A.a Sys Ard Thr Lee, Met Lee, A.a A.a Sys Lie Lea L
- The Ala Pro Val Fre Asp Tyr Cyr Trp Tyr In 6 GV 3 in Fer Inc Asp  $\alpha_{\rm S}$
- Fro Ala Tri Oly Pres Act, Pris Asp. Pres Typ. Thr. Type Hr., Acq. Acq. Phot. Pres. 198
- Him Arm July Johnson (1984) All Thomas Arm Arm Indiano Lydro of the Arm July 1990 (1994)
- the Dearest Court Table 1 of the growth Alaba te gas tak
- er Atrana vy 4. 19 k.a. 1

Ala Asp Gin Ala Ari Ala Giv Giv Iro Ala Ard Iro Irp Arg Giu His  $\mathbb{R}^3$ 

Ser Met Ala Ala Met Lys Fro Arg Th: Gly Asp Gly Pro Leu Glu Ala 35 40 45

Thr Lys Glu Gly Arg Gly lie Val Met Arg Val Pro Leu Glu Gly Gly 50 55

Giv And Lee, Va. Va' G's to . The lee Asp Gue Ala Aia Aia Leu Gly  $e_{\rm S} = -2$ 

Asy To bealty, by You The Fer

par information rander. In higher

CALUTARITARE MARATTERITIES:

JAN DENGTH: 95 amount another

obt TYPE: amin: acid

FIG. CIRANU EDNERU: Single

off TOPOLOGY: Linear

(x., JEQUENCE (#GJPIPTION: JB) 15 Mo:89:

the Arg Wila Asa for less Asa. On the Ala, thy Aon free the Ara the  $\frac{1}{10}$ 

the try Art for the unit on the approximation that the the the Academy

is the size if the second of 
(X1) SEQUENCE DESCRIPTION: SEQ 11/ NO:90:

Met Thr Gln Ser Gln Thr Val Thr Val Asp Gln Gln Glu Île Leu Asn 1 5 10 15

And Ala Ash Cla Val GRa Ala Ray Mer. Ala Asparan in s Thr Asp Val  $_{\rm SC}$ 

From the from two teles to a time was Kasa byte Adm Ala Ala Sin, Oin 3%

X.: V.I. Bep. A. A. a. Rep. Ach. bet. And. God Eve. Feb. Alia Alia Oly. Alia (A) 60

in the Arms of Armstein Almanda in the Lambda Armstein A

, yr sily "Lu Vi. Asp Chu Sin Ara Ala Thr Ala Leu i Arm Asp thy  $_{\rm BC}$ 

Fig. alpoint Wall tim Also act the All Sig Also Velough City Asposer for the Company of the Comp

Not All and less than Aug. The Errowers Val. Also What All a Gly Glucia a second of the second of th

An ine Met aug in 1965 in www.www.inining.Adj. 1965.

odnostvi Albus i događaja Buo Pearlo v Alborovije sediniha Maa Tha 147

The The West of Many

\* \*\*\*

ACT INFORMATION FOR JEET IN NO. 9.7:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 263 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

DOI: 100.00EDOR TENCRIPOLORI DEL 100.001921

The Algebra Met Ser Val fine Algebra Wy dim Algebra Minise. The Algebra with  $\frac{1}{2}$ 

Fig. Va. And Va. Alia Alia Alia Alia Alia Lyr Grouthir Alia The Gly less that  $\mathcal{C}$ 

Vil Er. Pr. Pre Val lie Ala Un Afri Ari Ala Sin Lei Met lie Lei  $\Theta_{\rm F}$  .  $\Theta_{\rm F}$ 

The Air The Arm Dec. Lem Six of Air The Brev Air Le. Air Val Asmospherical  $\Phi$ 

Win Alastin Two sely of the Net Sep Alastin Ang Alaska. And Med Thomas

The Charles And the Annalysis Analysis of the Shares and the Share

s , and As . Fig. (a) CW to the contribution of the scale x - y and the scale x

and  $\mathcal{H}$  is  $\mathcal{H}^{(0)}$  and  $\mathcal{H}$  
210 . 1+ . . . . Ser Ser Sly Leadly Sly Sly Val Ala Ala Ash Leadly Arg Ala Ala 230 235 240 Ser Val Arg Tyr Gly His Arg Asp Gly Gly Lys Tyr Ala Xaa Ser Gly 245 250 Arg Arg Ash Gly Gly Pro Ala (2) INFORMATION FOR SECTION 129:93: A: SEQUENCE CHARACTERISTICS: (A) LENGTH: (A) amprovacions (P) TYPE: write as 11 (A) STRANCEDNESS: Single

TOVERNEY TOTAL COMME

(x), DECTENCE DESCRIPTION: CEC ID to 1931 My Ser Tyr Sty Miy Va. Fir to See Fine Ata Miss Alla Aspe Mis Sty here Atlanta 1990 en Dan Ser En Lour Fe. Mystimes Inc. of Der Thr April 1984 William Charles and the Property Charles and the Charles

- Ala Eric Ala Pric Arg Pric Lyr Ero And Proc Tyr Gly (In Tyr Gly Arg 165 176
- Tyr Gly Gln Tyr Gly Gln Tyr Gly Val Gln Pro Gly Gly Tyr Tyr Gly 180 185 190
- Gln Gln Gly Ala Gln Gln Ala Ala Gly Leu Gln Ser Pro Gly Pro Gln  $\frac{195}{200}$
- G'n Ser Iro Sir Pro Fro Gly Tyr Gly Ser Gin Tyr Gly Gly Tyr Ser 210 - 220
- Ger Der Erit Ger Jim Ger. By Jer. By Tyr The Ala Him Free Ero Ala 240 \$235
- Gli. Fro Fre All reln ser G.y Ser eln Sha Ser His Gli elly Fra per 145 255
- As a Gy The Gly See Sim Ata Gly See Ata Fro Val Ann Tyr Ber Arn  $\mathbb{R}^{26}$  .  $\mathbb{R}^{26}$  .  $\mathbb{R}^{28}$
- Proder Cly Cly Cha Can Da for Ser Proday Sty Ata Pro Vaco 200 300

### -. INFORMATION FOR SE, ID NO:91:

- CONTRACTOR OF SEARCH SERVICES :
  - A' IENSTE: Softman carry
  - (B) TYPE: part for inf
  - TO STRANGE PROPERTY OF THE PARTY.
  - The Party Transfer of the fire

that is to set out the set of the

A N	M W	Y. 1. 1	$N_{i}$	7:	77 7	WST.2	W.: :	• ; •	32	·:: '	7.	$\mathcal{M}_{i}$	• • •	AT 14	$\cdot, :M$	i :
ůž.	TTKT	'GG I	with."C	1175.	iji u	MCT.	És									
(2)	INFO	RMAT	'ION	FOR	SEÇ	1D 1	10:95	);								
	(i)	( <i>1</i> ) (1)	OUENC ) LE TY ST ) TO	NGTE F#: RAUF	i: 16 amin Eleik	в ал (с. 40 (30):	nine De Simi	acid	ls							
	( × : 1	71E),2	115,1461	E DE	2.7E:	DTII-	11: :1	F .: :	E No	; a6;						
	Me-+ 1	1 73	Most	* .1 1		. :	110	.v. a	Ž1 1	17	1.8.		Ala	/v) 4	74. a	Als
	Пo	87	Fil i	A	ĀΙα	j.". 1	Üу	Tal.	Tr. +	11+13	11:	Mot	$K_{ij}^{ij}$	Jly	Gly	Pro
	V a 5	<u>, ()</u> ;	77.1 31.		:·	ş: 1.	1 .	** 4		11.00	eliy	Ala	1 ; . 41	, e <sup>2</sup> 1	£x	Letu
		1:	Fi i	e . ;	Å. i	Ε;	Ai j	* * *	Ť	11.5	AΓι	A. i	1.1.	Îm i	Tir	200
	. 12 . 1		26 4.	:	100	• • •	÷. :	1:	.= '-			1:.	ist. i		. y .	$i + \gamma^{\nu}$
	. '	1 .	9			·		1		1.1		F. F. F.	Ar r		Addi v*	ένε")
		· ·														Î în
	•												• , =		·, ·	i ve

1.3	T. F. Nov. 1	1; -	5.1	3 x	1.1177
1 * 1 /	1/1/4	11.	114 1	10 1.00	1 .1 . 1

- (b) TYPE: nucleic acid
- (C) STRANCEDNESS: Single
- (P) TOPOLOGY: linear

#### (xi) SEQUENCE DESCRIPTION: JEGS HE N 196:

· . = 0:1.10g1, 5.55	حادثيا الربعمين بب	trace of 11 b	a Maria	1-8 WORGOT	£.(±
- Archerant	ATTENATOR	1, 17, 17, 17 T	ently of Parkets and a sing		1.1
CACTTOAMID	$\mathcal{M}_{\mathcal{M}}^{1,1}(\mathcal{M}_{\mathcal{M}$	F. C. C. M. A. F. S.	ADECEMBRICAL	Arriveque	186
$v = X^{\alpha}(0, 1)$	07.12 X1800	· · · · · · · · · · · · · · · · · · ·	WASSIESTOR		745
Catematan.	$A \subseteq \mathbb{C}(\mathbb{R}^n) \subseteq \mathbb{C}(\mathbb{R}^n)$		AN WASTATT	AARETTATG!	. + 1)
COGCGACOUG	GCATCGTCCC	C ISROCIA IS	CONTACTOR	CH KITCCTCA	360
rerogador.	TOTAL ST.	. wateruly	TO TANKE	Q11 HOTCO:	426
CATCTONIA	(CAATTY) :		(1) 1 (1) (1) (1) (1) (1) (1) (1) (1) (1	""C"A WEC	430
PROPAGOT					£3±↓11
	CACTONIANT CACTOMACO CACTOMACO COCCTOTO COCCGACOCG COCCGACOCG	CACTOMAATT AN MEGA DOT CACTTOTACO COTTOACCACT COCCTTOTO A STOCKTTOM COCCCTTOTO A STOCKTTOM COCCCGACCCG GCATCOTOCC TOCCGCGACCCG GCATCOTOCC CATOTCCTAC COCAATOCCTC	CACTURATO ACURSENTO A DAGINGO CACTUANO DOTTACONOS ENGUARTOS CONTINENTATOS AUGUSTOS ANTOS COCCETUTOS AUGUSTECO A OUNCE NA COCCETUTOS AUGUSTECO A OUNCE NA COCCEDA O GOATCOTOCO O UNA CALCUTO O CAACOO O COCCEA O CALCUTO O CAACOO O COCCEA O	ACCESAACE ACCESACE ESTACEMENT CONTRACTOR CEVAACENA CACCESACE CACCESACE CONTRACTOR CONTRA	TOOGGERANT AND SHOW IN A PROPERTY OF THE SHORE THE RESPECT OF THE AND SHORE THE SHORE

## OF INFORMATION FOR MEDITINATED

WO 98/16645 PCT/US97/18214

130

Sin Arr Ala Ala Net Ala Ala Sin Net Sin Ala Val Fr Siy Ala Ala 90 76 80

Sin Tyr The Gry Lew Val Giu Ser Val Ala Giy Ser Cys Asn Asn Tyr 86 90 95

(2) INFORMATION FOR SEQ ID NO:98:

(1) SEQUENCE CHARACTERISTICS:

(A) JENSTE: 154 base pairs

(30 CYFE: modiero adid
(5) CTPANDEDNESC: Single
(7) U 491/OFF: linear

AN WARAST AG WANTE THOUTH AND WAS IN THE STATE THAT WAS CONTACT AND WAS CONTACT.

\* TARGEMATICS FOR THE II INSIDES

A DEPUTE CHARACTERISTICS:

OAS LENGTH: 'I ambies a fide

OES TYPE: ambe desire

STORAMEDUEDE: District

TO SHA FREE FUREER

(A) LENGTH: .m. feare pairs (F) TYPE: max.e. racid (C) STRAMDEDNEES: single (D) TOPOLOGY: linear	
(xi) BEQUENCE DESCRIPTION: DEp ID No.100:	
CUSTOCIUS A CTTUCASUTE A INVIGANA E LUGUMERUCE NO ESCADADA AU TRAINSAAÇOT	{ v ' }
TOWARDS FOR FATABLES A RETRIBUTE A DESARROSE TOTAL STRUCTUMENAS ADRICTOMORGE	îz iI
AGTOCAAAACA COOCA A COO GACAAACA TO CONSCIENT COO ACAAAAA AAAAAA AAAAAAAAAAAAAAAAAAA	186.
A MATERIAL COMPUNITA A CONTROL OF COMPUNITATION OF A CONTROL OF CONTROL AND CO	4.3
ACCOMMENDATE OF THE CHARLES OF A CHARLES ARE THE CONTROL FACTOR OF THE	.* •
A, INFORMATION FOR SECTIONATION	
AN LENGTH: 3-58 have paint AN LENGTH: 3-58 have paint par THE: northern and f ed off-Annellifour comple- 10 Topological lane er	
al delement three *later of the first te	
All about the analysis of the first transfer and artiful to the second	i
	*

WO 98/16645 PCT/US97/18214

	THE PROPERTY	AIGAGETT N	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	Construction of	G Angerengåre	*: ()
AGGTT COSST	Tarverond	a i ma saan	Contractive Contraction	GCTGA CCT	: community GG	720
TGATCGCCGA	GAACCGTGCT	GAACTGATGA	. TICTGATAGC	GACCAACCT(	TTGGGGCAAA	780
ACACCCCGGC	GATCGCGGTC	AACGAGGCCG	AATACGGCGA	GATGTGGGCC	. CAAGACGCCG	840
COGCGATGTT	TGGCTACGCC	GCGGCGACGC	CGACGGCGAC	GGGACGTT	Crabogarca	900
AGGAGGGGGC	GGAGATGACC	Validia in La	TENT TO TA	GCAGGCCGC.	· FIRSTONAGG	9 n C
Accordance	CANNA 12 12 12 12 12 12 12 12 12 12 12 12 12	fin il. W. W. C.	TALANAA	" TOOMS CAN	· .: o :::::::::::::::::::::::::::::::::	1020
Accidacecta	icantroca.	10 WYW) 1	· 17771 · 7w.	CTT. TOOTT 130	TOTAL USA AGA	1680
gestataco)	$\mathbb{R}^{N} M^{n+1}(\mathbb{R}^n) :$	TOTATOA	ACMT-137" T	UATHGCONAC	AATCACATGT	1140
· Wî  \Ac wa	Teleja i Tega	let Witt Mide A	AWARTT WE	THE BATCETT .	Lavis (CTT)	1
CINTAGONO.	30 200 100 <b>2</b> 0	05. 370 AAA	CCC 1 0 197A	AAACGGGCTC	CHETTGATGA	1260
nomentary is	WATT TOTAL	* T.	the contract of	Tancorocci	GCCAACTTGG	1320
ama, garagar	1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1	TENTEST +	Tartin Variation	Takijā vasi	DOADCAACC	1380
Maria Marcin	250 130 00 14 F	;::\;: <b>:T</b> (; \ \:::	. w.w.:	W.J.W.	GU MAAADAG	1440
**************************************	avi alice e		***************************************	* (.8. 10.2V.)	maionwri.	11.00
**************************************	2007 10 10 10	****	4. 1111A1 -1	MT I NI WI	m minning yan	1
* * * * * * * * * * * * * * * * * * * *	. :Zv: :::::::::::::::::::::::::::::::::	A M N D THE	17 31 + 47.1	artera	n m stæm	100
77 · · · · · · · · · · · · · · · · ·	X TALAY	a were	* * * * * * * * * * * * * * * * * * * *	·1 *** . 75 (****).	Towns own	:
11 Silv Mi. s	AND AND A	r, will be "	zana andara	to the DATE of Co.	Allogram.	1" 1 .
, · · · ·					11.0	•
* 1	. 1				e	* .
* 4.00	. (1)		* •	··. ** ** ** ** ** ** ** ** ** ** ** ** **		
8						

GCCGGCAACA	ACATGGTGGA	AWCCA, WGC	e engem	PMOTINGS.	CTOACACCA F	234(
GCCAAGGCCA	GGGACGTGUT	STACGAGTGA	AGTTCCTCGC	GIGATCOTTO	GGGTGGCAGT	2400
CTAAGTGGTC	AGTGCTGGGG	TGTTGGTGGT	TTGCTGCTTG	GCGGGTTCTT	CGGTGCTGGT	2460
CAGTGCTGCT	CGGGCTCGGG	TGAGGACCTC	GAGGCCCAGG	TAGCGCCGTC	Clicgateca	2520
TTCGTCGTGT	TOTTCGUNGA	- FUNCTION TO	GAGGAGGCGĞ	ATGATCGAGG	CSCGCTCGGG	2580
GAAGATGOOG	ACRACGT YM	TICCCTCI	TAUCTOTOGO	TTGAGGCGTT	COTGGGGGTT	2640
CA CAGBITT	ASTIMATION	$iiK(\mathbb{N},\mathbb{T}(\mathbb{N}),\mathbb{P}).$	Color DANG(e)C	GTUMACUCCA	FAGGTCGGT	2700
cennaghadara	TOTAL CONTRACT	· · : BOCMOTON	GROGATTIAG	TOGGTCAGAG	CGTGGAGTAC	2760
CCGATCATAT	TOSCHADINA	THATTO JOS	ricanting.	TCGTAGATGG	APTICACCAG	2820
GHT0EGCNCC	CASCIGITA III	Index CTT (27)	::::::::::::::::::::::::::::::::::::::	ATUMATTÖG	TOTAL	38×1
PTTTTTGČAG	SACTROCARD	COGCTHOGHE	"An Physical	COMPAGE (F)	CAPTAAGGCC	394¢
C2007900073	TEGCTGSTWA	1 Marrachari	n indefactivic	nchessecta.	acvastacaca	3000
BARSAAÇO :	NOTE: TO ST	* * *, *, *, * * * * * * * * * * * * *	or radio Weathur	Association of the second seco		30 In

INFORMATION FOR SEQ TO NOTICE:

## go. DEPUENTA MAPANTER BUTTODE

- (A. LEMATH: 1:1 amin anida
- .F. TYPF: andrew artis
- POLITICANI RIMERINI I INGGO POLITICANI PROPERTO POLITICANI PROPERT

WO 98/16645 PCT/US97/18214 134

Ala	СlУ	C.n	Ala	alu gņ	lart.	-11.r	Αla	r Ala	9()	1 7 1	Ьr	Val	Ala	A1a 95	ı Al
Δla	Tyr	Glu	Thr 100		Tyr	Gly	Leu	Thr 105		l'r o	Pro	Pro	Val 110		Ala
Glu	Asn	Arg 115		Glu	Len	Met	11e 120		lle	Λla	Thr	Asn 125	Leu	Leu	Gly
Glin	Asn 130	Thi	1:0	Alu		Ala Lie		Ä. т.	:11-1		Gla 140	Ty:	61 y	Glu	Met
Tri) 145	Ālπ	ČI.	As jo		74 150	$\Lambda_+^+,$	14	1900	H;	Ty:	Ala	Ala	A1-4	Thr	Ala 166
Tir	Al i	Thir	712 e1	ll.: Terr		1 - 11	ł i		31 a 170		A i	::	Hi	Met. 175	
51	Ala	·	31y 180	heet	lert.	d):		Ac. 4		A1.1	V.1.	1731	;1+. 190		ger
Asp	Thr	Aia 195		Ala	Arn		iem į		Alin	ASI	Val.	Free Å( 12	viln	Ala	Leu
6:1+4	01:. 21(	Desi	Alsi	ŝin		The		÷Î*;	The		Pr 227	11.0	1, 1	· 5'-7	Len
	Gly	<u>Dorig</u>	7:40		1:.* / 302			i	His	A: 4 234	1	1.7	:10		Arm . 40
1)		, 1, + +	∵, t	Alm .d	A. I.	A. 1.					Flar	in: te		(1) y (	
	M av		A. A.	1:.2	* 1		:		: 0	: ::	: y		٠ <u>.</u>	1:3.	A. 1
							• 1 .							,. 1	
••**															
						•								:	

Thy Cly General Gray Varies as Are Val to Fro And Pr. Tyr Val Met Gro

Pro His Ser Ero Ala Ala Gly 385 390

#### (2) INFORMATION FOR SEQ ID NO:103:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1/25 base pairs
  - (b) TYPE: numbers asid
  - (C) STRANDEDNESS: Single
  - J. Tapolatur linear

THE STREET PROPERTY ON THE BOX OF

SACOTAMA VA COMORCULTON AGRECOOMA CARRAGADO PROPORTORO A DECMAGADO. APOTOCOTO COCTOTO DE GUOGTOGATO PAGACTICATE DESCRIPTA AUTOCAACTA 120 ATTICUTT VA AGTICCEGO I ACCEATAG VA COT MOVALT COLLABORIA CONTRACON 139 FIRESTITE FREE TO WEST FREE FORWER ALCOHOLOWS TANDED TECHNIQUE OF POSTUAGEOG 240 AATTEGORES TA CETTA TE AATTAAT AATTAA TETOAKATEN DEGOREGEEN ET WETTEGORE 340 APARATA PARTET TERMENT NATIONAL APPENDIX TO SERVICE n er ny sa pengentah darih da namarayan sinasaya nagaraya TRANSPORTED AND THE PROPERTY OF A SAME AND A 4 24 TATRAFATT STANTAL ASA STANT - AA S STANTA AS AS AS TO 1..; . .

. : 1	$GOBIG_{\pi}\Lambda\Lambda$	125 132 x 25 1 x	NATURE OF	$(1K^{\times}h^{\times})^{*}$	Artigura raz	LatinAcona	1140
GJ	CGTTGCTG	CCGTTCGAGG	ACGCCCCACT	CATTACCAAC	CCCGGGGGGC	TCCTTGAGCA	1200
GG	CCGTCGCG	GTCGACGACC	CCATCGACAC	CGCCCCGGCG	AACCAGTTGA	TGAACAATGT	1260
ĢŌ	CCCAAGGG	CTGCAACAGC	TGGCCCAGCC	AGCGCAGGGC	GTCGTACCTT	TTT CAAGCT	1320
(3-)	GPGGGCTG	TGGACCTTO	TOTAL COOK	TUTGICGC	CTCAGCAACG	TCAGTTCGAT	1340
A 7	CAVACAVU.	CACATOTOR	TOWTO KOMA	a sperimen	ATGAGGAAGA	COTONATO	1446
ંત્રે	DOTT MAG	CLESTIAN COLO		TOWNST .	GAAACCOÇCE	CGUAAAACGU	1500
(1:1	rogadan	7/7/17. 2 mg/s 1 mg/s		1	1 st 1, Gi (CT)	TTT-CG-98TCT	1500
: 10	1 1111111111	.:::::::::::::::::::::::::::::::::::::	A TOTAL	7.000 2700	TENERAL CER	TAIMGATGAC	1.00
: `:	Wilats:	-1 71	$K^{1}$ , $M(1)$ : $\mathbb{N}$	$X(x,x,t): x \mapsto$	**************************************	TWOTAN	1450
M	TOTGACC	AGCCCCCCCCC	AAACCETTTT	TOATAL MITT	angga		

## SET INFORMATION FOR CFD IN NOTICE.

FIRE SEQUENT THARACTERISTICS:

TA LENGTH: 10th and the act. I.

Re TYPU: amin. artif Co Japann Elmenda

He T F 14 Wr linear

# er in the second of the second

Sometimes and the factor of the contract of the second

 $((x_1, x_2, \dots, x_n), (x_n, x_n), (x_n, x_n)) = (x_1, \dots, x_n) + (x_n,  

					11.0	P170 11	 100	$LG$ $\alpha$
	1:			1.0				

- The Asn Arg Thr Glu Leu Met Thr Leu Thr Ala Thr Ann Leu Gly
- Gln Asn Thr Pro Ala IIe Glu Ala Asn Gln Ala Ala Tyr Ser Gln Met 130 135 140
- Trp Gly Gln App Ala Glo Ara Met Tvr Siy Tyr Ala Ala Thr Ala Ala 145 - 150 - 155 - 166
- The Ala The Stu Ala beh length the 31r Asp Ala 95 (bet II) The 145 ( ) 120
- Now the Giy Novi to Lie . To a function Was with wait with Clu Tin Alacine . Fig. 188  $_{\odot}$  . The  $_{\odot}$
- Asy The Ala Ala Asa Ash Ash Israelle Ash Ash Ash Ash Ash Israellen
- Glin Glin bed Alia Glin Free Alia Glin Gly Va. Val Err. Feer Nee Ey. Leaz 216 - 227
- Fig. 61y Dev. Trp The Alm Vel Der in the Fig. 10; near Fr. 10, show Aug. 2.0  $$\rm PP$$  . P40
- Value det det fle Alia Ash Ash Hill Mit Ver Met Met Gly fmr. Gly Value  $\mathcal{M}^{\alpha}$
- . The first sense the decomposition of the first sense that the first sense is the sense of the
- All Art Pin Ara Jan is a few Alastic list Arm soy its High Ara Me
- and also as a significant distribution of the significant and significant and significant and significant signifi

114.1	 1.04 2 1 (2.1	

- (t) STRAMSEDNESS: Single
- (12 TyPOLyW: linear

## (xi) SEQUENCE DESCRIPTION: SEQ 10 NO:105:

ACTICACIOS AGAMBATEAN	TOMOGRAPIO	THEMAN	GROTRAGACA	AC DAACCAC	$\hat{\epsilon}_{i}$	
METONANO MONANTA	WASTYGWG		#GCCCCCGAA	PMHAACCCG	137	
COMANDOM ADOINTE		94 ( N. W.M. R.	CCCTGGCCCCC	Y VITTONAGO	180	
STOATE TO TELL AND THE	727327(17VII)	Washawar II.	CCCCGAGGAT	ATTOAGACTG	240	*
STANGACLIC MADAGTATIC	ALAS MATOS	may babas	ar a maga ng		*00	•
Arustons Armint Sound	GENERATORS	CONTRACTOR	CCAAGACTCC	COACGATTON	360	
- ATOGOGU TO DOMOTOGICA	THATHTGTG:	7, 7, 5, 1	- T. MO CO DAT	WOWLTO FESTE	4270	
FIAACALMAT (MIGGORADIN)	AACCOASTA	***; ***; **; **;		GC PJAGCCAA	480	
GENAAGGT FIC ATTAACATGA	tolomos from titte	. mm -a.a. a.a. :	TOTANGAN.	AC MO JOSOG	540	
THE WINDSAM AGOST YOUTH	I,MARTINA	)   W W Y	-Wall state	With Collaboration	$\hat{c}, O\hat{c}$	
- 'AC MARKETT - ETCHAA Mort	"WAATET II	13.11 000000	Amil em Aamil	**************************************	tiere	
19450 Week AA LAB WITT	:=1:::		1.4.11.2	Tay Windaa	· ,· .	
TO THE RESIDENCE OF THE PARTY OF THE		***	AAAA DEAAA SEE	iá il jan ik		
e ko sa se sakaalat s	7-871. v 1, .	· (P)	24 274	hama in	1-1	
** ** ** ** ** ** ** ** ** ** ** ** **	+ • <sub>0</sub>			· · · · · · · · · · · · · · · · · · ·	• •	
				* **	* .	

Al-GAGAGCTC	AWA! COOP	Aro (coco)	CAP POR WORL	. ::T*,V	CCUAAGGTYCG	144
ACGGCAATTG	GCTGATCACC	AAGTTCACCC	CGOTTTAGGT	TGCCGTAGGC	GGTCGCCAAG	150
TCTSACGGGG	GCGCGGGTG;	CTGCTCGTGC	GAGATACCGG	CCGTTCTCCG	GACAATCACG	1560
GCCCGACCTC	AAACAGATCT	casacaarat	CTAATCGGCV	GGGTTATTTA	AGATTAGTTG	1620
CCACTGTATT	TACCTĞATUT	TCAGATTOTT	MOUTGGATT	TACKTTO 3CC	GCAGGGCGGC	1680
TGGTGCACTT	TONATORNIC	CHEOTE WELV	TTTTTACAGAA	TTTGACCTCT	TUCCIGACCTT	1740
oll : Jacoba	AICATT	TTARTAC	80.5446(0.004)	AGGATIATIN	AAGTGGTSGA	18140
TTORGER	mry 21.V . 2.1.	· · · · · · · · · · · · · · · · · · ·	= } !\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!	WCGOVEEN.	CGGGTTGGGC	1860
orean a	3000 COM	A WE DOWN	+7v: 577,110 i	ADDIA VIGI	TTTeSecce	1920
CIPCUS CUTTY	CAGTOGGTG I	THURST	(MC,MUU,M)	T ZITTINATA(*	frr Grager	1980
GUSTOTGATS	OTGGCGG MG	TENERAL TENER	gerrander:	ATOAGCGTCA	n isneegga.	2040
DOWNAG THE	Accordant	Tarley Comments	TOOTSTOOK	GCCTACGAGA	CGGCGTATGG	2100
CENTOWNS IN A	. 5 5 7 7 7 5 7 5 1 4	I WTM 11.	MA STEETE	CANCTONTON	TTOTGATAGO	2160
«WAVAAAAA	TPSWETTMAA	WW. Control	*9.7 · ; · : TFC	AAGWAG YW	AATACGGGGA	. 220
(WIOI333)	7474-74	emeration t		jimaquunun.	3, 7 7 7 7 3 7 5 8 7 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2280
18.4	11.1 41.1	Print Strain	As CHARACT	Marin Tolar	`` : ``	2.44.1
7. 1 2 1	*********	7. 10 122 175	With the section	thought with	i "A. WA WA	. 4'.
ne Transfer			7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7. 7	* *** ***** ***** ********************		7461
Nº 1 'L 1 AVA	and the second	·	V			
					. * : *	
					*	
		•			* * * * * * 1	
	•					

INCORRORD CTATALORS AGRIANTS

30.

#### (10) INFORMATION FOR SEQ ID NOTICE:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 396 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: Linear

### THE SEPTEM HE ENDINE HE AT MELLER !

- Val Val Amp The Gly Ala Lembritis (GD: The Aun Jen Ala Ang Met T
- Type Alamely I: Sly Ser Alamer her value at Ata Ata Ata Ata and the Cry  $\mathbb{R}^n$
- App Ser Val Ala Ser App Leu Pha Cer Ala Ala Jer Ala Phe Gin Ser 35 45
- Value Value Topology and Observation with the displaced Gap the GDy when Ber Alfa GDy section
- Let Met Val Ala Ala Ala Ser Er Twe Va. Al. Err M t Der Val Three  $\mathcal F$
- The UV sector and the Section Annual Equation ( ) and the Variable Annual Annu
- After the constant that the contract of the section of the constant of the co
  - The first term of the second s
- and the second of the second o
- $\frac{d}{d} = \frac{d}{d} \left( \frac{d}{d} + \frac{d}{d} \frac{d}{d} \right) + \frac{d}{d} \left( \frac{d}{d} + \frac{d}{d}$

J.I.	31:. 21-		Ala	··i:.	i: .)	T1.:		73.07		111	1:0 270		- Aury	· 11.1:	: Letti
Her 225		Leni	T.T.	1.ys	Ala 230		Ser	Fro	His	Leu 235		1°r.,	Leu	Sei	Agn 240
Ile	Val	Ser	Met.	Lou 245	Asn	Asn	His	Val	Ser 250		Thr	Asn	Ser	Gly 255	Val
fier	Me∙t	Ala	.363 1.63		Len	H15	25-7	Met Obs		Lys	617	The	Ala 27g	Ριΰ	A1α
Ala	Ala	(11n 295	Ali	V i	41-11		Ala		#11n	Aran		V : 1 (*, t	-IIn	ΔΙι	Mert
	2° 1 2°0°		(1) \$		-(-	(n.			. + 2		11 y 3 (10)	. **1	. 11	Нÿ	Len
saise Nota		Υ	VII.		Ali ili			y		Ali	ř:	$\mathcal{L}(0)$	Vid.	4, у	Ser 324
Leu	Ser	Val:	Pro	41n 4.35	Al i	Tipo		ÄГэ		Asn	Чn	Alı	V :1	Thr 335	Pr.:
Al a	A ju		A K40			tuet.	The		100-1	Th:	Sirr		Alia Stat	-31r	Thr
ž4* 11	; • . <sub>9</sub>	1 7	1117:		len.	117	11.0	Istati.	i i	1.* -1		. t.	1	Tir	Asti
:	Cly	oly	dv	i :.	:119p				ji T.		500	7.1 :	po t	:·· :	11:
7 + 1 7 a	ă.,	u	V si	<u>;</u> 1	1 .	ř: i	•	-	*	A. I	* 2				
1.1	·	11	i i	÷ .	11	:	:								
		1	• -												

TG ,	TEGIGETET	GMAACGCCGA	G TENER CONT.	. :::',:'A.' AA:':::	GGAGGGGGG	AGCCTGGACT
GC 30	CGCAAGCCGC	CACCCCACGG	TOTTOCGATO	AGECCAAGAC	GOGTCAAČAD	GCTACAAAGG
AT 36	CCAACCACAT	GAGATCGCCG	GTCGCTGCCG	CCACGACGCC	CAGGCCATGG	GGCATACACC
ΓT 4.2	CGATCGCGTT	AACACGATCC	CTTCGGTATC	CCACCAACTT	GTCCTTACGG	CACCCAGGCC
°A 48	TOĞAGGTOTA	COCCTGGCAA	GVVCCACCCV	TCCGTATGTG	GATTATTICA	ĞACCGAGATG
	CHICLATCCT	( CTACCAEA	- AOMA POR	ACACGETTI:	ACCGCGGTTA	ccasgecgau
vG - v()	C TEL IGCAG	PAATGCCCT	OF THE PERCENTAGE	PIM NA NAS	A ACCORDO	TGATCCCG
iΛ 66	FACTGGGTGA	ACCC.CGGC	4 TEATTAG	"to introduce	HIT INVAIL	
G 72	/ Extractor	CAGGGACCT	TANDUT HING	AUCTOA : "A	CUGATOUAGU	BATGAGCGGC
Т "Н	A MTG IGOCT	MAR COCC	$\Sigma_{k,k,k}(M^k,\mathbb{C}M_{\mathbb{R}^k})$	$((0,\pm 0)NV(0))$	OGCACÇGG ; ;;	ownerse.
G 840	COMPRESSOR	· ATCAGGO	11 July 2 17 July 17	HIGAN YOM!!!!	A stocket if	om marke
C 401	aa waxaaa	NGTGGTTGA	7° 10° 12° 12° 11° 11° 11° 11° 11° 11° 11° 11	ASTIGOTACI	c7000016006	CRIBGSCOTG
i. (46).	es arractar	an Madeet	The state of the s	AMM : Top	W. Takina	To the state of th
1000	accasagagaga	GGAGGGATGG	ted assites:	endomini in	0. MAZ 13070	GOGATOGE II.
) ; (····	2.10.12V(1.14A5V, 1			alan karita	ar in Merit	F. BATTANA
* 1 14	· · · · · · · · · · · · · · · · · · ·	7:: 77: 37:	70 10 10 10 10 10 10 10 10 10 10 10 10 10	A. TOWN. A	dix i dix edi.	TWO WAYN
N 12 6	tha thaa	anna ivin	7. 11. 11. 14. 14. 14. 14. 14. 14. 14. 14	· · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	Mark by the
• • •	, which the tW. $^{\prime}$		AAAAA TAA	TIWARE.	A 11 1 N. 1 11.1	is the probability
	110 50.11		to a refuge		* * . · · · :	
		÷.				
	** *	Vicinity of the second		w. A		A 7
,	*		, a.e.,X			

### THE LITER OF THE PROPERTY.

### (xi) SEQUENCE DESCRIPTION: SEQ ID NO:108:

CTAGTGGATG	GGACCATGGC	CATITTCTGC	AGTCTCACTG	CCTTCTGTGT	TEACATTTTG	6,0
GCVCCC, ZEL	COAAA CAACO	CACT MOUTE	BAABAACGS 1	TO COTOOSA	TATOGTOGG	120
A RELICCATA	CONTROL	G CGHAAGAG	mportectiva	TCGGCCGCCA	CSACAMOCTO	180
TAWAY YA	N. WW. 1779	TAAACAJSA 1	AAAPELISA L	ACCUA TOUAA	, (TC)AA(Te;	.140
a mmaArrann	ementte ba	ATTOTAMON	AACT TROOT	Jaires Cal Valle	WACATOTON	£0.
**************************************	1777 11127 134	ADDAM T	A TOTAL	is meather.	vest meagy;	16
TICTICAGO	: ATTOTTC	TRIBOTA :	:::5)::::1	ATITITION	W.Moodaldet	42).
TAGUCCTCO	ħħ.					432

## AND THEOREMATION FOR BEING ID NOT TOOL

- REPORTED CHARASTERISTICS:
  - (A) LENGTH: 367 amand consequences (P) 1918: amino unio

  - STEEL STRANDED DEBUGE SINTER
  - L. Bart. Fr timesi

# x. m. maka sasakisi da sa, d d adea

The	i∽an Ala	Mari A	La Thi	F1.2	1:	711	1	120	1.1-1	11.	Ali	A1.1	ASD
		100				$1 \mathcal{O}^{ t_1}$					110		

- His lie Thr Gln Ala Val Leu Thr Ala Thr Asn Phe Phe Gly lie Asn 115 120 125
- Thr lie Pro the Ala Leu Thr Glu Met Asp Tyr Phe the Arg Met Trp. 130 140
- Ash Gln Ala Ala Leu Ala Mer Glu Val Tyr Gln Ala dfu Thr Ala Val 145 - 150 - 159 - 160
- Ann. The beautiful Wallys here our Pro-Met Ala Ser lie ben Asp Pro-165. 175
- Gry Alasker out for The The Arm Fro 11e Phe ody Met Fro Her Fro 187  $_{\odot}$
- Why Ser Jer Thr Fre Valody on Lod Fre Fre Alazha thr the tar 1995
- Lem Gly Gin Lem Gly Gin Met Jer Gly Fre Met Gin Gin Lem Thr Gin 210 - 255 - 256
- Fig. Levi A.n. on Mal Char New York Physics (Hin May Phys Thy That Gay 22)
  1.64
- Giv v Ad. in A. a Ary . He life Al con the congress of the life with the life and the life of the
- for the Control Annal Design Annal William Styles for
- Alamay Alamas and a same and what is a solution of the same against the same and th
  - The same the Capture of the second of the se

- (1) SEQUENCE CHARACTERISTICS:
  - (A) DENGIE: 100 amine ablors
  - (B) TYPE: amano assai
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SED IN NO:110:
- Mot Ala Glo Mot Lws Tor Acp Ara Ara Thr Leu Ala Cin Dro Ara Gly
- Ash the GHz Ara II) Ber GHy Arg ben by Thr GHz IIe Asp GHz Value  $\frac{1.7}{1.5}$
- call Sec Inc Arabilly Sec I.e. Since by Since tryeary are Arabilly  $\frac{37}{37}$
- The Ala Ala Can Ann Ala VII Val. deed the Gliu Gliu Ala Ash Lys SC  $_{\rm C}$
- win Lys Sin Stated Aspecia the best in Ash Tie Arg ain Ala Gly (6)
- Va. bin Tyo Ser Ari Ali Aspesia alto lin din Gin Ali ber ber ber
- alm Martinary The
- MI AMAÇIN ESELSE, NEW ELLE
- THE THE HAPA HERELT CON-A LETSCHELL OF THE PLANT OF TWENT HE STATE OF THE

TG	Nacaa	ET 1	MΨM	GMU	W G	W.Z.W.	E3A 1	` AN	MAY V	MEZA	±7507	ħ i A	377	.". ¥.,;	Muni		304
73(113)	I ITAT	CG I	AGCICT.	CK.	IC M	narij(	'TAAT'	. • .¥(	i-i-JAZ	SATG	$\mathbb{T}^{n}\mathcal{T}_{h}$	137111	TAY	ATT		11.	36:
CTTC	GACGA	GG G	GAAG	CAGI	d cc	TGAC	CVVC	СТС	CGCA								396
(2)	INFC	RMAT	NOI	FOR	SEQ	11) N	10:11	2:									
	(i)	(A (B	MENC ) LE ) TY ; ST - T()	NGTH PE: RANI	r 60 Aman EDNE	um: o ac DO:	no a Id Ling	aidr									
	.X.*	. This, '	· E.C.	r.	J.Sk.	! 7	N: U	o. i	1 7. 4	: 11.	:						
	110	ileri	F. y	A.B	Den. D	î.y.:	Thr	31 ta		A:*! 1 - *	1.1.	Vil	· † † 4		11.1	$F_{4.0,4}$	
	Gly	Seri	Leu	iin ≠	Gly	Gin	::P		Сту 15		Ala	Gly	Thr	A1-4 36	Δ!n	Gln	
	7.1 :	ħl.:	Val 35		As I *1	Elar	• Î î î	117. 411	**: :	A	A. 1)	Trye.	311.	ly.	Ha	Olu	
		Arj.	7.5	7:0	, 5, 1 ;		Arm e.e.		Ast 1		řv t		<b>V</b> i	Hr.	Tyr	A Property	
	14.4	25.1	farif -	٠	1 ° 1	(), i.	: † •	: .	2 1 4		,:. ·	,*. ÷	- 11	t.t.g+	1 /	: 100 t	·
	10 -	MAC	. · · · · · · · · · · · · · · · · · · ·	· ·	Ю,	. 1. 17.	:1:	* :									

WO 98/16645

### PCT/US97/18214

1.47

CCTGGTACGC CTGTGAAGCT CTACGGCTCC ACGGGCTCCC HAGGTT CTC AGGGACTGCT	,7 4 i L
TCCCCTCGTC AAGGAGGGAA TGAATGGACG TGACATTTCC CTGGATTRAG CTTGCCGCGG	300
CCTCGATACC CGCGAAATTC CACTGCTGCT CTGTCATGTT TTTGCTCCGT TTCTTTTCGT	360
ATTAGCGGGT CAGAAGCCCA TTTGCGA	387
(C) INFORMATION FOR SEQ ID NO: 114	
(H) SEQUENCE THARACTERISTICS:  (A) MENOTH: 072 base parts  (B) TYPE: 1.0010 or activ  (C) STEAMSEDMENT: single  (D) TOPOL, CY: linear	
(xi) SEQUENCE DESCRIPTION: SEA 19 N 0:114:	
SOADDOORS TOTTO TOTAL SECUENCE SECONDARY STOEDSTATE SCADONOOD	60
TROSEST OF ATHETERIST TESTICIAL CONTROL TO ANTICATOR ACTIVITATION	170
TTO/COOMSCT TTESTTOCOT FROTEIN SIA CARSSOCIAMI COOCCOCCES F. ACGISEGIAC	140
COMMENTACE OF FOR ADMINISTRATION OF PROFESSION OF ADMINISTRATION OF ADMINISTRATION OF A STATE OF THE PROFESSION OF A STATE OF A STAT	. 40
er militat provincia (Militate M	.: /.
THE PRATTIENT FOR THE BEST OF THE FOREST STATES	
THE THE MERIT HERE ARE A THE SHOPE IN THE SECOND STATE OF THE SECOND SEC	

and the second of the second o

(a) SEQUENCE CHARACTERISTICS:

A. LENGTH: 15 amin. Silate

1.48

```
(P) TYPE: amino arid
          (C) STRANDEDNESS:
          (D) TOPOLOGY: linear
    (xi) SEQUENCE CONCRIPTION: OF, IN POSITION
     Ala Vai Glu Sor Gly Mot Low Ara Leu Gly The Pro Ala Pro Ser
T. * INFORMATION FOR JEG IN NOTE.
     il sequence chara tericilo:
         (A) hELVIII leamin and.
(B) TYPE: amino and
         (G) OTPANPERNESO:
         (1) TOFORWAY: Linear
    SE SECTION CONTROLL NOW, IN MICHAEL
    Also Also Mercury (i.e., Area Inc. ) y Aria (y y Fr. ) perces i Also Also Inverse.
    H. Hy and
   THE FMATER FOR THE FOREST
```

```
GA DEMOTH: 14 main. activity
```

- (P. TYII: imit.o actd
- (C) STRANDEDNESS:
- (f) TOPOLOGY: Linear

### (x1) SEQUENCE LEGGRIPTION: TEQ ID NO. 110:

App lie Guy Jed Gla Der Ihr Gla Asp Ölh Gin Xaa Aid Vai

# (3) INBORMATION FOR SECTIONARY :

- OF SEQUENCE CHARACTERISTICS:
  - (A. LENGTH: 1) aming aring
  - .BT TYPY: amin. actid
    (C) STRANDEDURSG:

  - 415 Policiosa: limbar

# (x) SECTION OF THE SECTION OF SECTION OF THE

Ala Cu Alaska Leeska ing Kaasan Kasa ito Walija.

# CONTEMATE NOWER OF THE NEEDS TO

- INTERNATION OF SEASONS FOR THE POST
  - A LENGTH: 10 miles and a
  - De MARKE IN THE LEVY I

  - THE SECOND REPORTS OF

And the second second second

WO 98/16645

150

PCT/US97/18214

(B) TYPE: amono acld (C) STRANCEDNESS: (D) TOPOLOGY: linear (xi) SEQUENCE DESCRIPTION: SEQ ID NO:122: Ala Pro Lys Thr Tyr Kai Gla Dia Lee Lys Gly Thr Asp Thr Gly 10 12) INFORMATION FOR DEC 12 120:123: TEC DEQUENCE SHARAGTERISTION: (A) LENGTH: (9) amains a wide (B) TYPE: amino acid TOT STRANDERNESS: (D) TOPOLOGY: Ilmear CXII SEQUENCE DESCRIPTION: CRO IN Terminal App Pro Ala Ser Ala Pro Asp Val Er - The Ala Ala dia let The Ser Des Des Arn Jerrica Alla Alla Producti Valider Inc Alla Arno on INDEMAND A FEB. 15 quarty qu THE THE TO MEAN TO BE TO THE Proceedings of the second of t

```
(.) DEQUENCE CHARACTERISTICS:
     (A) LENGTH: / amino acids
     (B) TYPE: arano acid
     (C) STRANDEDNESS:
     (b) TOPOLOGY: linear
(x1) SEQUENCE DESCRIPTION: SEQ 10 No.125:
Amp Pro Bly Cyr Thr Presidly
1
THEORMATION FOR SELLID BOOKEN.
EFF DEQUENDE MARATERIOTION:
     (A) DENGTH: I amin for m
     (B) TYPE: amano acid
     (C) STRANDEDNESS:
     (I) TOPOLOGY: linear
(ix FEATURE:
 ty OTHER INFORMATION: Shows "The despirite Like Name Her Rither a
TWEET OR, MEDITED THAT IS THE LAR. IT DO HAVE I
Real Mark Gir Place That of your control for the
1
FROM THE WAY OF STREET
     e et gal.
Si Singing te elle Mer elle elle elle elle elle
```

```
BUT INFORMATION FOR CHO IN N :1198:
```

- (3) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

### (x:) SEQUENTE DESCRIPTION: SECTION::128:

Xãa Xaa Xaa Slu Lys Pro Phe Leu Arg

- IT INFORMATION FOR SEC II No:12.3:
  - (1) PEQUENCE CHAPACTERICILISE
    - (A) LEWGTH: 15 amorth across
    - (B) TYPE: amino avid f() DURANDEPNESS:

    - (D, TOPOLOGY: Linear
  - (X1) SEPTENCE LEDTERATIONS OF, INC. NO. 18
  - Man App Syr And Lyu Ser Ara the size time Val Through As a ser I  $_{\rm c}$
- . THE EMAIN OF A BOOK, IN Mark to
  - THE SEQUENCE THAPA TORRESTED OF A 1831 Mills III american and a fifted and live a
  - er eggstatet and the commencer

```
(B) TYPE: amino acid
```

- (C) STRANCEDNESS:
- (D) TOPOLOGY: Linear

### (x1) SEQUENCE DESCRIPTION: SEC 10 NO:131:

Ala Fro Glu Ser Gly Ala Gly Leu Gly Gly Thr Val Gl<br/>n Ala Gly 1 $^{\circ}$ 

### (2) IMPORMATION FOR SEQ ID NO:130:

- (1° SEÇUDNCE CHARACTERISING):
  - \*A) LENGTH: 21 amino acids
  - (P) TYPE: amino acid
  - (J) STRANDEDNESS:
  - -D) TOPOLOGY: limear

# (RIB DEQUENCE DESCRIPTION: SE, in to :13, :

Mag Tyr the Ata Tyr Xia The thr Ala Gly lie Valler. Gly lye lie A

Ash Val His Lew Val

### LI MEDEMATOR FOUND IN 11 KG

The state of the s

THE CONTRACTOR OF A STATE

GBCCSSGTAC	CTCA-TCCGG	673637366A	- 17(1133)7177(	GCCTOCCCC	WWG IGCCGG	30(
CCGTGCCCGG	TGTTGTGCCT	GCCCCGGTGC	CAATCCCGGT	CCCGATCATC	ATTGCCCCGT	360
TCCCGGGTTG	GCAGCCTGGA	ATGCCGACCA	TCCCCACCGC	ACCGCCGACG	ACGCCGGTGA	420
CCACGTCGGC	GACGACGCCG	CCGACCACCC	CGCCGACCAC	GCCGGTGACC	ACGCCGCCAA	480
COCCOCCADO	SACCACGCCG	GIGNOCNÈGO	CGCCAACGAC	GCCGCCGA:\?	ANGCOGGTGA	540
CCACACCAAC	AACGACCÓTC	OUCCCGACGA	acercañace	GAMGACGATE	POTOCGACCA	600
CONTROLLER	GACCACGITT	GCTCCAGCCA	Colora Medicina	CACGACCCTT	GUTCCGCAGC	660
Pargara		CAADIAW	interiografia	AACCTAG 'A ‡	waa ngree	726
HONICGEAGAC	GOTGGCGCCO	DOTTOGOAGO	(13C) (1T(3C))	TGUCCGCAAC	nur Andågen	790
Unionackann	ATT 3000000	CTCTUATOA	annacient	TCACTA.W.IT	`` JAHGACAT	840
GGCCGGTGAT	GCGGTGACIP	TGGTGGTGG	CTOTITOAAC	GL		882

- 4. \* INFORMATION FOR SEQ IS NO:134:
  - (.) SPQUENCE CHARACTERISTICS:
    - (A) LENGTH: 815 battle pairs

    - TYPE: macloss acid
    - Co Colocosy: Linear
  - ... MEDEMONE TYPES DNA perviole

### HOUSE THE THE THE THE HEAT OF THE HEAT

grander er egger i by element element og kring gren er er g

WO 98/16645 PCT/US97/18214

155

ACCEGETATE	TOTOMOTE	CCTCMACCTY)	CATCATGATS	TOUAGGTJAC	IGTHACCGCG	600
CCCCCCAAG	GAGGCGCTGA	ACTOGGCGTT	GAGCCGATCG	GCGATCGGTT	GGGGCAGTGC	660
CCAGGCCAAT	ACGGGGATAC	CGGGTGTCNA	AGCCGCCGCG	AGCGCAGCTT	CGGTTGCGCG	720
ACNGTGGTCG	GGGTGGCCTG	TTACGCCGTT	GTCNTCGAAC	ACGAGTAGCA	GGTCTGCTCC	780
GGCGAGGGCA	TCCACCACGC	STIGOSTONG	CTCGT			815
(2) INFORMA	TION FOR SE	O ID NOTE H	):			

(i) SECTENCE CHARACTERISTICS:

(A) DENGTH: 1152 base pairs

(E) TYPEY route; said

(C) DTRANDEDNESS: sinfle

(E) TOPOLOGY: Linear

Till MOLECULE TYPE: DNA , delomic

ACCORDENCE DESCRIPTION: USED IN INVIEW:

ACCORDENCE SECTEMBER: TOAGATORIA WATERTONER ACTORDORGE SCRITTCAGE - 60

CTTCTCCCAG AACAACTG TO WAACATORIC GOOGGEGAAA WAGGGGGTWA TTTWA GGTE - 120

TATWADAAGA TERTE WAS DESCRIPTED WAATERATA TARKAANA TEREGORGA - 140

CTTCTCCCAG AACAACTG TO WAACAACTR WAATERATA TARKAANA TEREGORGA - 120

CTTCTCCCAG AACAACTG TO WAACAACTR WAATERATA TARKAANA TEREGORGA - 120

CTTCTCCCAG AACAACTG TO WAACAACTR WAATERATA TARKAANA TEREGORGA - 120

CTTCTCCCAG AACAACTG WAATERATA TARKAANA TEREGORGA - 120

CTTCTCCCAG AACAACTG WAATERATA TARKAANA TARK

WO 98/16645 PCT/US97/18214

156

GITGANACIIG	CTGGC GEGT	t il valtulit.	MATSTAAT	CCCPTGCSES	CAGOCCGGCC	960
CGČ TG 3CCGA	ATACCAGCAG	ATCGGAÇAGC	GAATTGCCGC	CCAGCCGGTT	GGAGCCGTGC	1020
ATA COGOGG	CACACTCACC	GGCAGCGAAC	AGGCCTGGCA	CCGTGGCGGC	GCCGGTGTCC	1080
GCGTCTACTT	CGACACCGCC	CATCACGTAG	TGACACGTCG	GCCCGACTTC	CATTGCCTGC	1140
GTTCGGCACG	AG					1152
(?) HIFORMA	ATION FOR SE	Q 10 NO:136	·:			

- :, SEQUENCE DEARACTERISTICS:
  - (A) LENGTH: 655 base pairs
  - (B) TYPE: numbers and d(C) STRANCEDMERO: cinqle
  - (D) TOPOLOGY: Finear
- (11) MOLECULE TYPE: DNA ( periond "

# CARCARTERS SCHART RESCRIPTIONS SET THAT OF TOCATIONS TOCATIONS TO CONCATON ASSESSMENT OF THAT IS NOT A CONCATON TO CONCATON ASSESSMENT OF THAT IS NOT A CONCATO

(ii' MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEO ID NO:137:

( , , ,	.,,,,,,	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	. 02	JOINE.			., .			•					
Asn 1	Ala	Val	Val	Ala L	Phe	Ala	Val	lle	Gly 10	Phē	Ala	Ser	Leu	Al a 15	Val
Alā	Val	Ala	7(: 7(:	The	11.	Arq	Fr ()	Chir 25	Δìa	Ala	Ser	Бүз	P1++	Val	Glu
GIY	His	Bln m,	Asa.	Ala	G.1.		Cly	Lyr	Ph	Met	Pro	1.6.11	letti	Pro	Thr
-31 n	G1:.	TI n	Al÷	P* .	V.::	; · · · ·		Pro	:'1 ·	; I * -	Asp on	Arŋ	President	Thi	Ala
@1y 65	1411	Oln	GLV	ary.	Thr 70	110	Pint.	ΑĒτ	val.	41n 75	Asn	kal.	Wei	Ero	
Pro	÷1γ	Thr	Ser	Em.	Gly	$\nabla a$ :	1) y		T!11 40	Pro	Αlα	ma	Pro	ДЈ а 95	Pro
1. g	Ala	f <sup>4</sup> r	Ala Lor	Tal	11:	GΙγ	7.1	Tai Tor,	: 1	Ala	iro	Val	iro	He	Pro
7.0	(**	115	ļ	11.00	I : ·	i*: -	En-	1 1	81 <u>y</u>	7:1:	0.i.	Prin	÷Гу	Mint.	Pr.×
*** ·	1 l	ir	tin	<i>I</i> .1 →	! :	1 : 1 :	i:.•	1.1.2	: 1	Tyl	Thr	Har	J.O.Y	Al i	Titt.
1 d	t · »	· ·	i in:	11.1	1)	: :	Lit	1.1	î ;	7::	3-7	11.,	11	Ere	Th::
	٠.			* .				ņ ·						: r : ***	11.1
	** 1	*			1 -	*.)	; ·			,		::···		75.1 d	::

The second of th

# (2) INFORMATION FOR SEQ ID NO:138:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 174 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (P) TOPOLOGY: linear
- (11) MOLECULE TYPE: poptide
- (x:) DEJUENCE CERCRIPTION: DEP IN NO.136:
- The Ash Gib Pro Lou Ata Pro Lyp Ala Pro Pro Asp pro Fro Bor Pro I
- Pro Arg Fro Fro Val Pro Pro Val Fro Ero Deu Pro Fro Ser Pro Pro 20 25 30
- See Pro Pro The Gly Vep Valler. As ratio Lea Lea Fro tro Trp Lea  $3^{12}$  .
- Ala Sly The free ProcAla ProcEro Val Ero Pro Met Ala ProcEro Pro 50
- Pro Ala Alaur Leon in the Leon bracks. Leon inc. for the Leon from the uu
- Short Him I'm The Arm Free Energy And Free Energy Alastes (Free Aras) and the Section (Section 2017)
- All the are the called the second of the area at
- er er die de la Santon en de l Santon en de la Santon en de l
  - , and the second of the secon
- Here Allen in the Conference of the Allen and the Allen an

- (A) LENGTH: 35 dmine acids
- (B) TYPF: amake acid
- (Č) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (xi) SEQUENCE DESCRIPTION: SEQ 19 No: 139:
- Oln Pro Pro Ala Glu Val Der Amp Oln Arg Val Ger Gly Leu Thr Gly I Bo 15
- Also Val. 30 n for der fire Arg Thi Thir Also Glin Asp for Air Fro Arg 20

Ash Aru Ary 35

- (2) INFORMATION FOR SEQ ID NO:140:
  - \*: DEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 104 mane nords
    - (B) TYPE: amino acid
    - .C) STRANDEDNESS: Simple
    - 'D) TOPOLOGY: linear
  - Ar stroughter representation
  - omic objects in the first series of a constant
  - And Alle Ser All the Ser and the And the Ser and the Ser

  - Provide Anna Bara, the principle of the control of th

100

- (2) INFORMATION FOR SEQ ID NO:141:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 53 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: other nucleic acid
    - (A) DESCRIPTION: /desc "PCk primer"
  - (vi) ORIGINAL SOURCE:
    - (A) ORGANISM: Mycobactorium tuperculosis
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:141:

USATURIATAT GGGCCATCAT MATCATCATO ANGTHATONA MATCATOGRA ACC

- (A: INFORMATION FOR SEQ ID NO:142:
  - FOR SEQUENCE CHARACTERISTICS:
    - (A) DENGIH: 42 base pair.
    - (B) TYPE: nucleic acid
    - (C% STRANDEDNESS: Single
    - dry TOPOLOGY: Linear
  - (14) MOLECULE TYPE: other matters and
    - (A) PERCEPTIONS Form "PT Firmer"
  - TO REGINAL DOUBCE:
    - MAI ORGANICM: Myoula terrino tur orbal acro-

NAME OF THE PARTY OF THE PARTY.

- - MERMAN BOOK OF THE EAST
  - TOTAL A CHARLET PRESCRIPTION

    - Z. Tar Modelli. In the second of the CVIII of the second o

3.1

(xi) SEQUENCE DESCRIPTION: DEQ ID NO:143: GGATCCTGCA GGCTCGAAAC CACCGAGCGG T (2) INFORMATION FOR SEQ ID NO:144: HIT SEQUENCE CHARACTERISTICS: (A) DENGTH: "I have pairs (b) TYPE: numbels wid ofence subgrouped contin ala ToleHCarr linear The Mantewood Table of Leading Court of the Auto-٠. v. Mildinaly J. URCE: A GROWNING Mycons atoming turner and using ax. Sequence procedures: TE: 10 10:144: CONTRACTOR ASSESSMENT AND THE SERVICE OF CONTRACTOR FOR SEASON TO 1401 THE THE TENENT OF A PART OF A LOSS O A LIMBERT CONTRACT E IVERT BUTTO L'ORIGINALE L'ARISE I I HOUSE FARMER TOTAL TERMINATE AND THE TRANSPORT OF THE PROPERTY. e e s é, sign.

(VI) ORIGINAL SOURCE:  (A) ORGANISM: Mycobacterium tuberculosis  (XI) SEQUENCE DESCRIPTION: SEQ ID NO:146:  SAVINGAATIC TOAGAAGCCC ATTEGRAGS A A  3.1 INFORMATION FOR SEQ ID NO:146:  (A) SEQUENCE HARACTMEISTIMA:  (A) SEQUENCE HARACTMEISTIMA:  (A) SEQUENCE HARACTMEISTIMA:  (A) SEQUENCE HARACTMEISTIMA:  (B) STERNBEIMANCES: STERNE  (B) SOURCE: STERNE  (B) SOURCE: MYCOBACTER TORS DESCRIPTION:  (A) ORGANISM: MYCOBACTER T	( : · · )	MOLECULE TYPE: Street musicus acid (A) LECURIPTISM: /desc  "FCR primer"	
CONTINUENTATION FOR SECULD NOTIFICATION  (CONTINUENTE TO COME SECULD NOTIFICATION OF THE SECULD STREET TO COME SECULD STREET SECULD STREET SECULD STREET SECULD STREET SECULD STREET SECULD SECULD STREET SECULD STREET SECULD STREET SECULD SECU	(v:)		
CONTINUENTATION FOR SEC. 10 NOTE 14:  13 SECURITE 190 Come pairs  (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs (P) Type macled and the pairs of the pairs (P) Type macled and the pairs of the pairs of the pairs (P) Type macled and the pairs of the p	(x1)	DEQUENCE DESCRIPTION: SEQ ID NO:146:	
Degree of the parameter of the part of the	-3AGAGAATI	FC TCMGAAGCCC ATTTGGGAGG A A	3
(A) DENGTH: 1900 tomor pair: (B) TYPE inder[vi   redd (C) STRANDEDNESS: Junite (J) TOFOLOGY: lines)  (J) MERECOLE TYPE: DNA feeren; (A) ORGANISM: Mycoba derion tomor mileste (X) ORGANISM: Mycoba derion tomor mileste (X) HEATURE: (A) ORGANISM: Mycoba derion tomor mileste (X) HEATURE: (A) HAME / KEY: Clos (B) LO TATION: TYPE: Clos (B) LO TATION: TYPE: Clos (A) JUNITED TO A GARAGE CONTROL OF THE	\$.11 INFO	RMATION FOR BEC 10 MD: 14/:	
(A) ORGANISM: Mycobasterior tober milesion  IN. FEATURE:  (A) NAME/KEY: Club (B) LO WITCH: I'  (X) JELDENCE BEDURINGE DE CLUB DE STELL  (X) JELDENCE BEDURINGE DE CRUB DE CLUB DE STELL  AND	- т J	(M) BUNGTH: 1990 come pari: (R) TYPE: nucleic acid (r) STRANDEDNESS: cinibe	
(A) ORGANISM: Mygrobaldering tober milent:  (A) NAME/KEY: Childer (A) NAME (A	. 1 1 )	M SECULE TYPE: DNA ( enren) )	
AND MAKE / REPORTED TO A LITTLE OF THE PARTY	$_{i}\vee _{x^{(i)}}$		
TRITTETTE AND THE MET A SEA CAN THE STATE OF THE MARKET PROPERTY AND THE MET A	tx.	(A) NAME/KEY: (CLU	
ANNAL COMMANDE COMMAN	) X	JENTHENZE BERTHEITER TER D. D. D. HOLLEGE	
	Coloration is	A COMPAND FOR FAIRANCE CONTROL A SERVICE OF THE HERA	. *
	72 - 74 . T	a saas martin in land as time of an old of a continuous and or mineral or	1.
	1. Y. 1/4/4/	AARAN MAARAMAA MATEE TERRAHAMA AMA AMA AMA ATEE SERJER METERSERIES SERJER	
			• (1)

				Tyr					ili					Thi	in jij Gly	41.
			Gly					Ala					Asn		GGG G1 y	460
			GCC Ala				13] 11									५ () भ
			ET:													556
			OTH Val		$\cdot : : \cdot_1$			V .								σένή
			(Vi.) (In									$\operatorname{Fr}_{2,n}$				652
			11 *			Ä		: :								yniv
H S pt	761 + 741 (‡ 1 + 1	p (c)	ow Aug		: · · · · · · · · · · · · · · · · · · ·	(1-1) -41-; -1-;	ida Istoi	7 1::	i '	 i •• :	Edin Edin 174*	personal de la companya de la compan	West des	Tāsi Tyri	mist Dem	१४ कु
			- Will Festi	1.1					×							
			i ş								e †			· ·	a.i.	÷::
		. * .														
i.															-y	-0

	181.										. 1					2 3 3	
				ger Ser		-11e					${\rm Fro}$						1684
				GAG GLu 315													1132
				And The													1130
	A. T.			71 · · · · · · · · · · · · · · · · · · ·													12 8
4				ndel Lyr													
	ľAC.	•••• /4,	TT (	MANT.	'EW'	1 000	74° 1740	i''AA		347.3	(1137)	`= 'A.C	, G131)	C.P. 64		acuca	1373
	FAT	arre.	;/ ;/ · ·	X		T GA			ī. ·	•:.•		v, :::	AI. T	ii.i T	HIT	JGGTL	1,693
	r iAT	A 197	n)	i itt it.		: - "	1 1111		7::		: :	RTTG	7	ro n	THAT	CATC:	145.5
j		CAT	277 7	n ko W	7 7.	• 0 • 1 • 1	· WW	<b>ا</b> : ::	7:17	STIT		700	2013	(* ):	7.7.7.6	W.TI.	1111
	77(3)	'AA'	7v * .	TA 13	. 7	i. i.	. el.l	4T 74	٠.:	. · · · ·	17	; • • •	AT W	·	· ; ;, ·	1.5.77.8	11.00
	"I W.	11 (11)		T. L.	: '; 1	/ T		:: ;; <b>:</b>	• • •	÷	ā, 1 1		PART		• • • • • • • • • • • • • • • • • • • •	777 13 v	1.00
		: · ·	٠	:	474		; **:			7.3	*2 · /· ·			. • •			÷ ,
	. 1-55	er tr	*	100	i A. i. ji.	7:1:	17 - 4	Milit			1-20-2		50, 1	4.			
				·													
									. •								

	Mar	 2/11		40.00				
	18° 4	1.	**		***	1 1 7 .	 	
		 2. 4. 4	, .					

### (x.) BEQUEEUE DESCRIPTION: UEQ ID NO:146:

Val	Lys	11e	Arg	Leu	His	Thr	Leu	Leu	Ala	$V_{\rm cl}$ l	Leu	Thr	Ala	Ala	Pro
1				5					10					15	

- Leu Leu Leu Ala Ala Gly Cys Gly Ser Lys Pro Pro Ser Gly Ser  $\frac{20}{100}$
- Pro G.u Tar Gly Ale Sly A.a Gly The Ye. Ala The The Inc. Ala Sec. +6  $$4.5\,$
- For an Wall the left Ala over the Try for the horizon for the horizon  $\alpha_{\rm c}$
- law Alb 180 Tay New York After Head Row And Typ Proc After Val Times
- Also dly Thr Ma. Arm lie dly Alasker Ada Alaskyr few Ger witz gly 100 .  $$10^\circ$$
- As a Din Din Ma. And Typ Admiller are styled for stir Baselin by: The District District Medical District Distri
- Dealest Anglight Mat For Ata Alas Marin de Milliold The Called Inc. 14
- The Alicense Park are the Arte Double Control for the Maximum District.
- est that a control of the control of

- \*

Pho	Len	100 275	Fre	Ás. I	ř <i>i</i>	i, n	260 260		(1 r:	Al i	A) a	A. a 285		÷γ	l'ne
Ala	Ser 290	Lyt	Thi	Pro	Ala	Asn 295	GIn	Ala	He	Ser	Met 300	He	Asp	Gly	Pro
Ala 305	Pro	Asp	Gly	Tyr	Pro 310	11€	He	Asn	Tyr	Glu 315	Tyr	Ala	He	Va]	Asn 320
Asn	Arq	Gln	Lys	Asp Ras	Ala	Ala	T).r	Ala	GIn .335	Thr	Len	Gin	Ala	Phe 3 yb	Len
hi.	1:1:	Ala	110 346	/m	/sitp	olγ.	Arn	170	Ain	der	File	Less	Adp.	Hin	Val
	Phier	ili iti	1, 1, 2	11	i'r a	Þο	Ana spir	VAL	V + .	Гур	Len	Ser Bek	Asp	Ala	1.6112
1	A1.4 +7g	Phr	: 1+	1 + 1	. 933										
(1° r	ine	TAMA	TON	F13	BEÇ	IP t	19:14	· 4:							
	(1)	. :		MGT: THE: BAMI	i: 15 macl EDNE	TERI 93 i efe ec: Line	450 10 - 1	joili i	÷						
		٠.	÷	* ::	. 11	11	i.: ."		• ;	: 1 : •	:				
	11:					' v 1951 v	والمناوع				A 78.1		9 3		

$\mathcal{O}(\mathcal{O}(\mathcal{O}(\mathcal{A}(X)))$	: TACONALI IN	CHAT MAAA	11 11 12 17 17 17 17 1		AASTORS II	66
COCCAGOSTS	Macaramas	PIACIDEG (T	AUTOMA	CACCCCICAL	ANGGGTOOGG	7.2
TGACACCTTC	TTGTTCACCC	AGTACCTGTC	CAAGCAAGAT	CCCGAGGGCT	GGGGCAAGTC	78
GCCCGCCTTC	GGCACCACCG	TCGACTTCCC	GGCGGTGCCG	GGTGCGCTGG	GTGAGAACGG	846
CMACGGCGGC	ATGGTGACCG	GTTGCGCCGA	GACACCGGGC	TGCGTGGCCT	ATATOGGCAT	90(
WW.TTCCT:	"PACEAGO" A	-CONACTOR	ACTOCK WY	CONCAACTAB	GCAATAGCTC	96(
TRICAMITO	TTTT PER TE	h(x) = h(x)	MIRWILL	**** **** *** * * * * * * * * * * * *	TTCGCATC	1020
MAAROON I	1;C;;;;;;(C;;;;;;;;;;;;;;;;;;;;;;;;;;;;	SATTTOSAT	MTD,MC.TM	ornang si i	AGGIGGTACCC	1077
MI-MICANI	TACGASTANG	AKTTGGTTAA	AA 2000'AA	AAGGACGÇOĞ	- Anogogoa	1140
$X = X \times X$	57x177174	ACT GERVEAL		As a PANCE FOR	CUTTOTTCGA	1200
HAW HALLING	end(2523	Til ver tamb.	HT HE WAL	TTOUTCAS	TO TOATCOC	12n0
GANGATT 10 1	AGCTAPTT	SELLINY (MU)	ACTINIA, ACT	AVYTOCHT	HERCCATCGG	1326
1.5 . 1 . 1 . 1 . 1 . 1 . 1 . 1	GGAGLAT ET	Thursday.	::::::::::::::::::::::::::::::::::::::	tt ware and Little	C. GGCATT	1350
	71 1.3.3M77.1	N 10211177	n. Togarita	CONTRACTOR	GETECTECTC	1449
	T NAME OF AN	93.11 () 13.11 ()	A PERPARAN	HTT JONETT	CTTCAGGGGC	1500
ASSIMATERAL		·,, 21, **,,,,	7.0% 1.31 I	hopping Works	Harry Mir.	4.5
**********	mij militaria.	paggara.	a sweet tan te	9, 71 THE	2.1 The Williams	1.0
4		77.71	A MARKET	· · · · · · · · · · · · · · · · · · ·	mir day till	÷
To be with the	1 7 1 1 1 m	and the Figs	"Zvi-2" - Enit Zvi-	// · · · · · · · · · · ·	i Jam Sty.	-1
* : *: *		111			e	. 6
· · · · ·		4.				
	$\mathcal{I}_{X_{i},Y_{i},Y_{i}}$					
7.8	· · : s			*	in the second	*
	• 8					

. ...

xi) SEQUENCE DESCRIPTION:	: SEO ID	NO:150:
---------------------------	----------	---------

Mert 1	Lys	116	Arg	Leu 5	Hip	Thr	Leu	Leni	A1a 10	Val	Leu	Thr	Ala	A1a 15	l'r
Lesti	Deta	Levi	Ala po	Ala	řil i	Тij	C7.3	: '' Y		i yn	Pri	) In e	30 r 30	Пу	Sec
1111	::I:.	Tran 30	.177	Ala	ij, y	$I_{i}1_{ij}$	о . у 4	Thr	Viril 1	Аlч	The	71.1 41	3,10	Аlн	See
	Fr .	val.	ma <sub>1</sub>	Zer:	/s1 a	*: 1 \. :	The		(h-)	Thr	11	1 1	Tyn	Pre	Let
Pare 60	$A^{*}t$ .	Lea	111	ŧγ	1 F 2 G	is *	1 200	i	î.,	ĀI I	771	Ĭ:	ASH	Vali	1 h t 80
1100	r.,	ÂΈ	Glr.	Gly As	The		.i++t	117	Alia An	r	1	Ž.	w.	āla ge	Ala
$\tilde{c}_{X}$ . 1	Gly	Thr	W 1 1-7	A. Tr	1.7	· 7	* - 1	.`**1 ** <u>1</u>	A. p	Alsı	171	·	. in	47. h.	Ну
A.p	la	Alai Lista		Hiri	Lyr		1. (		Art.	11.	$K_{ij}^{\dagger}$	1.0-1	Al.	He	ß:
ā ,	*: · · · · · · · · · · · · · · · · · · ·	. \$	V si	Asst.		A :	0	î:	,		160	* - 1	īi .	In a	,.
# 1       *	Ä.∵.		* * * * * * * * * * * * * * * * * * *		1 + 1	ň.	2.1 t	11	411	+11.	i i y	er,		ē	1: -
:::	şs it	:	} •	:	. •							•			
: •										;				* \$	٠

- Ser Sin Arg. My Sen Cly sin Aba Gan Lee, aly Ann Ser Ser Gly Ann 263 260 Phe Leu Leu Pro Asp Aia Gin Ser Ile Gln Ala Ala Ala Ala Gly Phe 275 280 Ala Ser Lys Thr Pro Ala Asn Gln Ala Ile Ser Met Ile Asp Cly Pro 290 295 300 Ala Pro Asp Gly Tyr Fre the Lie Asm Tyr Glu Tyr Ala He Vil Asm 315 511 Ash Arg Gin Lys Asp Ala Ala Int Ala Sis Thr ber oin Ala Phe Len 330 His Phe Gin Ito Let Pro Iro Air Vi. Vi. Eys Lou For Asy Alic Lou-355 lise Ala Thr like Der Der 3777)
- . THE THE PRACTICE FOR SEQ ID N : 151:

. 5

- Fit DEGUENCE CHARACTERISTICS:
  - (A. LENGTH: 1777 base parts
  - (F) TYPE: modely and to surprise property and
  - THE FAMILY CONTRACT

orași (La Para Cara Brazil Na La Princia de La

TOATETTACT	$T^{\prime\prime},T^{\prime\prime},\Gamma^{\prime\prime}M^{\prime\prime}M^{\prime\prime}$	CAPCATOLLS	76-11-170 JAAN	talantaalkani	i Gerage, Leec	*: ()
TGCGATCTGA	TCGGGATGS	COCGTCGGAC	AAGCTCAGCC	CATCGGGACC	CGACCGCTAT	660
AGCTATGGCG	AGCAACGAGA	CTTTTTGTTC	GCGCTCTGGG	ATGCGCTCGA	CCTCGGCGAC	720
CACGTGGTAC	TGGTGCTGCA	CGACTGGGGC	ressected	GOTTOGACTO	GGCTAACCAG	780
CATUGCGACC	GAGTGCACCG	GATKIGGGTTG	ATSCAASCCA	TOGTCACCO	GATGACGTGG	241
O GGACTGA	in Chira	1 % 600-11.1	LI ZAMBGUT	TOOGATOGO	TEAAGGCGAC	900
a Www.i., i.	TO WOOM TO	· · · · · · · · · · · · · · · · · · ·	4 (17.25) 1 (1.37) 1 (1.37)	tesandere.	GATCCTGCGA	4(3)
$m_1 \dots n_m$	$\lambda_{i}^{i}$ An man."	- 1717 * * * * * * * * * * * * * * * * *	"NEED TO AT	TUCITOAACGI	CFGCGAGGAC	1920
THORONA	· 33 7277 377 ;		11 7 1 3 3 1 CM	Variatios Williams	SHOURAGOTO	: Ch.,
41.46 3.77 43	$\mathbb{T}_{\tau}(YX) \cap \mathcal{Y}_{\tau} \cap \mathcal{Y}_{\tau}$	13210000	To see Here's	COACATEC	-WAACTGTT 1	1141
ATOMACIONI	A: "1" 10"1"	MTTATEA T	BUILDING.	THACTATET	CASGAGCTGG	1200
GCDWC AJA	FIC WAST WIT	Assistanti		TICAGGAG A	TAGCGATGGC	1 460
orogran der		d validas an	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	* 18** * 10** * 10**	CATTOMACGA	13. /
W. MARAAT	** D: #WY ID. 1800	$(\pi_i \mathcal{A} W : h_i) : \dots$	in the transportation	GTCAACTCAT	AAGWYTTCCT	1380
· · · · · · · · · · · · · · · · · · ·	W.M. T. W	e e waya aza e	A MATE	A ***C0.1;1 **	TTOKANGA	1440
TT AND.	1	A CHAMAGE	1		/	1 10
	e production of	Tagar Valla	#***WEW	· · · · · · · · · · · · · · · · · · ·	for 1 m 1971 117.	:
	W. W.A.	is www.iiii	i AA Veet	The same T	·*·····*10····	1.4
					A 1 1 1 1	

CREAT SEQUENTE SECUREFICAL SEQUENTES IN A SECURE	
GACATIGAAL CSTACCGUIC TUCUTAGOG CTCC MOONE TGAATGCO'A TATCACGCAC	60
GOCCATGTTC TGGCTGTCGA CCTTCGCCCC ATGCCCGGAC GTTGGTAAAC CCAGGGTTTG	120
ATCAGTAATT CCGGGGGACU GTTGCGGGAA GUCGBCCAGG ATGTGCGTGA GCCGGGGGCG	180
OSCOSTOGCO CAGGCCACOS CTGGATGCTC AGLCCCGGTG CGGGGACGTA GCCAGCGTTT	240
RETIRE PETER STONAMANTE STACTORS IN GAINAGES REPOSESTORMENT GRENAAGAC	10.1
COMMAND OF CONTROLLING AND	35.4
(20 INFORMATION FOR SEC ID NO.152:	

HOUSENCE TEARACTERISTICS:
 (A) LENGTH: 1338 house pairs
 (b) AVHIS masters acris
 (c) STRAUDEDMESS: Singles
 (d) TSIOLOGY: Timear

AACTACOMERCE DESCRIPTIONS OF STATE STATE AACTAC TO SACCOME AACTAC

WO 98/16645 PCT/US97/18214

172

CACA RACE	$C(\mathcal{H}_{\mathcal{C}}(AG)) = .$	G. J. TACTOC	CONTROLATIO	GAAAGTTITT	$= \psi_{+,\infty}(W, L(\mathbb{R}))$	(+   } (
CGGTTAACCG	GCCAGAGGGT	CTGGACGCCA	TEGAGGATCT	GCACATCTGG	ACCGCCGAGT	960
CGGTGCGCGC	CGACCGGCTC	GACTTTCGGC	CCAAGCACAA	ACTGGCCGTC	TTGGTGGTCT	1020
CGGCGATCCC	GCTGGCCGAG	CCGGTCCGGC	TGGCGCGTAG	GCCCGAGTAC	SGCGGTTGCA	1080
CCAGCTGGGT	SEAGOTON 3.1	CTCACCCCIA	COTTGJCGG:	GCCGGTGCAC	GACGAGGCC	1140
ogopou ngov			ADDITE	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	SCATCCCTT	1200
	TWIA GUI W	1-14-1		(171.7 <b>2</b> 1363.7	T. T. COTTOC	1260
I o - STOMAT	Million of the state of the sta	A SW AKS B	i waliganya	GUMATANT .	1 Figura soca	1320
adecotado i	MACAACCI					1 5.2H

THE INFORMATION FOR PERSON SHIPLE

### . F REQUENCE CHARACTERISTICS:

- (A) LENGTH: off base pair.
- (B) TYPE: nucleic heid
- (a) ATRANDEDNEAU'r minite
- 113 TORYTOWY: Time or

# A. Symple of Solity to W. A. Saleste

(x)) SEQUENTE DESCRIPTION: SEQ. 10 NO:155:	
GAAGACCCGG CCCCGCCATA TCGATCGGCT CGCCGACTAC TTTCGCCGAA CGTGCACGCG	60
GCGGCGTCGG GCTGATCATC ACCGGTGGCT ACGCGCCCAA CCGCACCGGA TGGCTGCTGC	120
OGTTOGOCTO CGAACTOGTO ACTTOGOCOA AGCOCOGOCOA ATGACAGGO	180
GGGTCCAGGA TTCGPGTCCA AAGATCCTG TCGCAAATCCT CCACGCGCA CGCTACGCCT	240
ACMA MUNACT TOORDE NACE ENTER ENTRE THAN IN THE GREACE VEL TITTOGECOGN	300
HARTATTAT : GROTORIGH - FILLMARINA GUATORIGGA TTTOROGOGO TURGGOGCACT	360
TOUR TOOTER TOO TENTER TO EVENT UP OF THE ARATEMETER OF TAKEOTOCTOR.	420
ADBOTT FOR LEGISLEY AND AS A WAS TO AND STORE STORE ANA COORDINATE.	4.9
ACCIDITAÇÃO E ON	497
(2) INFORMATION FOR UEST 10 NOTICE:	
THE TEQUENCE CHAPACTERISTICS:  (A) LETSTE 63 and contract (P) TYPE: amino wid	

# A ATTEMAT CANADISCIPLINE SECTION 1999:

-0% DURANDEDNESÖ: US USBSGAT: Linear

The wave too and be wall of districtly what has been districtly the first first

	2-47					u* e							115			
::: y	' Âsp.	Al i 115		Val	117-	Gin	nii. 120		Ain:	Fre	Thr	Ser Lab		Tyr	Le	
Trp	Arg 130		lle	Met	Pro	His 135		Glu	Gly	Leu	Gly 140		Leu	Val	Al	
Cyp 145		Letu	110	Ğly	Mot 150		Ala	Ser	Asp	Lys 155		Ser	Pro	Ser	G15 160	
Pro	Val	Vi-j	ryo	::•)   6:5	Tv:	477	111:		Arr 175		Pho	len	Phos	5.5 175		
Ггр	Asp.	Λla	1901	Auţ	Letti	diy	A sp	His IF	Val	Val	l e u	V +1	Leu 190		Asq	
Vip	+ - 1 - 7	: 4.	Al a	i • · 1.	1 1		A3] -	;,;	žil i	Artri	::::	Han	Arq	Isti) i	Δra	
V <sub>a</sub> 1 i	Gli. 210	GTY	Tio	Isla	11	Mast diffe		Ail)i	120		Ti.r		Mort	71.:	T 2 7	
&   σ . 2   γ	Asp	-11	Pres	1,155	Ala	val	ht q	at y	Va.	The 235	Gir.	GIY	ine	Arq	Ser 240	
Free	GIH.	Hly	: [ * 1	Pro 345	Mart	A. 1			Hr.	A a.	I i e	î be-		51u 25¢	λiq	
	1 1	Γ:	115	iver	: • •	i*.				r. i	λj.»		The		Åtn	
i' * .	777	At a	-62 4	Ĭ:	Etier	1.1.	Art.		.8	41.	ă. ţ	asl ()	Ä: 1	l i	Ent	
i	1'+ 1 2 +	111	÷·	mî 1	:	ė · . '	1 :		. 1		* - 1 *.1 *		e.) :	,	T er	
					* =				Ya:	4.						
															ii i	
		·. :	;		* :		* *		٠		.,.					

- Lie Ard Leu Pro Ala Ero Gly Arg Asposer. Un Gly Lyd Gly His Gan 405 410
- Ser Gln Pro Leu Fro Ser Gln Arg Gly Arg Gln Ile Tyr Val Ala Gly 420 425
- Gln Arg Ser Ser Tyr Leu Fro Ser Glu Leu Val Ala Ala Fhe Leu Trp 435 440
- Ala Gin The Glu Gir Ala Glu Ard He Thr Arg He Arg bed Asp bed 456 436 46
- Tip Acm Arrity: Env Elizer the the ter Let alm alm Arricly Leu 47.1
- Len Arm Ang Fr. The The Br. In Thy Tys Ber Bin Arm Ala His Mot 496 .  $4.6^{\circ}$
- Tyr Tyr Mal Len Len Ala Er Len Ala Anj Arn Glu Gen Val Len Ala 500 500 500
- Ang Leu Thi Per Sla Sly The Sly Ala Val Ene His Syn Val Inc Leu r ≥ 1 r; c = 2 r)
- din App Ser Pro Allesty Arr Arq 5.30
- (.) IMPORMATION FOR JEC II DESIRED.
  - ... SECTENCE CHAFASTERS STICKE
    - A. LINCTH: 194 arct. actet
    - de Trificación acid Convinción descri Control de los controls

ti.										74					튀(:
©1 y	Giy	Leu	Tnr	V.i.) 85-	Азр	1:1	Llys	₹1.	3e1 90	Trj	Fro	Arc	, GIn	Arg 95	(G)
Ala	Thr	Val	Leu 100	Ala	Ala	Va l	llis	Glu 105		Pro	Pro	Пlе	Val 110		Hi:
Phys	Leu	Va1 115		KsLG	Less	See	(11.1 11.1		Arq	Pro	61 y	02 n 125		Pro	Ph∈
Amp	L.v.:: 1.30	Дзэр	V.11	Vil	1	H:.	Arq	Hi.	11;	] 1	A.a 140	De a	At i	ÁI I	Зeт
o.11 .45	Thr	T.e.u	:41 u	Hip	Chr Phr		H i M	±;].ψ	Ara	Arq		ji7 ±i.	Ari	In to	Λr.:
11.13	Ai i	1.5	žsty)	Asy-	/61 :	¥ 1.•	h.		An q 1 Te		l:	17.54		Ser 175	Va !
A.A.	Mest	*****	Vall Per	programme (	i i	Val	1, 11	Alar 14		At I	Ata		Fr.	Val	Va1
Gln	H1.3	61n 195	Туз	His	Y-4-,	Tig.		ii u	Vil	F1.12		11. 29%	Pr	Giu.	Arg
#?u	ain 216	l.V	**************************************	,	Leti	 215	ŽV I	::	Abt		A1a 220	V-r]	GI y	Stot	Ārq
I:;	A. :	11.1	1701		Arq . «	$\tilde{\Lambda}_{i}^{j+1}$	Al.	Hiri		Acq.	ligt.	i de	Ali		Hio .147
۲.	11 *	÷		i .	17.		Ϋ,	Art.		Ā ţ	.° +1	<b>5.</b> :	1:	***: ****:	
h: :	Tar 3			•	i.				-:	i. :			1:.:	* ·:	
÷ 1		• * 1		٠.	, 1				•		:				
		٠,													
	6.														

WO 98/16645 PCT/US97/18214

•		$\neg$
п	- 1	- /

ATHAACATUT CGTCC	GT-KIT - FORT TECMAS	· · · · · · · · · · · · · · · · · · ·	37017003033	CTACTOTO	$\dot{\ell}_1$ :
SCCATGCACG CGATC	GCCGG TITCTCCGAT	anaraaa. w	MAGAGCTGC	-GGTAGCGGA	120
ATCGCCGTCT CGGTG	SATECA CEEGGEGETG	ACCCAGACAC	CGCTGTTGGC	CAACGTCGAC	180
CCCGCCGACA TGCCG	CCCCC GTTTCCCACC	CTCACGCCCA	TTCCCGTTCA	CTGGGTCGCG	240
GCAGCGGTGC TTGAC	GSTST GGCC				264
(D) INFORMATION	IVŠK DET ID NO:158	<b>.</b> :			
(A) LE (B) TY , 4 L	E CHAPA TERISTE Y NOTH: Illicate p PR: miller ferid RALSE NESS: Jinear PK: YSS: Linear	411.5			

ext. DEGGESTE DESCRIPTION: SEG 119 Notiles:

+,2)	. P. W. C. W. C.	MANGENTA MAGGACIA	ATSANGE ENGINERALI	TA moderna
170	- ET PRICAAGO	1970 (774) 77. 001 (800) 779 (501	TATERANTEN TELATORIE	AABTOGET
1	IF G TATAC	H MACHAL LUHTEG MA	CONTRACTOR OF THE STATE OF	ASTRONICA
	1 1 1 1/2 17. 11.	IIVI ta i sembiai t	TA PRESENTATION S	
,	i i i da · · · ·	. WATER OF THE TOTAL OF THE TANK OF THE TA	emagnest tallamini	t to the
	* · · · · · · · · · · · · · · · · · · ·	ZINDING GAINING	ព្រះ ស្រី សម្បាល់	1 5, 117 7, 1
1			rand and all All collections	Assertation of a
				v 3 - 1
			eu tue	· 10 1
• •		* 2. * * * * *	· · :	

GAGTACAGCA ACATEGORGE CITODA SERVICE TO ESTEMACIT ACESACOGITACE SITUCAAACEG.	102			
CGCACCACCO TCGGTTCGCA CUTACGGACC GGGTCCGACA CCATGTTCGT GGCCCCAGTA	1086			
ACCATCGGCG ACGGCGCGTA TACC3GGGCC GGCACAGTGG TGCGGGAGGA TGTCCCGCCG	1140			
GGGGCGCTGG CAGTGTCGGC GGGTCCGCAA C	1171			
(2) INFORMATION FUR SEQ ID Notled:				
THE REPORT OF ARROTHE DELITED AND A LENGTH: 12T Frank pairs The TYPE: nonless acrit (1. OTRANCEDNESS, single 11. Indicases: linear				
JAGO JEJUENTE DES FIFTION: JEJ JOHN 1866: Flamaggues taung egge Googsvanga agag til em intertagnta geologicag	(./)			
2002 20000 2000 2	12.0			
TOACTCAABL TICTIMOTEL AARICTRIA ARTSONINA CIRTIMOTEN GRANDAAAC				
TINAMATER TERMAK BELUM ARAKAN MENANGUN TINAN	180			
THE SWALL HER RELEGIES TO THE STATE OF THE S				
THE STATE OF THE S				

٠.

179

1 17 17	
* • <b>3</b> ( 4 )	47) 2
	11.7

- (2) INFORMATION FOR DED ID NOT162:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 1439 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear

# wir deluence feldelphiem wer in final.

Offishanana.	GEOGAGGETT	TATAACAAG	A WA WYORA	TWG~FF	Company Control	£ ~ .
	ATTOWN WILL	$\operatorname{GoG} \times \operatorname{Al} \times$	: harris Moriet	COMPLACE.	· ···· · · · · · · · · · · · · · · · ·	1.70
TATE	-BFCTAGGAA	TACTTOCT	H-WARTER IN	TOTAL TELEPOOR	910/0410000	194
ST 30 TOAGTT	STTTWAN TOE	A MAGILATIG	T WAS REPORT	COTOLIACET.:	TGGAGGGG	240
CGATTONS	diffatAtetVo	MOTGATO	altanin Will	CATOTTT W.	MAGA JOGAGA	3(17)
ASTRATON	SW WAY	GG GKWY 130.	74 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	"MT"TAT-#"	CANGAAACCA	₹ <b>6</b> 30 (
TTOWNSACA 1		GETIAN IATUN	APT TEAT	CCACC TATE	GACAACAAGG	12. 1
The Control of the Co	T0 7441 7 7	WINCELLS	$(-1)^{2} + (-1)^{2} \sqrt{1 - 2} = \tilde{\mathcal{E}}_{2}$	1 M. A 1M	1 OTAPA ***	480
h = h + h + h + h + h + h + h + h + h +	promote and an experience	-1 (CTT 122 )	* ******	iann inar	Weilland, "	; :
W 11 11 11	111.			1900 to 1900 to	A. PWOTEN	•
Print DAVID	· · · · · · · · · · · · · · · · · · ·	1 5 1127		37x1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1 ->
				٠.	7	
*		4 * * .			:	* 1

CURRECTACY REGULARISE REPLACED FOR A CORRESPONDING FOR ${\cal P}_{A}$ and ${\cal P}$	į,
GCGGCCAGGG CGGTGCCGGC GGCAGCGCCC GCAACGGCGG CCACGGCGCC CGTGCCACCG 13	20
GCGGCGCCAG CGGCAAGGGC GGCAACGGCA CCAGCGGTGC CGCCAGCGGC TCAGGCGTCA 138	3 ()
TCAACGTCAC CGCCGGCCAC GGCGGCAACG GCGGCAATGG CCGCAACGGC GCCAACGGC 143	3 (4
(2) INFORMATION FOR SEQ ID NO:163:	
(if DEQUENCE THARACTERISTICS:  (A) LENGTH: 313 name pairs  (B) TYPE: nucleus actif  (C) STRAUDEDINGS: single  (L) Translow: linear	

# RE CENTER WE REPARTED THE BAY HE BENEVAL

APADOGRAM GEOGRAFIE TOTOGOTON TEATT TO OUT TEAT AND SOCIAL AND SOC

# FOR SEMANT NOR BODY, IN BOSTORY

(:) SEQUENCE - TEARACTERISTICS:  (A) LENGTH: 392 Fase pairs  (B) TYPE: nucleic acid  (C) STRANDEDNESS: single  (D) TOPOLOGY: linear	
(xi) SEQUENCE DESCRIPTION: DEG 10 Not 165:	
GEOCUTETETE GENETICA A EL CERENTATTO EL DACETTO GEOGGECAAT ANGUARNICA	(,
AGICCTACTA TTIACCITÜ JAGGACCCCI GCATCAAGCT GEGGGTCAGÉ GCCAAAGCAA	7.21
ICAAGOTMAT INDA MEGRHANI ELEKATMHA ELEKATMHA ELEKATMHA AREJOOMELEE ETABGAMEEG	18:
CONTRACTOR TO STORE SECTION AS THE SECTION AS THE SECTION ASSESSMENT ASSESSMENT	, 40
CAMARAGERAT GEORGICARO MACTERSEN MACTERSTAR MENERSTES ANGGCAGGA	361
CONCERTATA I CERTUTIARII CERTOTORCE INGURANARIA IN RIVATRACIJA IT MINGENATO	360
Sucharas e enambastele aa bafindtea era	392
(D) INFORMATION FOR OF, IF Notion:	
All DEROTER CHARACTERICTION:  All Expoler formation:  bl TYPE: protor mend  DEFARCHMENT: einote  de d Holero: Turear	
as a significant statement, and seems	
with the following the state of	

ACCUSAGGIG COGGCGGUTC FUG GCGGA: AAC STAIC F UTSCITCITIT COGGG 53%

(2) INFORMATION FOR SEQ ID NO:167:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 690 base pairs(B) TYPE: nucleic acid(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

UKUN SEÇTENTE DELURENTE NE SET IN MARIETE.

CONSACCE CON CORRESPONDE A CONSIDERACION CONTRACTACION ACCIDADOS ACCIDADOS ACCIDADAS ACCIDADA ACCIDADA ACCIDADA ACCIDADA ACCID 1.73 CHOCOTOGO CAACCHIST GURGELIN GELANDINA GARRESTE COMPITAT . . CHARGOGAAC CEGAARGOA TÉREBAACO DE ENCORA DEL EMPROCGAACE ACCGOMACEO 186 CACGAGGTAC CCCCGATGTGC GAACGTCGTT ENGONTGTTG TCGAACGTCE CGCCTCAGGT 240 CAT THYOGAT ROOMERS OF THE WARRANTA WILLIAM THE WOLLD THE WOLLDOW CHECKERACOT. 4000 PRAGOCIOTO TOCCOGCANO IGUTERACIONE PROGRAGATO CASCIGOCOS AACCOGCCCA 360 PERCENTION CONTRACTOR TO PERCENT TO THE TRACTOR OF SERVICE AND CACCAG ACHAILLEA AH WALTAN LINESHEEL TRANSPERS BRANDER BRANDEN THE YOUR 100 AND A COMPANIES OF MARKET OF MICHAEL CONTRACTOR AND COMPANIES OF MARKING A REMAINS FROM AND THE STATE OF T 0.11 Beer the court in toward in the contract of the magazawa take

A Section of the desired control of the control

WO 98/16645 PCT/US97/18214

## 183

A. Towns		Sir Francisco	AAD AAAA C	11 #4 <b>AA</b> 75 ****	* 18 1 x 1 x 1 x 1 x 1 x 1 x 1 x 1 x 1 x	(5)
TGGGCACAG	CGGGGGTGGC	GGCVACGGTC	egalocaticate	TANGSTA T	COACGTGGCG	120
GCTCTGGGGG	CACCGGGGGC	GACGGGGGA	ccagaagaa	TGGCGGCCTG	TTAATGGGCG	180
caggagaaaa	GGGGCACGGT	GGCACTGGCG	GCGCGGGCGG	TGCCGGTGTC	GACGGTGGCG	240
gegendoegr	a geologices	COCGGCGCA	ACGGCGGCGC	CGGGGGTCAA	GCCGCCCTGC	300
TOTTO MORE		DETGTAGETER	12/3 y 1-2 x 2/1/2 x 2 x 2 x 2 x 2 x 2 x 2 x 2 x 2 x 2 x	WikiGATCO.	GETEGEGGC	360
or and a	. 47.22 . 11.22 .(	A	Tollistomiczki			40
ARCHILL LES	MATI'N FWE SE	D 10 10:16	<b>'</b> :			
.:	FACENTE THAN A DESCRIPTION FOR STRANDER (TO STRANDER (TO STRANDER (TO STRANDER (TO STRANDER	4+# klast pla orlosis actid NESS: Pingl	() ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )			

.

# C. SECRETE PROCESS IN NO. 11 (1990) 11 (1991)

75 - 10 - 10 - 10 - 10 - 10 - 10 - 10 - 1	C. 30737 TORGET MA	: NW1 5	13101111111	$AA^{\dagger}AACATerr$	<b>,</b> (8)
		and the second second	T. WW	E. WEWWE	1.5.
AM, Mr. j. mm.	nder tymaer.	American		TABATATA	;=,
	ningh with the	r. • • • • • • • • • • • • • • • • • • •	. :: 25 - 1 - 20	. ; . IV., MYX,M	14.
1 10 m . 1 11 - 11 m	tinin na ana alumbi		1.::	of the children	
				, * ·	
	<i>i.</i>				

ing affilia in Auropia, bu bus 🕸 s

The second secon

(x.) SEQUENCE : ENGRIPTION: SEQUENCE NOTE NO.	
GGTGGTAACG GCGGCCAGG3 TUGCATCGGC GGCGCCGGCG AGAGAGGCGCC CGACGGCCCC	60
GGCCCCAAT3 CTAACGGCGC AAACGGCGAG AACGGCGGTA GCGGTGGTAA CGGTGGCGAC	120
GGOGGCGCGG GCGGCAATGG CGGUGUGGGC GGCAACGCGC AGGCGGCCGG GTACACCGAC	180
GGCGCCACGG GCACCGGCGG: CGACGGCGGC AAQGGCGGC	219
(2) INFORMATION FOR CEC ID No.1771:	
(i) SEQUENCE CHARA TERRICTICH:  (A) LENGTH: 494 bare pairs  (B) TYPE: Lundlein ariss  (C) CTRANDEDMECS: simple  (b) TOPOLOGY: line ii	•.
(xi) : FQUENCE DESCRIPTION: SEQ 1: NO:171:	
IARUTECCORT HANDOÙSTA AGNOCEDUA NA HANDONA E ENNERGARE HANDONECCO	•_^_

ET CATOMONIA		E . 1 . 170 . 241				
TR -CITCON	HANT KAUMITA	AGE FOR JOHN A	ha Hasaniya i	100000000000000000000000000000000000000	A Mary 10 tooks	•=^
JANCAGTTO:	m m (mm)	(1227) W (2.18)	*111 110 10	-510157115913	TO SEE VICEOUS	1.10
75 7 3 7 7 7 7 7	7777 (1767A	Act in program	> 1 * = 10 - 45 *	11 51 12 A 11	anagan KWV	150
7 771 Mat 2 1,37	#*** :: * * ***		ý., 1 · · · .	42.11 110 <u>1</u> 1	ning shaaring.	- <b>1</b> 1*
77 77 77 W	11111111				Aprila.	
WWW.	;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;	**************************************	F		11 - 1 - 1 - 1 - 1 - 1	3 + 1
HMA NA MININI	Lukerini esti	////**********************************	11. 17. " . 17 . "		****	:,
						1

(X1) SERVENCE DESCRIPTION: SECTIONS: IT NO: IT S	
GUGUCGGTOG TORUGAGORE HARRTTTUA ERICHOGOACH OGGGGGGG FE BREGTTGGGG	·
TTGGCGGCAC CGGCGGCCAG GGTGGGGCTG GCCGTGCCGG AGCGGCCGGC GCCGACGCCC	120
CCGCCAGCAC AGGTCTAACC GGTGGTACCG GGTTCGCTGG CGGGGCCGGC GGCGTCGGCC	180
GCCAGAGCGG CAACGCCATT OCCGGGGGGAA TAAACGAC™C	2.20
(2) INFORMATION FOR CEXCID NOT172:	
FIR CEQUENCE CHARACTERICTICS:  (A) LENGTH: 388 Face pair.  (B) TYPE: northet and t  (C) CTEAMER MEGS: childe  (I) TIN GROOT: linear	

(XET LEGISTER DESCRIPTIONS LEGISTATE CLASSIAGNAS LOCANCISCO 60

FERDA FORTE DESCRIPTION CONNECTE CLASSIAGNAS LOCANCISCO 11.00

THAT THAT I A FORTE AND CONNECTE CLASSIAGNAS TO MARKET CONTENT OF TOTAL CONTENT OF THE CO

TO STATE OF STATE OF WAY.

TANCO AA D	$::\mathbb{R}_{+}MG\mathbb{R}_{+}$	Www.naccast	As a summa	11111	:: :::::::::::::::::::::::::::::::::::	13
GCGJAAACGG	COGAAAACGC	GCAGACAACA	CCACCACCGC	caccaaccaan	ACCACAGGCG	240
GCGACGGCGG	GGCCGGCGGG	GCCGGCGGAA	CCGGCGGAAC	CGGCGGAGCC	GCCGGCACCG	300
GCACCGGCGG	CCAACAAGGC	AACGGCGGCA	ACGGCGCAC	CGCCGGCAAA	GGCGGCACCG	360
GCGGCHACGG	THEACTETCA	GGCACCA"	or emedical			400
i. THEORMA	TION FOR A	, in Neil	:			
.: , YE	CHUCE - THAE	AMTERISTICS	·:			

(A. LENGTH: 5.35 have pairs
(B. TYPE: hypers of r

CTEANDEDWELL: Christ
(CTEANDEDWELL: Christ
(CTEANDEDWELL: Christ

(x) SEQUENT DES WIPTION: SEQ II) N :1/o:	
GRANCESTEE TECCECERS CAR TOTTON GRANCESTAGE CONTRIBUT GOODTESTEE	ψi
TTGGCGGCAC CGGCGGCCAG GGTGGGGCTG GCGGTGCCGG AGCGGCCGCC GCCGACGCCC	120
CCGCCAGCAC AGGTCTAACC GGTGGTACCG GGTTCGCTGG CGGGGCCGGC GGCGTCGGCG	180
GCCACGGCGG CAACGCCATT GCCGGCCGCA TCAACGGCTC CGGTGGTGCC GGCGGCACC	2 3 9
(C. INFORMATION FOR SEQ ID NO.177)	

(E REQUENCE CHARACTERIOTICS:
(A) LENGTH: Gibliote parts
(B) TYPE: nucle.c acid
(C) COFARMETHERS: gandle
(C) TOPOLOGY: Linear

TWO SECTION IS INCIDENTED BY CITY IN THE

AGRAMATICITÀ D'ORTRERO ACOSTROLLO CUERTO IL CARACTO AL MARIO MUARAGORI.

DOCAACAGORI STOTOGORIGO D'ACOMATON D'ALLEGAM D'ANTIGORIGO ACOMATON DE COMPANDO ACOM

ACAMO - LOCACAGOS OCOCOS DATOS

## (2) INFORMATION FOR SEQ ID NO:178:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2138 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: sinole

(D) TGPOLOGY: linear

·*!		THE THE .	ee ib Shila	n:		
CJ;MC(M),	The Contra	*************	17/41.5111	ATTOX (T)		ij.,
OGAGCIAAA E	OR TANK IA	Time	·W. C.V. bear.	241 - A. Marie 11 - 11 - 11 - 11 - 11 - 11 - 11 - 11		12
AT IAMOGGG:	GGCATCAAA?	TARRAGES	J. CHTTCMCT	TTAGENGAGIA	TAATUSCTAT	180
AGUAUTAAGG	AGCATHATON	-JATAT-JAC'; ·	ATT TEMORY	CTOTOTO	MICAGCAAG	24(1
AAFTTTTAFA	nWilki mwy i	"TAGRIPOLIA"RI		nuwinézenu	ACT M. (C.C.)	5 . 11
-YATCA(Á	GLOGGAWGL.	ACTIONSTITA	AAAACCTCCCC	VVVO.II	TATTITOG	Žn0
no ananar	eral WW. 7.	-1112 Table 1	3.7 s OWAR W	1: ' ( A' ())	Teach Actual	4.1 -
v garana	111111111111111111111111111111111111111	71777	ALL TITLET WE	iw jari	A" " " " " " " Title"	10
AMACIA TIL	1966-61	70 - 75 i - 35 i	W.J., M		111 0/2 1/2 111	
1.2.1	AA TOWNER	··· a 1 1	· · · · · · · · · · · · · · · · · · ·	Lawest WA	TO MORANTA	,
n man, ian ir	17000 8 4.2			*1.2,* : ** :	1.1111	
				· (		
			E		********	- :
		, ē: · · · ·		1 1 4 4		

DERRIAT I	. Varanta	A.C. CITTER	' CIAN DOM''	t the part of	· Farmonia	11.0%
CTGATACTGC	- GRONCAGOTG	ACGTCCCCTC	GTTGGGAAGC	- GCAGCGPT(	Tallycogang	1260
TGGCGGTCAA	AGCGGCATCG	CTCGGTGGCG	GTGGAGGCGG	CGGGGTGCCC	TUGGCGCCGT	1320
TGGGATCCGC	GATCGGGGGC	GCCGAATCGG	TGCGGCCCGC	TGGCGCTGGT	GACATTGCCG	1380
GCTTAGGCCA	GGGAAGCGCC	GGTGGCG6CG	ccacarraca	CGGCGGTGGC	ATGGGAATGC	1440
CCATGCSTCK	316-10CAT 1A 1	C INDAMIGOR	CONTAINED	CWAGGGTTOT	``ARCAGGAA:	15.0
ACSARGES T	"	WT 16.1 EWT	GWC GMSCC	OT ATTOUT	AATGETTGGGC	1560
37743434749	fWA HWY: "-)	AA ITSASWT	STANGAATTS	GAC CG Willia	7.7900030310	1620
1771-W 11773	:57 : 37 : 37 : 37 : 1		./2 4/2 - 3/3	A 1-0000 AAT 11	AMARAMA	168**
Talkin, III.	5 · 7	z. MAGINTA	W. C. T. WI	off own town the	A. Lara Tra	1 140
	High To the	T MAAGATGB	olo kolgaa i	AND THE YOUR	NOAGEN ST	1800
3 · 1/2 · 1/	GT: WY GWY	AKDĀCH II	J.: L:YC!	transmitter in	"TATAA" CA	18(0)
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	GAPTA POTGA		APTER YALE	TOPHINA	TOMANGANON	1.420
AATTGO TEAA	*ion ATC III	क्षं कृत्रम् स्थाप्तारः	1.47. 177.0207.	0.73.50.50	KCCONAATON	1 145
7 1	13/10/2004/1	A 1 POTON 1	TATE WITH	WATGITT!	A.M.B.C. MITCO	j - 1.4 εγ
1 11 11 11 11 11 11	Webster T.	77. 77. 11. 17.	15.47(1.71) ·	:	ATT CONTA	. * * * *
Table 1 Table 1	THE THE ITEMS	TAND WORLD				

TI DE PRIMER RELATIONS NO SERVICE DE LA COMPANSION DE LA

o de establica e la capación actual de la

		4.					<b>4</b> )					; ·			
hou	Va.) 50	Leu	tion	Ala	Anp	(1) (1)	Met	Arq	Giv.	Tyr	1a	i hija	Äla	e Giy	/ A1
Lys 65	Glu	Arg	Gln	Arg	Leu 70	Ala	Thr	Ser	Leu	Arg 75	Asn	Ala	Ala	Lys	A1a 80
Tyr	Glv	G¹u	Val	Asp 85	Glu	Glu	Αla	Ala	Th:	Ala	Len	Asp	Asn	Asp	Glv
Glu	Gly	Thr	Va1 120		$E_{-1}^{1}$	11. 1	sher.	Al i	«Hy	Ala	Val	4.7	Пу 110		tier
Seet	Al i	3112	Leta	Thu	Asp		110	At 1	Vil	Ala	Thr	Ala Las	.;1у	din	Pro
Ach	11.0	Met	ž.Ų	10001	1.7.2		Al a		Ar q	Eyn	1+1		1:.;	J.v	Añr-
145 145	Miy	Ala	alera.		Al: 1°,±1	Hir	Pile	ÀГа	ΑŢ	1. y 11.6		A	thr	Fine	Auth Turn
bedi	The	Len	G.n	Gly los	Anjv	Va1			11169		G) y	F'hee		Acri 175	
-1-1	HV	201	Asta Cor	Αlτ	7.1.1	A. i	***	ř	ř. d		1 < .1	Aust	Gra 196	Glr:	Aru
(1)*.		1 1 2	1 1	.1:	ide i	1		. '.	٠	75 I .		i iest Z	A. +	i ya	( - 1 fr
Fu		14.	T il	мlт	-:1:1	! ···.	14	٠٠.	1	Ay a	`.÷ :	Ac. 1	( ):	11:	Fr
	.::				2 * 1 5	w	e 1	•	3 :						19 24 c
			ž ,												
												:			1 -

Ala dia Leo Thi Ser Ala dly Ang Cla Ala Ala Leo Ber Gly Asp 340 346

Val Ala Val Lys Ala Ala Ser Leu Gly Gly Gly Gly Gly Gly Val 355 360 365

Pro Ser Ala Pro Leu Gly Ser Ala 11e Gly Gly Ala Glu Ser Val Arg 375

In Alastiy Alastiy Asporte Alastiy Leustiy Cinsby Ang Alastiy

ony diy Ala Ana ieu oly Ciy iny aly Med iny Med ino Med Giy Ala 4 1.1 410

Als Bus Gir Gly Gir Gy Gly Alallyr Der tyr Kly Der Gir Gir Gir Gi 4.74 4.30

Any sid Alo Des Cyr The + 5 Asplicate Alo Tep The SE, Ala Valleria 43%

GLy Act Aru Arı Arı Gu Aq yer iv. Gu Yer Ly.: 402

ONE DEPORTATION FOR DEPORT IN THESE

( ) JEQUENCE CHARACTERISTICS:

(A) LENGTH 177 amin. and:

F. TYPE Samine Two is the Property of the Canada Mariana

The Table 1 with the second section  $T^{-1}P^{-1} = \mathbb{R}^{n+1} Y$  . The section  $T^{-1}P^{-1} = \mathbb{R}^{n+1} Y$ 

Ang Asp Clin Ser Len Len Len Len And And Gly And Val Asp Len Asp 100 105 Gly Gly Gly Arg Leu Arg Arg Val Tyr Arg Phe Gln Gly Cys Leu Val Val Val Pho Gly Gla His You Lou Arg Pro Lou Lou 11 - hou Arg Val Blo Arg Slu Am Fol Val Ala Sy Arg Ara Val The Arg Val Lyo Pro-The Glu Free Act Tv: Val Fne lie Ser Ara Met The free Pro Ser Pro . 17.\* Him Va. And let with App The Letter Letter Lip His Armitter Alla 164 Gin Phe Gly His Val Gla Ty: Pr. Lett Pr. Lett Lett He. Ali Ara Ser 1.35 Eric Leis Asp. Val. Alia Leis Com and Som Val. Him Der Val. Eric Eric Ele And Ly. Vil Angled Carrelly, whater Tall All And The Contract Pre-.\*\*\* 3 The Africane file (In the file of the file of with the one file of yield). THE PARTY MADE TO SERVICE No exercise and a second second Control of the second

- For Asy fire derived for version set her laws Asia values of the styr styre  $2\sigma$
- Ala Glu Ger Ser Ala Ser der Ger Ala Arq Ser Gly Asn Gly Ser Arg 35 40 45
- Trp Thr Ser Met Pro Ser Gly Th: Arg Pro Gly Pro Arg Arg Ala Thr
  50 55 60
- Set And Asp Acp And And Set Ala Thi Set Val Lie Printer And And  $\frac{1}{80}$
- Ser Van Ala Fr. And Ala dan Phe Cly The Ang Leo Ala Jer Has Ang Eo To ge,
- Ala Ser tin der Amm All. dyn 1. Val Ara Lie var ihr ber Ala Ber 140 - ten 110
- CDy Arithmodice for the front of the Val Arithmodish type Val 115
- Ast Lys Ash Gly Arg Arg Typ A. . Set Bly Tyr Arg Arg Lou Ann Arg 130
- Ala Ari Mer Mer Mer 118 Ala Ala Ari My Ari Mir ile Miy ile Pho 145 - 150 - 150 - 150 - 160
- And Ari Set Ard Tyr Con And Jes New Ard Year Den Tile Ann Jes 166
- His values His Survey Value 18 Survey Values (1888) Also 18

. THE PREST WE HAVE, IN MALLS.

THE THE MEAN THE PROPERTY OF A COMMON TO THE COMMON THE PROPERTY OF A C

		31					700					45			
Arg	Ash 50	Free	Ārq	Azu	.303	.5 i 5°.	ža g	Ārq	Авр	Ala	61 u 60	Asj	Arg	Arg	Va
Ile 65	Phe	Ala	Ala	Thi	Leu 70	Val	Ala	Val	Λsp	Pro 75	Pro	Leu	Arg	G1 y	A1 &
Gly	Gly	Glu	Ala	Asp 85	Gin	Leg	114	Asp	Leu 30	Gly	Ųa I	Сув	Aro	Ara 95	Glr
Ala	11.9	Å; 1	V 11 1 (0	Ist 1	At 4	11.7	H:	1.19 1.14	in u	His	His	Ārq	His 110	Arq	Ніз
Hri	11 y	Ala !!	Aia	1:	ži, (ļe		Ar 4	ñi -1	At 1	ÁΩα	Ar	His 124	Arg	Arq	Va;
4.1.	(4.1. 130	Him	Ar į	At i	· .	7. i. 1 •	+.f I	∵ e i	7.: i	⊆In	:: :46	तंत्र	Arg	7 y r	Val
31:. 14%	Thir	Al:	Him	Hitt	Ist 1 1 - 1	Ar i	File-	Ali	Atst	11:	Act; x	Arq	Val	hi-i	His
Halls	Viii	Art.	"ау	Pro No.	1 - 1	Art.	H	6.1.1	An 4 100	Arq	Arg	Val	Tyr	Arg 175	СТУ
Ar i	11.25	361	Hly lh÷		eri y	11.7		1 1	i i	t' y	G) y	Ala	31 y 190	i. Des <b>r</b>	V-1-]
117	r. ;		Δ÷:												
1215 F	MAT I				• •:	: 1 -	:								
		net. Pyl	 1 :	1 /	.F: .	: :									

Ş......

- 4.6 And oly Gry Lyn Amp Him Fribles I we had Arm Valoring Rep Sty 50
- Ala Thr Pro Trp Val Thr Trp Gly Glu Phe Val Glu Thr Arg Met Leu 65 70 75 80
- Ala Glu Tyr Arg Asp Arg Arg bys Val Pro Ile Val Arg Gln Arg Ala 85 90 95
- Alic the Glo J. r Lea Arm A. r Arm the Arm Lea Arm Tyr Pro Lea Alm 150 150 150
- His led and Ir. The led Ser Thi His Giu Ard Asp bed Thi Met Gly. 119 125
- Mr. Who will also of a less in Chap Alas ten Wal Throtte Ara Win Gig 130 130 130
- Fin Ata L to let ely Liq Wila Art Trp Test Ala Let Less Val Pro Ami 146 - 181 - 185 - 186 - 187
- Were Also Artic Gly Also care two Artic Levi Gly The Thir App. Val. Also 100  $_{\odot}$  175
- App Lea Ara Ser for Ara Sto N . Are Ara Ari Sty Fre city  ${\rm Arg}(v_a)$
- The App STy I of App Val His Lear Data Fro The Fig. App Leaf Alia App. 197
- As placed for As placed Association of the Associat
- The Alle And Acad Type a land of the South Review of the Stationary
- (a) A disemply As a Asia of the Asia of the term of the same of the ways of the action of the same 
T and T and T are the second T and T and T are T and T and T are T and T

.

(13)	TYPE	: :::::	ينيم إح	4 11/1

- (C) STRANDEDRESH: single
- (D) TOPOLOGY: linear

## (x1) SEQUENCE DESCRIPTION: SEQ ID MO:184:

CTCTTGCCGA	TTCGJCACGA	GTTGAGCAGC	CPAAdGGGCC	-attracocava	GTCATCGAGG	66
CAPI DECOGA		SWMGGGA	AG WAATOVA	$(ACCMCC^{\prime}A)^{\perp}$	AACAGCCTGT	120
a pateratal.	DAADHTTTT	AATTAG DESC	ST POCSANT T	STTIGHT:	STACGCAGCC	180
LindoRadVAL	gorran ord	CTACATOAS	ACTA COMO	ornachtair (	TTWAACAAGA	247
A*γ*! 10 33.W	GT (CACCGA)	Matter, take	ATMONIBLE	CONCERN TO	'AACTONA I'',**	21,00
AGCVATT CGA	CAGCTTAGT	HE GIOLOG	ALTERNATION TO SERVICE SERVICES AND ASSESSMENT OF THE SERVICES	: "WCAAGAA"	CG MAGGTOC	₹+-()
THACTCATIM.	NETCAATAAT	missionus.	$T^{*}(X)\cap X+Y_{X_{m}}$	migarêwa	COMMATCOST	420
TRAIG SIT	GCAGAGCT.Tr	CTOCACATOO		elenogerogger	ATTAACCAGC	450
CTTA-CAT:	BATATAT KUT	manipully of		stiffcV.AtVY).	TT YPWAACC	540
SWINGART	CATES MAGE	TRUATTOTAL	A STANDARD	201 10 1mm 7.5	CANGAGTORG	600
. A. MWA.L'	77-11-25-775	("" 1 1 1 1 V.	and a second of the second of	WI WARID.	AMTANTITY	total (
Comment Band	Wolf Fig.	AT AT UT	NAME OF THE PERSON OF THE PERS	**************************************		
Who contains	NWINTER T	AA * 11 - PAC.	) % ' <b>A</b> '' 'V. '		AL TITAPET	٠
$H_{2}(X) = 0$	distant	PERM ATT	Alloration.	and off the	tation of	** **
				**.	t the second	
						••
<u></u>		* * * * * * * * * * * * * * * * * * * *	9 10 (3)			• -

Trocerecti	Same Standards	CAROSTA	CONTRACT	HIG WY WY	FACCGATGC	1440
AATOGGGGGT	GCATATCATO	CCACAGGGCC	TOGGTGCCAT	GCTGGCGATG	CCGATCGCCG	1500
GAGCGATGAT	GGACCGACGG	GGACCGGCCA	AGATCGTGCT	GGTTGGGATC	ATGCTGATCG	1560
CTGCGGGGTT	GGGCACCTTC	GCCTTTGGTG	TCGCGCGCCA	AGCGGACTAC	TTACCCATTC	1620
TOCCGACCOG	COTCHEAT	AT THE CATER	GCATGGG TG	CTCCATUALU	CUACISTOGS	1680
CONTROL	EAGACETT;	COLONAL!	ADMICONTE	(10077) (175 );	STGATCAGCG	1740
COMPONANCE	THE STATE OF CONTRACT CO.	TOGATAGENIA	1. 1972	True promise	TYPECTACC	1800
ACTTOAATOA	De a langue d'	ATTENANT	PAARWAAP	TRATEM	PAGAGAGTG	1868
ir yn Maai	· [ :- : :		17000 170 170 170	YYYYXX YY	AACTTTTGCG;;	1925
STAATET	C "ATGAC "T"	Table Menoral.	ACANGROS	ATTOUTOATA	S THE TOTAL	1980
TACTGSTCT	NACCOLL-ME.	\$15 TENER; AT	Ty Marin WA	AMARINA KONT	AGTCATTGA	2040
·MAČAC-IGTT	BOTATO, NOVA	TOA STOP 16	1.7			2000

# TO STATE SHAPE OF THE PROPERTY 
# (1) SEQUENCE CHARACTERISTICU:

At LEMGTE: 1909 kare pains He TYPE: no deno unli-t. STRANDELARIO: Banane L. 198 (1997) rinear

THE REPORT OF THE PARTY OF THE PARTY OF THE

文 (本) ない a colonia (本) a colonia (精) (\*\*) a colonia (\*\*) a co is the little to the first terms of a wear to

CONTRACTO	STEETTIT'R		$= G(M^*(M^*))^{\frac{1}{2}}$	) AGAANGANG	TOGCGGCGAT	540
GAACABOOCC A	ACGGCAATCA	COACCACA	F ATTTUCCA: 1	CATACCCTC	COTACOGCTG	600
COCCOCCOTT (	GTCGATCGG	TOGOATATO	ATGGCGCCGT	TTAACGTAAC	AGCTTTCGCG	660
GGACCGGGGG T	TCACAACGGG	CGAGTTGTCC	GGCCGGGAAC	CCGGCAGGTC	TEGGCCGCGG	720
TCACCCCAGC T	CACTGGTGG	ACCATCCGGG	TGTCGGTGAG	CGTGCAACT":	AAACACACTC	780
AAGGCAACG G	STTTTTCAG :	GPACCAGCTC	AğıtıTCGACC	CGCAATCSCI	CGTACGTTTC	840
-W.15000000 A	: Acret 1990	<b>₩</b> 740043017	11111111111	GOTTTCGO	1GAAGCCGAC	900
CARSCARGE T	A STITE OF	CACLOTUAU	Anorganos	GCGAGGT-GGT	CGGTCAAGCC	960
GENATITING "	A PRATOCA	$-1.5(3) \approx M_2(0.5)^{3}$	un onnaaki	.NEWATOTEC	CHAAGTAGGC	:020
GTRAUGRA A	WHERE WAY	M. M. P.		15331074434	** AA. A.D.	
TT YAYARII I	.707 ) 1 4 3A	TAMATAM	r EV Valuet	A 1 19 NO BIA	, Googe Core	1140
	OCMANDO E	COAATCCCOG	AACCAGCA D	CTESTOTCTC	CTGCGATCAC	1200
an and the co	ATOGAGOGT	Figure (AMA)	ACTIVISTOR	TOAACCER	TAP GUACCT	1•••
TTT1117771711	TA NATAAN		OWA TWO	· WAT: ""	misercare.	1.320
TOWARD TOWARD.			M********	* ). (1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1	OCCUAAGCCA	1386
Adiron from the	William Heatter	(r) defit (II.)	A-ATHILTE	" We Willer Will	CHTTTTCC	1440
AT CHACKA A	TTTA YVAT	sayarın ey.	181111111	124711 - 174	7 E W	· 'w.
Altinia * b	1.12 11.11	APPHHI	ada Bariti	1.011.78	11 110 17	1
and the transfer		7. * V***W[ .		in the Public	· · · · r.p.r	1
	e	1.00				· .
					** .	
	: : ".				**********	: .

- (10) OTRANDEUNFRU: Finish
- olo ToPolo Wi Timesi

## (X1) SEQUENCE DESCRIPTION: SEQ 10 NO:186:

CTGGCGTGC:	· AGTGTCACCS	GCSATATGA	TOSTCATIC	AATTTCCCG	: + nacaecaeea	60
CCC CCC	TOCAMITTO .	ACEMONT E	or Reading	-xcavagata:	CAVGTGCGT	1277
GAN AATOR	GTUTGHT	TOTTVAA VA	. Albertick;	TATOURTE	· · · TACTCCCA	18.0
AAPAA SICYA	OTOTAGRAAT		TOUR WHATT	a Walter	TYTTGYTAGCC	240
NO ATOTT	"MGACG"AA."	46770377A4	TO TOWN YOU	n in Willia	o un a pagno	₹170
Transpart St	GOWAATTAG	AAAACCC Ess	British Shirt San 1	SHATTOTA.	"A" ATOTO	(A)
AST HITTGHT	"FETTOGTCTT	A TIADURAT	AT IT NATION	ELL MOVELL	TENTOTAGAA	-4,1-3
SGCHACAACS	"GONTTTEST	TCTTCCTCC		"TMATHMENT	GOTGGGGAAA	4 H C
1.2 relativita	CG GUAATOST			Lo Altan <b>LC</b> errai	anna impera	1.4.
STATOCIATÓ	ATTATGAGAA	ed switten	* * * * * * * * * * * * * * * * * * * *	$M_{i}, \dots, j_{i}, M_{i}$	TOTOGATOTO	600
CARLANTA,	AAMO VILLIJA	. 14.29.139.1	*******	4 17 <sup>4</sup> 0 3 4 W	a waterwe	***
and the state of	A17,11141	1.5. 1.5. P	·	17 V. 11 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	A ' " A " The second of the se	
A+4.11.132		• • • 00 , • • • • •	etin two		j	** -
		** ** 1 ** . *	* (1) *	1.7.1 h. 7	n i MATHEL	4 ;
15/3 - 5/17/64 - 2	11.11 - 11.11			t that any a	era established	•
• •	. =				' =- · * '	

± 00

	(x.) S	EQUENCE 1417	NATION: 3	EÇ He Nor18	<i>7</i> :		
CCGC	CTUGTT	GTTGGCATAC	TCCGCCGCGG	CCGCCTCGAC	CGCACTGGCC	CTGGGGTGTG	6(
TCCG	GGCTGA	CCACCGGGAT	CGCCGAACCA	TCCGAGATCA	CCTCGCAATG	ATCCACCTCG	120
CCCA	GCTGGT	CACCCAGCCA	cosseccits	TGCGACAGCG	CCTGCATCAC	CTTGGTATAG	181
CCGT	ngagaa	ncand mag	GAACTTUTAG	TACTOROGOA	CCACCTGGTI	ACCOGGACGG	241
GAGAI	NOTTO	GERCH CAACEET	$\operatorname{Coloc}(\mathbf{AT}_{i}^{*}(T))^{-1})$	perentica Actes.	AGTTSACCO	MAAAACCAGA	300
nin 1997	""GGCA	GGTGCTTCGG	and maddwd	ADGAGAAAACC	Ca∧raceaa;	ATAGGTCAG	359
1.17 1	DEORMA	TION FOR DE	4: 110 (0):14 °	· :			

DESCRIPTION OF ARACTERISTICS: (A) LENGTH: CO have pain (B) TYPE: nucleis Acla C: CTRANDEDMESS: Simils (C: C-40.000: Unsur

• • • •

TWO STATEMENTS OF STREET AND THE STREET

AA TROOMER LOOFER WATER OF TOTAL CONTROL OF THE ASSESSMENT FOR CAAS PATRIANST PRABATO OF MARKET MARKET AND ARREST A e sparent i di travara i di illariti. Partiti ala citta qui, iti di tripare i f = (1.47 + 2.46) + 0.01 + 0.024 + 0.01 +.14

1	Deeli.	Fire	i leyer	Gily G	· frc	Efr	' An ')		i Val	. îl•	· i	t Ali	i - Phi	A1.	i Asij
G1 y	Leu	Ala	Gly 20	Lys	Gly	Гуз	Gln	11e 25	- Asn	The	Thr	Leu	Asn 30	Ser	Lei
Ser	Glr	Ala 35	beu	Asn	A1a	Leve	Asn 40	Glu	⊜1 y	Arg	Gly	Asp 45	Phe	Phe	· Ala
Val	V + 1	Αrā	Sect	1.01	Ala	lent ci	i tipe		Assi	ñ:a	Log	Hīs	$\overline{v}$ : [ $T$	Asp	Ası
iln G	v;``:.	19.44	73.	A1 :	l≠yr ⇔g		î ye.	Iv 31)	Len,	dales	Gi.	110	Thu	Anp	Ar q
1.01	en.	[]: *	*. *	A up	es d	Ärit	;,- 1 <u>;</u>	Set	Agita ive	Alu	lie	l.n	Hr.	The of	Азр
.4-1	1.001		A. a. I. (	7.1	1.1 1	773 1				A. 1.	77	Å. 1.	At 1		Vall
D 91	1:.:	Hirr Lit	Ārģ)	$\nabla_{\mathbf{A}^{n}}$	Astr		1. H 1. * *		74.2	$\nabla_{\Theta_{n}}$		71.: 125	The	Ízlati	Lou
il.	14: 13:-	āt;	÷ 1++	1 .	î. "j	eli.v 134		9 2 1 - v	. *13	( i i	1.00. 147	H	11	Pho	Pro
7to. 141	1	AT s	Aş.	Æst.		/s.***	· . 1.	l + 1,		H:. ;:::	i٠.	71.7	His		Gly Sur
	7.	, ,	13.33		Pri i	l ji -	* • • • • • • • • • • • • • • • • • • •	í.		Ali	Ārt.	· :		*1. 194-	1:1.
	· e		*:		8 11	Λis	*		žet t	i I	\$ ty		1111	÷	:
				1				t			ē				

	25 <del>4</del> ()					201			7 11 1	• • •	300				. 1 * * !
G1 y 305		şin	Th:	Pro	Pr (- 310		iro	Pro	Asn	Ala 315		Asp	Glu	Туг	Prc 320
1 s.V	Leu	Pro	Pro	11e 325		Leu	Gln	Ala	Pro 330	Gln	Val	Pro	He	Pro 335	
Pre	Pro	P1 +3	il y i46		Anj	Vil	110	Fire 345	Gly	lro	Va!	Pro	14 o 350	Val	Leu
Ž., . i	Ā. i	i i ez	6 I.	1360	1:00	Aist	App.	Ãi a	120	Ala	Ali	11		Äim	Phro
Ar).	77: 336	Mirt.	Ωy		1.41.	1.53.	1	, lest	i i i i	υ· γ	eq.	nta	Thr	Phe	Leu
8 1.++ 38 <sup>1</sup> 4	- 4	9			, 1-1 4(p <sup>2</sup> )		/s.,,,,	At 4	· . 🖓	Shi ana	Mer	h	A-Y	λεγ	H.3
Tarl	Lem	The	1-7 %	401 i 401	1.0	The	Πlγ		11.	I · ·:	Ho	Ala		Phe 415	Val
i i d	h :	19441	(1 p)	Tyr	Ad 4	Thr		H 1.7	Fr"	Dett.	Sie	Asp	Mert 430	Arg	Lesi
Đì.⇔		Αυτ. 4 - Έ	Ara	Alb	Val.	niı	C.:.	Ā. i	n't.	Er t		145	Thr	Va2	Len
	1 1 'a		:	1		4. ;	1 :.	· .			1: 46%.		Pt.		iin
į.			H .	* 1	, (° ).	1:	*	٠,		† ·	. )		: .e.		1 : 4 = 1
	. 7	f	· · · · · · ·	1a 105		*	*: •		1.0		A. ·			ph •	[5.7
	٠												•• .		

640 Oln The Aon H., Ser Shu II.e He Ala Thr Ala Lyo Lyo Val Ala Len 595 600 605 Thr Pro Glu Ser Gly Ala Gly Arg Gly Ala Ala Val Asp Pro Ser Ser 615 Leu Pro Arg Gla Tha Asn Phe Ala Ala Gla Leu His Asp Leu Ser 625 How Ara Lym Ala Va. Val For Va. 11r Ala Thr Ala Leu Val VAL Ser The Loughbourn Ale Ale Fie best in Lyr Cln oin Alader H.b Arg Arr Ala Pro Lea Lea Jer Ala I, INFORMATION FOR SDC IF NOTICE (1) SEQUENCE - HARACTERISTICS: A, LENGTH: 126 amin carri THE TYPE: amin's anii. TOTHAMITEIMESŠ: I\* T FOLCER: Linear THE DESCRIPTION OF THE RESERVE OF that also the pays that day the that the product of a deal what places Mit tall of the Ala Val. Tallor of a week to the only Table. It

The Ari Arq Adj in Ari Co Ari

- (2) INFORMATION FOR SEQ ID NO:191:
  - (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 89 amino acids
    - (B) TYPE: amino acid
    - (C) STRANDEDNESS:
    - (D) TOPCLOGY: linear

ME AMAMENIA TABLEST TABLES, II MARIATE

Les Alastyn Glin Tyo him Arg Ara fyr Ang Va. By I'e olin Pine Arg I

oly fro Ala Sy Es. Val Ala Chi . In der sly by Es. Sy Fro Eag. 20.

The Ala Giu Siv Ary Sin Val Arg Ala Sin (v) Siy Ala Giv Phe ben 39

GE. Ar | Ard Er. Alle Vel Ger. Siv Aver Sep Fred Fred Adm Adm Acad Sep 50 65

iro Giy De Artusa Art Al. And Asputer In Art His Lei Lei L.

 $\hat{T}$  is a limit of the map  $\hat{T}$  and  $\hat{T}$  and  $\hat{T}$  and  $\hat{T}$ 

TO THE EMPLOYED LIBERTY MINER A

- His Leu Ala Met Lie His Liu Ali Win Leu Vil thrown Pro Pro Gly 45 J.J
- Gly Val Arg Uln Arg Leu His His Leu Gly He Ala Val Ala Pro Gli 50 55
- Pro Gln Glu Val Val Leu Ala His His Leu Val Thr Gly Thr Gly 65 70 75 80
- Glu Val Gli Gly Glu Jly Arg His Vil Ala Ala Milu Val Val App Pro (4)
- The Ash of hiller ben Ara Sin Val Lee Siy Pro Air Fro His Asp Lys 1.56
- fr Alp Artony Levely Win

TO INFORMATION FOR SECTION 11941

- 1: SEQUENCE CHARACTERISTICS:
  - (A) DENOTH: 11% aming it is
  - (P) TYPE: amino anid (P) CTRANDETNESS: 1. TOPPOLOGY: linear

WENT PROPERTY IN CONTROL OF COME TO SERVE

And Ala Ara duy Bus Ara dan Serity. The dan deartha of a Harisday

V. T. J. Barrier, M. W. J. Barrier, and S. Salaman, A. S. Salaman,

#### INFORMATION FOR JEG IN N 11941

#### (+) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 811 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (K) SEQUENCE DESCRIPTION: SEC 1: 10:194:

	os IMUs Mo	CAATO CITT	COTTABAGAGAT	(37.37.747 37.0 <sub>3</sub> )	GCCT#GCT-#	TJBCGATGGC	$\epsilon_{\rm c}$
	ti maa prod	MACONING	west with	. I I IN W. T.	102/1/5 . 1	WAMAGGCTC	120
			KAN TOT VAN	$\gamma_{i,1} = \gamma_{i,1} = \gamma_{i,2}$		7 Wy WOCKC	1 ਲ-)
	. Maga sasa	COMMITTED AND	CCASCIONA	4- A4 : 493-1	mannan in	- GOEWGGCT	)* <b>4</b> **
	ATOGETT,	OFFICE OFFI	GGREETA	:::::::::::::::::::::::::::::::::::::::	1° WYWY W	-14:3677736	17) ,
ñ	in total all a all	namena n	MARTEMAL.	<i>z.</i> ::::::::::::::::::::::::::::::::	1	m manager (40)	÷ € •
	CGAÇCI OCU	CSACCCABO	dodana noss	Sample : V	· · · :(:::::::::::::::::::::::::::::::	CINGUISTGIC	420
P <sub>i</sub>	. p.W. i p.W.c.	OMEGANATA	om more to the	<i>7</i> , ************************************	·*************************************	TATGOTGGGC	480
÷,	WWW WILLS	515965W55	attermining t	" "- "AT 12"		ECONTRACT.	:, <u>, , ,</u> ;
:	11.1111111		ett. EMITA 1		114 174 11 1	111777 (P.7777 )	r + 1
	Altin I Wy		· Priview.	***		1 : "370 7	• . • 3.
			;;; .fe		v •		

(xi) SECTENCE, LEG TRIENT DE LORGE LO DE EL PAR

STORREGESAT	· 51000000000	ATGACTITTE	G 7Wr W. 7. 16	COTAGIAGI	GAAGATATCG	'nί
GACTTTGTGG	TCCCGGTGGC	GGGATAGAGC	ACCTGTCGGC	GTTGGTCAGC	GTCACCCGTT	120
GCTCGGACGC	CGAACCCATG	ĊTTTCAACG1	AGCCTGTCGG	TCACACAAGT	CGCGAGCGTA	180
ACCTCACGGT	CAAATATCGC	GEGGAATTTC	GCCGTGACGT	TCCGCTCGCG	GACAATCAAG	240
GCATACITCAC	TTACACGCGA	ALAN THE	CH STITIOAT	::GCTTTCGG:	CTGGTGAACG	30€
TGCCGGTGAA	GOTHUN 750 Y	onwine we	ANYANGA, AT	$-\mathbf{A}.(G\mathbf{T}\mathbf{T}^{T})(\mathbf{A})$	CAGGTGCACG	Soil
AKDADRAKUSA	COGACGUATO	OWNER WAS IN	OPETETOCAN	GGCGTGT56"	GASSTGSTCS	420
A MACHICIA	TIME WELL BY	n atmater c	······································	CAMATOST	FOGATOACCĞ	4-4-5
A TWIGAPAT	MINNITT	·		7.11 3A(137)	TTG-WAGTTC:	540
h-gu sasadan.	CaACTGGG	FULAPTADE	T. 1964, E.M.	CTAGTTTTT	SACTOMATT	$\epsilon co$
PORTAGE AND A	PAME CHIAT		UTWA W WIT	W. MARKET	ACCOGNIGG	មួយពិ
NEC THE NA	Tunton	; PYCEGAAT	MARKET AND	TAAGAGUUGU	TABÉCACAAT	7.20
TCG-FGFCGA	CONTRACTOR	JACAAACH JI	A SAATOATOO	AGACATTOO	GAGCTBAAGT	780
4 70002.TAT	7377 Y . 1911	TOTAL YMTT	MI-WWC-L-W-	MATACUTTI	SATELATOCOA	** -1 (
	TANTA WE	. Will DI	- 577, N FYT N	Wented far	0.1.1.844.0.70	*1,13
Martin Prans	Transfer	v:	Astronomy	tit rankari		÷.,

and the effect of the state of the state of

creeseeeer	GCCACCGTT	· **C**CGFF**c	TATAANATA	n genaamoon	COACCCTTGC	240
CACCCACGCC	ACCGGCTCGG	CCCACCCCCC	-dgacaddaad	CGAGCTGCCG	CCGGAGCCAC	300
CATCACCACC	TATGCCACCG	ACCGCCCAGA	CACCAGCGAC	CGGGTCTTOG	TGAAACGTCG	360
CGGTGCCACC	ACCGCCGCCG	TTACCGCCAA	CCCCACCGGC	AACGCCGGCG	CCGCCATCCC	420
agaagaaaaa	GGCGTTGCCC	CCCTTGCMGC	COTTECCGAA	CAACAACCC	CCGGGGGCCGC	480
aarraaagan	Conditionality		PETONO SEL	.:::::::::::::::::::::::::::::::::::::	CTGCCCCCCT	540
TATES CONTRACTOR	ACTIVATORS	igi - TUMYA	W.C.3337.2	TGCCTCGGGA	TC1GGGCTGT	e () ()
MAACCT CO	$\partial \mathcal{M}(0) \mathcal{M}(0)$	7.10c(C)d(/),	in regression	CCCCCCCT;;	GCGCCGTCAC	660
ACCUATACC	ACCOUNTRA	manana.		AT WOMEN	AGMACCO	
Similar (Max.)	ATIONNIA	4.774000000777	14 WW 1777		PERCHANGE	780
Adeceeant	ACCACUGAAA	1201176017A1	CAN THACA	OTTOGENETE	COGATOTSTA	840
SGAAAGCCCC	GCCTCCGGCG	CCGCCGCTAC	TAACTO ACT	GPTHOTAGET	ACACCGTCGS	9()2
Wattiggaa	A SCANCASSES	= :MAA(7); : (3)	1 777777	$(1+\alpha,\mathbf{k}^{\mathrm{H}}),(2,\alpha,\omega)$	Aimmoundgr	46,00
Court dealer	927377377	making kali	PHILAMPATATE	FOUT WOLV	a pagoyacca	:020
	$\mathbb{F}_{n} = (-M_{n}\mathbb{F}_{n})(n)$	The state of the s	1.1576.33.	than, nami	JUACCEGITCE:	l okt.
Caracada i	AGUGO WILLE	; * [	(-1)(-1)(-1)	and the	THA NATION	1 1
· Marine	*****		*	1	A''' 'A'''	;
	1300500		gr. 17. 15 1	THE SENSE OF	** 1,7, 15, 1 · ·	it o
			* * * * * * * * * * * * * * * * * * * *		*	: .
. *				* * * * * *		
				1.1	+ 0 - + - + - + - + - + - + - + - + - +	:
			• • •			* *

CGGNURATT	$K_{i}^{2}$ . The $i$	ACCERGAN	ARCOTT ATAKE	WANCE WAN	re Cal ACTTe.	1860
CCACCACCTG	Cegna pagea	ACCVOLVVO	AMOTOSM	CRITCOGGATA	V PATATON	1920
CCGCACCGTG	TOCCCAGOGG	CCACGTGACC	TTTGGTCGCT	GGCTGGCGGC	CCTGACTATG	1980
GCCGCGACGG	CCCTCGTTCT	GATTCGCCCC	GGCGCGCAGC	TTGTTGCGCG	AGTTGAAGAC	2040
GGGAGGACAG	GCCGAGCTTG	GRADAGACOT	GGGTCAAGTG	GGAATGCADG	CTCCGCGGGG	2100
AGATGAATAG	GCGGACOCCG	ATCTCCTTG	1 ECTGAGTCC	CTCACCGAC	ACTAGAGCCA	2160
CHICAACKTC	TOPOSTICAL:	AWB 1911.11	A STWATER	C. F. TTT	TTGCACCGC	21.26
dylen Tarry	CSCATA NEED	AFCFCCT.AT	CWICGATAA	CGMOTTOUT	1CGGCCCCAGG	2280
CATCCOTC IAA	CTT FIT HUA	or And Ant	t to Maceur	YYYX YY Y. Y	WANTETACAON	234
· · · · · · · · · · · · · · · · · · ·	With the state of	N. 1975				1 × 1

#### A F INFORMATION FOR SEC ID NOTIFIE

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 376 amino acids
  - (E) TYPE: amuno acrei

  - TO UTRAMPROMESS:

# \* PART \* THE BATTER HE SELECTED IN HER ALL SELECTED IN THE SEL

- grafite Ara 💖 Åra tra i e Arg Ara Sar ole tek (v. Ara Marva).
- ordinate continuous flar control for the open of the top and

- Bly Gly Thr Vil Ala Ala Gly Ala Thr Bry Arg Ari Ala Gly Ser Ala 115 125
- Met Ala Ala Arq Ala Ala Val Ala Ala Gly Leu Ile Thr Asp Ala Gly 130 135 140
- His Ite Cys Arg Ata Val Pro Gly Ata Gly Arg Gly Ata Gly Arg Gly 145 150 160
- ile Asp Pro Val Cys Pro Gly Glu Ala Gly Ala Ala Gly Thr Thr Gly
  175
- Asa Ala Met Asa Gourdin Providing Val Ala Aia Val Thr Aia Arg Thr Lee 180 190
- it (App Ala Tym Cly Bir Ala By Ala Ala App The Ala Val Ala Ala 190 – 195 – 195
- Val Aladis Candis Devise Valds Ins Juy on Ala City Asi Ala 21c 22s 22s
- +High The The Higher Alsa Val Alsa Alsa Val Alsa Are His Pro Gly Arg 240 235 236 240
- Alr Ser Alm Ass Alsoely residences for Associate Arm Ata Val Ala 255
- The Val A. rays Sin with it. Also sly Arm Aim Arm Let Fro Gly  $c_{\rm SS}$  , which will have  $c_{\rm SS}$
- And the Value Ala Value No. Albertain and Alas Fig. (Abs. Ly  $^{\circ}$  And Evaporation (  $^{\circ}$
- $\gamma$  , and the Hammark late and the distribution for the following constraints  $\gamma$
- der Aber by die die treiben die Aerbeere in Armitem Her Die beer

- (P) TYPE: nucleic dend (CV STRANDEDRESS: cingle
- (D) TOPOLOGY: Linear

(xx) SEQUENCE DESCRIPTION: SEC 15 No:198.

GREAMMAND OUTGOODER (	nandoochte.	3480 G200	ACGACGAGAS	CLC EECCAC	€(C)
Websing as well as the second of the second	TA, ' 'AT: :*'C'	Property (	11 1 - 11 0 1 4 TV	GOTHITGAMA	1,*11
COUTTOARC AA EGRITARE A	Armathagas	acom presta	CESTIFITORS	CGCGGGCCCC	180
errowalion oliveration o	TT TOWNS	.::	· AATUT · · ·	instant toWater	1.4%
er din e waker i bowan et nore di	AN TITOTAL	-in -v	MA VIII	WA See Was 1	•
ATTEMACU E SEARCATOS E	ACETUČAA	V.J., (1) th tart	EPARCOT 1755	TOCT 803TTC	$\{\mathbf{r}_{p}\}$
ACCOCAGOO TOAA AACAT C	COTTO NO	AA TA'' II ROCA	HIMACAA 1A	COCCITAGE	4.00
AACACCGOCA GCAACAA AT T	(Mishing), ith	AAMAC COAC	AMIRAN TI	AGSTATOUSG	480
CTCACGGGTA SOLVETTOTT G	acarrette.	SPEET WACT	$\mathbb{R}^{-1} \times \mathbb{R}^{1} \times \mathbb{R}^{2} \to \mathbb{R}^{2}$	CAACATC66T	404:)
Torrowant of cobactor A	YVG-11621.	Mand Wwat		WWW.Carre	$Q^{(i)}(\cdot)$
ANTOUNANTE POOCOAACA ER	TA MAGA T	::1,2,2,2,2,2	IN THE PROPERTY.	TANA	
A CARLON COMMISSION AND STREET	$h:MV(M) \to$	errani and	zeni melenyzy		
ROW MIA WILAM INTO PARA	W4 12561 /5 +1		A Wale 197.	$\mathbb{E} M^* M^* M^{-1}$	·
PRIMATE WAS ASSESSED.	TWML W	* * * : : 17.7.	7.11 W - 7.4	The grants	* 1
				* ¥'.	
X**					
***************************************	ži. Eleek		11 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	7.7	

TTCAATATCS	JÇA FECIANA	CATHOGAGTA	TT WATORO	omnico (ha	F CUTTO HAAAC	144
TACAACATCG	GATCCGGAAA	CTTCGGGATC	TACAACATCG	GTTTTGGAAA	CGTCGGCGAC	1500
TACAACGTCG	GCTTCGGGAA	CGCGGGCGAC	TTCAACCAAG	GCTTTGCCAA	CACCGGCAAC	1560
AACAACATCG	GGTTCGCCAA	CACCGGCAA.	AACAACATCG	GCATTCGGCT	GTCCGGCGAC	1620
AACCAGCAGO	GGTTCAATAT	TACTAGCGGC	THGAATTCGH	CCVCC&: AV	CAGCGGCCTG	1680
TTQAATTCK	G. ZO WATAA		FF: AA. GCGG		CHICCCCATC	1740
BUAAAATTOBA	DEADORD MA	'',:::::X.'	D. West Conf.	HACCON MA	TA SIGG WTC	18(ii)
TYMATERTG	MAD BRIDGE	ra marrani	may was a second	PODACTECAA	M. M. M. Comple	1860
TA MACACCÇ	AF ATMALL	*, 1.7.1.1.2.2.1.1.1.1.1.1.1.1.1.1.1.1.1.1.	TU WA YIKU E	TEAACA YYAA	VAN NEGOVAC	1000
TTOAA XIT	STGATACIAA	THE MAKET	William CV † .	ERMAN TAA	JA PESSTTE	2.350
TTOMATION	TAA TIT IM	The later	TTO WAY	- DACTIMA	WW. Charles	. 1.4.
TTGGTGCCG	GCGATANGCA	GGGCCAGATT	-ECATCGATS	ColoGoloVo	CATTCCATTC	0100
AFTOTUATAA	AGATAAA	FOTCAL DAY	1.7.27.50.14	TANCOA TIL	ALC: ALC: AAC	1 600
AT AT TACO	$I(CY_{\lambda},;Y;\ldots$	- 17 1 4 <b>V</b> - 2 17 17 1		. P. P. P. TATO I	UMGUNGTTTG	1.220
spinistration of the control of the	and the Tax	TO MANY TO TA	To A contain	$\mathcal{M}_{\mathcal{M}} = \mathcal{M}_{\mathcal{M}}$	TAL MOOTE	
A MADING 14	#. · · !A*· , *#*	7	A: A: Afi		TWW.kv. fr	
t-17.11m11.	monte say		$\cdots \cdots \gamma_{i,j}$	WWW.11.00	A 1944 YM	• ;
	DOTANGE	ATAMATET I	*) W. B.C. C.	MARCON, A.	and a painting	11.4
* * ***						·
*					Ŷ:	
	* .* .					1
				* - 4-		. 10
	. 11 40.				1. 1. E. T.	
8.4						

- (B) TYPE: amino agra-
- J) STRANCEDNESS:
- (D) TOPOLOGY: linear
- (E.F) SEQUENCE DESCRIPTION: JEC 15 NO: 199:
- Gly Gln Asn Ala Fro Ala II. Ala Ala Thi Glu Ala Ala Tyr Asp Gln I
- Met Trp Ala Jim A p Va. Ala Ala Met Ebe elly Tyr Him  $K_{\rm tot}$  Gry Ala 20 25 30
- For Ala Ala Vat For Ala Leo Inc Mo. The Sly (in Ara Leo Fro The  $3^{\rm tr}$
- Val. Also GLY GLY GLY Also bet Val. For Also Also Also a GLE Val. The  $\pm 6$
- The Ari Val Phe Arg Ass. For Gly Let A.a Ass Vi. Arg G.a Cly Ass 65.
- Value And Anto-Try Ann. Value And A. Scholand, Lett. By the Adam algebraic  $\mathfrak{P}_{ij}$
- Thy Arm Thy Aem The Thy New Thy Arm The stry Feb. (4) Ban The stry Feb. (19)
- Then  $A_{ij} \in A_{ij} \cap A_{ij$
- Fire the Even Time low feet Arm Arm (16) the first for the fire the lightest field the fire the first form of the fire the fire that the first fire that the first fire that the first fire the fire the first fire the fire the first fire the fire the first fire t
- ing the 11 per experience of the second seco
- and we have the second of the

- 5.9 Ser lyn Ash Pr  $\times$  Tly Ash ber Ash Thi Gly Sly Phe Ash Met Gly -250 -255
- Gin Tyr Asn Thr Gly Tyr Leu Asn Ser Gly Asn Tyr Asn Thr Gly Leu 260 265 270
- Ala Asn Ser Gly Asn Val Asn Thr Gly Ala Phe the Thr Gly Asn Phe  $\frac{275}{280}$   $\frac{280}{285}$
- Asn Asn Gly Phe Seu Trp Ara Siy Asp His Gln Gly Lea Ile Phe Gly 295 300
- For troubly the the Ash Per II. For Ala Pro Nor Box Gly The Pho 3.7 315 320
- Ann Jes G., Ala Gly ser A. a Ber Gly Phe Lea Art. For Gly Ala Asn. 305.
- Ash Ser sty Phe The Admil'er Ned Ned Ser Sty A. (1): Sty Admiles Gly 44. (1) 150
- Usu Ala Ann Ala Gly Mai heu Mal Ser Sly Ma. The Amn Ser Gly Amn Ser Sel Sel
- The Val Ser Sty Less the Ann Met Ser Less Val Ala The The Pre-
- And like the detaily the the Aon Throfty Ser Ann Het Ser Gly Phe
- The sty of the fact that the Annales of the Alaskan Arquity tap

  to the sty of the Annales of th
- e er er frage fra 197 Ad for amerikan 1970 frage fra fra 1981 av ja 40
- seriesy Am Val. dy Am Tre early a ten dy for a y Am poetagy to

	530					1, 7,					1.1				
Phq 545		110	Ala	303	-11 <i>y</i> 550		A 11:	Short	Πy	11h1 555		Arm	361	Gly	Leu 560
Phe	Asn	Ser	Gly	Thr 565		Asn	Val	Gly	Ile 570		Asn	Al a	Cly	Thr 575	Gly
Aon	Val	G1;	11+- 580	Ala	Acn	guer	σlγ	‴n: 585		Алп	71;	$\Theta^{\frac{1}{2}}Y$	Ile 590		7,STI
i') ~		The		Arn	T!.:		1.454		Astr	Ala		our our	7.72	Ash	Thi
	Elec n.u	Leu	Aan	Alm		Argo	11111	Ann	Thi		Pho 67%		Ash	Th:	Gly
r ,f,	lyn	As ti	71.:				intri	Va.		Ann e e.		Åed.	71.1	-11 y	Agri (-d)
Elies	Ait.			A-9~ 64.7	T**1	Am	: +		ord North		Essi	it		ASI NSS	
Arm	Thr		Pho Cur		Aan	1:-		Ant.	Y:*	Asi.	Thr		A13 623	Phe	Asţ
77.	(11.y	AB AB	Ffare	A Str	žu r		1 t.ε •,Ω;	i×		Иiч		Auje Par	FN-14	( ] . I :	$G_{i}^{*}\gamma$
11.1.				řst j		,* -1 r, -1	1.1.	11.	11.1	1.7.	Em-		1:	He	Assi.
		<u>.</u> į			; t	• •		. i.	X	• • •		; · ; ·	Нy		Auth 170
23-3	i.··	lnt		Thr	' î	• 0 1		tit		. 1.1.	) ·	-11,	. 1(1	11	Per
											-1-1		Y	i ÷	l to

- Also flow Gly Asia Geometry Fiber Glin Asia Gly Ger Lew Glin Ner Gly 835 \$840 \$845
- Trp Ala Asn Leu Gly Asn Ser Val Ser Gly Phe Phe Asn Thr Ser Thr 850 860
- Val Asn Leu Ser Thr Pro Ala Asn Val Ser Gly Leu Asn Asn 11e Gly 865 870 870 875
- The Ash Let Ser ely Val the Arresty Fro Thr Gly Thr Tie The Ash esb 890 895
- Ala Giv ben Ala Ash ben ety En ben Ash He Giy Ser Ala Ser cys age es. 910
- Ary Ho Ard Hir To Lea Ary The Val Jee The He Lee See Ala Phe 915
- Type Gry Ger Alt Ser Asp Gin Ser Ash Fro Sty Ber Val Ber Gin 190 - Ger Ger Ger Ger
- (L. INFORMATION FOR MED ID Notice):
  - 41) SEQUENCE CHARACTERICITECH
    - own throught of happy party
    - (B) TYPE: teacheder acid
    - Sets STRAMPEDMEDD: Sincile
    - "P) TOPOLOGY: linear

will depress a deposit from the first and th

FUNCTOWING PROTESTIVE AND ADDRESS TO THE TAR ADDRESS AND ADDRESS.

(i) BEQUENCE CHARACTERISTICS: (A) LENGTH: 31 base pairs (B) TYPE: nucleic acid (C) STRANGEDNESS: single (D) TOPOLOGY: linear	
(RE) DEQUENCE DESCRIPTION: SP, 110 K.	0:262:
STATE ITO A GUELTSMAAR STATEGAGES I	34
CNOORMATICS FOR THE HESSICS:	
(i) ORDENCE THARACTERISTICS:  (A) LENGTH: It is a party  (P) TYPE: Audible Asid  (C) STRANDEDNESS: Fingle  (I) TOPOLOGY: Linear	
(%) SEQUENCE PENERIPTION: NEW 11 Y.	a. 53:
Morraatto agorotuuaa atottooyoya :	6.
L HEARTATION BORGE, IN KARAME	
A DESCRIPTION OF A PROPERTY OF	
and the state of t	s - 1+
t. Ger Stroppy of the street	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:205:	
CGATATCTGC AGAATTCAGG TTTAAAGCCC ATTTGCGA	38
(2) INFORMATION FOR SEQ 10 NO:200:	
(i) SEQUENCE CHARACTERISTICS:  (A) LIMSTE: We take pairs  (B) TYPE: madely acid  (C) STRARRESERVE area;  (C) COPOLOGY: Timea;	
(xi) seggen a own biblious of the states	
CONTRACT DA OCCACDEGOO MUAACGEOO	30
CO. INFORMATION FUR SET ID NOTES TO	
** JENGER E HABA TEBIGTIÖD:  (Av DENGTE: Si kase palit  (B) TYPE: porter sin:  (C) JERANGENTENT: Jinge  ( D) SUCCEST Linear	
with 19, med house date of the income of the end	

CAGETTON C	OTTA WITTO	MARCE TO	- <b>W</b> 0.2352	CTTTCCTT	TOTTCCTTC	1.2
CTTTCTCGCC	Angringged	GCTTTCCCC	TCAAGCTCTA	AATCGGGGGC	TCCCTTTAGG	18
GTTCCGATTT	AGTGCTTTAC	GGCACCTCGA	CCCCAAAAAA	CTTGATTAGG	GTGATCGTTC	24
ACGTAGTGGG	CCATCGCCCT	GATAGACGGT	TTTTCGCCÇT	TTGACGTTGG	AGTCCACGTT	308
CTTTAATAGT	GGACTOTTGT	COTAAACTOS	AAQAACATTT	AAGCITATOT	CGSTCTATTC	Зυ
PTTTGATTTA	"MAGGGATTT	TG" "GAT" TC	GCTATTGC	TTAAAAAATG	AGCTGATTTA	4276
TTTAAAAATTT	AACGCGAATT	TTAAAJAAAT	WILLW WILLIA	ACARTTURG	GTGGCACTTT	4.80
TAAKCHOCT	CTSCSCGGAA	CCDTMITT:		TAAATACATT	CAAATATGTA	54(
i io mitampo	AATTAATTET	TAMAAAAAT	TATOWNAT	CAAAT JAAA	TGGAATTTAT	600
CATATORS	ATTATCAATA	TTTTATAT	**************************************	PITTINIANT	JAAGGAGAAA	r-to-C
MOTOMOGOMA:	GCAGTTYCAT	AĞGATGG YAA	GATO TOOTA	TOWN, NO.	ATTCCGACTC	720
HOGAACATC	AATACAACCT	ATTAATTEC	TODIÇAAA	AATAAGGITA	TCAACTGAGA	780
WT WICHTE	KII WUWI	SAATCHCE!	A WATER WAY	AMSTETATI	ATTRUTTOO	840
		COATIACET				17.
r'adwid wo	1 WWW.FTG 1	-HTW: NA	14.2 (34.402)	** <i>IMT</i>	ProbaGGW	-t <sub>1</sub> , ·
		11700				1 .
111 7. 1 ° a.	** A CALA	int. Train.	· · · · · · · · · · · · · · · · · · ·			10
		1 257 45 725 1	(14A7AAA). (c)	75 75	STATE AND THE	
	7 7 ···				* # 25	
			o. *		* ** * * * * * * * * * * * * * * * * * *	
	4 N X 1				i	2.55
	()	4.7		* <sub>0</sub> 1 *		

AACT TGTAG	CANCGO TAG	' Alandre "	PATTA TIANTO	· rgtia luko:	. Karran	1740
ASS SECCATA	AGTOGTGTOT	TAJCGRUTTE	. GACTCAAGA	GATASTTAG	` GGATAAGGCC	1800
CAGCGGTCGG	GCTGAACGGG	GGGTTCGTGC	: ACACAGCCCA	. GCTTGGAGCC	AACGACCTAC	1860
ACCGAACTGA	GATACCTACA	GCGTGAGCTA	TGAGAAAGCG	CCACGCTTCC	CGAAGGGAGA	1920
AAGGUGGACA	GGTATCCGGT	ANGCGGGAGG	GTCUGAAÇAG	GAGACCGCAC	HAGGGAGCIT	1980
CHASOSGGAA	ACCCCTGGTA	TETTTATAGE	COLUMNOCO	TTC%*CAVCT	FTGACTTGAG	2040
CITABATETT	ISTGATOTT	STONG EGGE	CONTRACTOR	GGAAAAAACU 1	CAG DAACGCC	7:10¢
CF "CTTTTTJAC	FFTTCCTGGC	ALL LOCATOR	C.TATTC	ACATOTTATA	TOTTSTRATIA	2160
TOTTGATT	CTOTGGATAA	COGIATTACO	\$CCTTTGAGT	GAGCTGAT <b>A</b> C	•CECTOCOCGC	22.20
APPOSAACGA	2018/2001/0	COAGTUADIO	AGCGAGJA/vd	COMMANDA	COT MIGGGE	80
TATTTTCTCC	TASTANT	GTGCCGTATT	TCA-MC-10FA	TATATVIGTO:	ATTOTAGTA	.1340
CAMICISCIC	TGATGCCGUA	TAGITALGCT	AGTATACA	CONCTATOR	TACGINACTO	.:400
a granacer	GNODOCHOMA	M.C.C.C.C.T.W.		tinatic military.	Girmrener	2460
GTT:INTA. W	TOTAL TIALA	W. White it is	117, 2017min 13	5 5 5 5 TO \$ 12.	$\mathbb{A}(\mathbb{R}^n,\mathbb{R}^n) \cup \mathcal{A}(\mathbb{R}^n)$	. 11 t
STITICATO	ADDOAT TADE	AACGCGCHA	A.M. William	TAAACOTUST	CA 1 CONSCIP	
OT WALLDAY	TWWMIGT	***************************************	7.1	AS TOTAL	-: :: 1 ~1 ··· /; -	1.1.4
Asset TO MAKE	10 1 - 11	il service Tribum,	. 35 111	ALCOHOLD TO THE		
PROWEL AT		AND THE PROPERTY.	1 11 11 14 14 1	OF TAAT A	TIS 1 MAST MATE	* .
A YOUNG A SACTO	AD FIDENCE	The State		W. *** *** 11*	1.4 TO 1894 11	.9 . 1
1 & 1 2					* ******	
· www.	AAST SALES	2.1		1 4 4 7	• 100 ex.	
					Y Y	

HAGTTGCAT	I ATAAA-AASW	A CAGTOVIAA	J TORGCOMO	: ATAUTCATO	. George Greek of V	342¢
CCGGAAGGAC	G CTGACTGGG1	TGAAGGCTC	r CAAGGGCATC	· GGTCGAGAT	CCGGTGCCTA	3480
ATGAGTGAGC	TAACTTACAT	TAATTGCGTT	r GCGCTCACTG	CCCGCTTTC	AGTCGGGAAA	3540
CCTGTCGTGC	CAGCTGCATT	AATGAATCGC	CCAACGCGCG	GGGAGAGGC	GTTTGCGTAT	3600
TGGGCGCAG	GGTGGTTITT	CTTTTCACCA	CGTGAGACCG	NATE AND AN	TTGOGGTTCA	3660
concergae :	CTGAGAGAGT	TOPAGNAAG	- durceaccer	GGT PPUCCS Q	AGCAGGCGAA	3720
AA1CUTGTT r	GATOGIG ITT	AAZG KOGGGA	. TATAACATSA	GOTGO TO S	CTATOSTEST	3780
ATCCACTA	CONCAMINA	GPATTAACST	RAGEOGGA	CTCTGTAAT	JCGCGCATT()	3840
postanyeaa.	CATCTOATOR	TT IGGAACSA	-MARIONACT	i naanatti	TOTE ATTOA	.•. 3903
GCATTINGCAT	GCTTWT A	AAAC CWA	131777707	·	1.10 0.11	39ыс
TOROTOMAS	TIMATTICMA	STAMMATATT	TATE OF A RES	A IU WW.	AJACTOGOG	4020
AGAGAGAACT	TAATGJGCCC	CODADAATOD	CCATTICCI	GTGACCCAA1	GCGACCAGAT	4080
acheevee	савтобрата	*CCTCTTÇAT	************	AATAMETET.	ATGGGTGT	4140
SGTCAGAGA"	TAAAF AAITTA	AACGCCGGGAA	TATTAOTOTA	CRACTI	A 'AG''AATGG	420C
CARCORGOT 1	AT TEA POST	TAGITAAT W	2.7. W. 7. W.	W. H. A. L. H. H.	M M MAGAC	4.160
	C:	THE		77. 1 W. C 🛶 *	$\mathcal{W} = \mathcal{M}_{\mathcal{F}} \cap \mathcal{M}_{\mathcal{F}} \cap \mathcal{F}_{\mathcal{F}}$	4.12
	(10 - 10 - 10 - 10 - 10)	1.2.11051.766		AALISHI .T	e van met y	; (4
11.77.17	HW MINNEY	V LEAMVE A	ratingana.	10 Per 45 - 1 - 1 - 1 - 1 - 1 - 1	A MT PTT NO :	€144.
	for the second					
		i		i i		; .
	*		0000		(* -	1.
	v		r #11		· V · · · · · · · · · · · · · · · · · ·	<b>5</b> · · ·
		442 T T T Sec. 1	· . · · ·			: :

AATTAATAGG AGTGAG	TTATA GGGGAATTS	i galeig attaå	CAATTCC: CT	AATAAA )ATO	5040
TTTTGTTTAA CTTTAA	ngaag gagatataca	Y TATOGGÇÇAT	CATCATCAT	ATCAGGTGAT	5100
CGACATCATC GGGACC	CAGCC CCACATCCTC	: GGAACAGGCG	GCGGCGGAGG	CGGTCCAGCG	5160
GGCGCGGGAT AGCGTC	GATG ACATCCGCGI	CGCTCGGGTC	ATTGAGCAGG	ACATGGCCGT	5220
GGACAGCGCC GGCAAC.	ATCA CCTACCGCA	CAAGCTCGAA	CTGTCGTTTA	AGATUAGGCI	* 5286
GGCGCAACUG AGGGGU	TOGA AACCACCGAC	COCTTOCCT	GAAACGGGCG	CCGGCGCCGG	5340
TACTOTOGGG ACTACC	T WE STRUTCHE	GOT GATUTIE	GČGGAGAGY	GTAGCACGC"	5400
GCTCTACCOG CTGTICA	AACC TGPSGGGTCC	GrandlacV.	$\mathrm{GAGAGGTAT}^{-1}$	OAOTOŌAADO	54(.0
CATCACCGCT CARUSCA	And F Transaron	~ 200AT 000 🕏	CACCONGCO	CCHGGACGG"	5520
PAACATTGGG GOOTE Y	SACE COTATOTOTO	1000000000000000000000000000000000000	$V_{\omega}^{2}$ , $i_{i_{1}}$ , $i_{i_{2}}$ , $i_{i_{3}}$	ACAAGGGGCC	$i,j,Q^{\nu q}$
GATGAACAIC GUICTAC	scur fatacquira.	1.77,100,000	TAMAACCIO	CCGGAGTGAC	640
CGAGCACCTC ALGCTOV	AACE GAAAAGTECT	GROWATE	TAP CANGOCA	CCATCAAAAC	5,7(a)
STOGGANDAN TURNASA	ABT CBCOLON.	· · · · · · · · · · · · · · · · · · ·	AA**(TGC:: <sup>‡</sup> \$	CONC. HCGGT	3 / 6/1
AGTTSCOCTS - Williams	mur i Amagamaaa	L MANAGARA	TOSTICAL: O	ACTACCTGTC	187
ltaa koaadati limo ka k	hor in Police AMedic.	11.12.331.3mm	rghayn Artiğ	TOGACTICCC	1881.
	TOWN IN TANKET .	$(\mathcal{V}_{\mathcal{M}_{\mathcal{N}}}, \{\gamma_{1}, \gamma_{1}\}, \gamma_{2})$	M - 17 0, 11 .	Contract of the	÷ (3.)
	CONTRACTOR	· 7:: * * * * * * * * * * * * * * * * * *	way way	"("("A)" ("C), (C) (")	it for a
Ammanna minak		The state of the s	en and de	ATTITIMM"	*********
	**** * ****** ****** *	vivivi i	- I TANK TO SEE	m was filtereday	. 1,
***				* * () * )	
eter a.	ž ' 1	* * * * * * * * * * * * * * * * * * * *	: *	taga ta	
	-				

Greenstand	ACCOMPGNA	( C. M. C. L. C. C. L. C.	TATTECHATA	CTTGTTC. 1 E	, ACCOMPANCE.	1,71 (t
GGCCGCCGCC	AACACGCCGA	. ATGCCCAGCC	GGGCGATCCC	AACGCAGCAC	CTCCGCCGGC	6780
CGACCCGAAC	GCACCGCCCC	CACCTGTCAT	TGCCCCAAAC	GCACCCCAAC	CTGTCCGGAT	6840
CGACAACCCG	GTTGGAGGAT	TCAGCTTCGC	GCTGCCTGCT	GGCTGGGTGG	AGTCTGACGC	6900
CGCCCACTTC	GACTACGGTT	CAGCACTCCT	CAGCAMANCC	ACCGGGGACC	CGCCATTTCC	6960
da ivayêrde	Coloration (Coloration)	O DAATGACAC	**************************************	dTeggeedge	TAGACCAAAA	7020
D.W.T.T.A. DEC	$N303.005\Delta M3$	CACCIMOR	PAAGGCTT ST	accadamps;	COTOGGACAT	7090
TENT MATTE	manara	ANG CHOKAC	· · NATVAA ·	- A BAAAACG	TUTCGCTTGA	7140
THE MARKET	masine and Stands	A. 250 N. WAS THE	227A 1 27A 37C 1	AACTTCA	ATCC FABRAA	2700
DOGANIGGO	Million Committee	CSCGCGTAAT	or think this has	om Magyara	CAPCUGACC	427,5
MCCAPS CAT	PAGCHOTOFF	TTOTCGTATC	arring war	FITAACAAT	CUCTGGACAA	1 4 17
aacageaace	AMIGNOCTICL	CHICATORA	1 1300 NUT DE	(C=\(\);	DOWNSEGO	7.160
0.24.20.60.11	1.TG*AGA6	endomprovide i	w. with that	uninggg <sub>W</sub> y	TOGETHER TANK	74.4+)
ADACDAC 2003	COACACO		W Supra Supra	daar in in	GATAT TOATG	750t.
. William bi	Arma AMe.	MCCACCAC M	0780 78000	· Trace	- TWATDAME	
	7 Y	ert ent book	$\psi_1,\cdots,\psi_V$	TAR TAR AT	with the transfer in the	*** k.; ***
T WW	* * * * * * * * * * * * * * * * * * * *		***************************************	18 - F. 18 - Y. 17 - 1	· · · · · · · · · · · · · · · · · · ·	11 .

AT THE GRAND BUT BUSINESS THE RESERVE

Later than the state of the sta

- Asp der Val Asp Asp Ele Arg Val Ala Arg Val IIe Glu Glu Asp Met 35
- Ala Val Asp Ser Ala Sly Lys He Thr Tyr Ard Ile Lys Leu Glu Val 50 55 60
- Ser Phe Lys Met Arg Pro Ala Gln Pro Arg Cly Ser Lys Pro Pro Ser 65 70 75 80
- Gly Ser Pro Glu The Gly Ali Gly Ala Gly The Val Ala The The Pro
- Ala der Der Fro Mar. Ehn Leid Ala Silu Thr Sily Ser Thr Leu Leu Tyr 100 100 140
- Fro Lem Phe Win Lev. Irp \$17 Fro Ala Phé Hie Glu Ary Tyr Pro Ann 115
- Val The The The Alaskin any The Tay der Tay Alas Jay 1.8 Alas Gir. 130 -130 -130
- Also Also Also Gly The Val. Ass. The Bly Also See Asy. A. r Tyr. Leaf Ber. 145. 150. 255. 166
- Glu Gly Arp Met Ala Ala His .ya Gy les Met Arm lie Ala Deu Ala 188
- The Ser Ala GL, who Val Aan try Arn Louding Gly Val Ser Hough.
- Telling Lemman, "Some byg Wife near Allo Alla Heather and the salvering plant."
- Ty the tension for the second and A + 1 + 1 we give the Values, if
- Both Marie Town The Aras Marie Marie Area (1900) and Area (190
- ing the state of t

				٠. '					5					4.44	
ijijy	Ffre	Ala	340		The	1:0	лl i	A.(n 345)		$\Lambda_{+}^{\circ}$	. Tie	.3++1	Мет 350		- Asp
Gly	Pro	Ala 355		Asp	Gly	Tyr	Pro 360		lle	Asn	Tyr	GIu 365		Ala	lle
Val	Ast. 376	Asn	Arq	Mir.	Түз	Assp Test	Ala	Ala	Thi	Ard	Gln Bro		beti	Gln	Ala
11.0 394.		Hir	î:;	7 1-1	110	mn:	Αψ	$44\gamma$	ht.	1475 394		Ser	Pho	Len	As <sub>k</sub>
Alm	., :	His	Files	1111	114	. •••1	110		Ala 110	Val	Val	Туг	Intu	Ser 415	•
<i>f.::</i> +	Letu		Ži. t		: [			A. i		1-1-1	. 7.=		Aup 430	Ala	$L^{1}$
Ť:::	1.001	Al v 435		V.a	Αlπ		A.4. 442	1100	J.1.	At a		145 445	Gly	Asp	Len
	Thr	Olt.	•	Авр	Od n	Val.: 4 ***	1.	ir r	Thit		91 y 466	701	letti	Gin	Gly
2.1. 41 1	1:10	Ar r	i.ÿ		$A^{(r)}$ :		Th:	i		7. r. 4 <sup>-71</sup>	Alτ	živa	741	Var]	A1 q 480
Eric	1.1-	1,12		ž i Ç	Ástita	i y.	* .:.		112	+ 1.	7 - 47	75 (I		1.200 4.600	i.e.,
÷		. •	ř.: 1	7.1.	Al a	. 1	Ve.	: 11 : -		, v :	At 1	i	7V*4 1		i.
* *.		.:.		i			. 1					1 · · ·	Pt.:	. 1. :	žv.,
						Ð., .					1				8 I
												· ·			
	÷														

11.7 625		Fire	1000		Fr. 630		Sli			Fro 635	Va.	And	Asti	$A_{i}$ , $p$	Thr 640
Arg	He	Val	Leu	Gly	Arq	Leu	Asp	Gln	Lys	Leu	Tyr	Ala	Ser	Ala	Glu

- Arg The Var Beu Gly Arg Leu Asp Gln Lys Leu Tyr Ala Ser Ala Glu 645 650 655
- Ala Thr App Ser Lys Ala Ala Ala Arg Leu Gly Ser App Met Gly Glu 660 670
- The Tyr Mct Fro Tyr Fro Gly Thr Ard Ile Ash Gln Glu Thr Val Ser 675 686
- les App Ali Am. By Mil Ber Sty Ser Alia For Tyr Chu Mal Eyn 602 602 605
- May Cor Pro Ala Air Ach Ala Fr. Ang Ala oly Fr. Fr. Din Ara trp. 725 730 736
- The Val Val Trp Lem oly Thr Ala Arn Ann Pro Val Any Lyn stry Ala -74%
- All bys Ala fee Ala Old De de de Ard fe le L Maj Ala er dre ber Men
- Alger Alger Alger Are there Alger Alger Alger Alger Alger Alger
- Fig. 185 Va. Albert Thronk of Properties for the Archive term  $\frac{1}{2}$

. 1 / /...

## CLAIMS

We claim:

- 1. A polypeptide comprising an antigenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:
  - (a) Asp-Pro-Val-Asp-Ala-Val-Ile-Asn-Thr-Cys-Asn-Tyr-Gly-Gln-Val-Val-Ala-Ala-Leu (SEQ ID NO: 115);
  - (b) Ala-Val-Glu-Ser-Gly-Met-Leu-Ala Leu Gly-Thr-Pro-Ala-Pro-Ser (SEQ ID NO: 116);
  - (c) Ala-Ala-Met-Lys-Pro-Arg-Thr-Gly-Asp-Gly-Pro-Leu-Glu-Ala-Ala-Lys-Glu-Gly-Arg (SEQ ID NO: 17);
  - (d) Tyr-Tyr-Trp-Cys-Pro-Gly-Gln-Pro-Phe-Asp-Pro-Ala-Trp-Gly-Pro (SEQ ID NO: 118);
  - (e) Asp-Ile-Gly-Ser-Glu-Ser-Thr-Glu-Asp-Gln-Gln-Xaa-Ala-Val (SEQ ID NO: 119);
  - (f) Ala-Glu-Glu-Ser-Ile-Ser-Thr-Naa-Glu-Xaa-Ile-Val-Pro (SEQ ID NO: 120);
  - (g) Asp-Pro-Glu-Pro-Ala-Pro-Pro-Val-Pro-Thr-Thr-Ala-Ala-Ser-Pro-Pro-Ser (SEQ ID NO: 121):
  - (h) Ala-Pro-Lys-Thr-Tyr-Xaa-Glu-Glu-Leu-Lys-Gly-Thr-Asp-Thr-Gly (SEQ ID NO: 122);
  - (i) Asp Pro Ala Ser-Ala Pro Asp Val Pro Thi-Ala-Ala-Gln-Leu Thr-Ser i eu-Leu-Asn-Ser-Leu-Ala-Asp-Pro-Asn Val Ser Phe Ala-Asn (SEQ ID NO: 123); and
  - (i) Ma-Pro-Glu-Ser-Gly-Ala-Gly-Leu Gly-Gly-Thr-Val-Gln-Ala-Gly; (SEQ ID NO 131)

wherem Xaa may be any ammo acid

- 2. A polypeptide comprising an immunogenic portion of an *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen has an N-terminal sequence selected from the group consisting of:
  - (a) Asp-Pro-Pro-Asp-Pro-His-Gln-Xaa-Asp-Met-Thr-Lys-Gly-Tyr-Tyr-Pro-Gly-Gly-Arg-Arg-Xaa-Phe; (SEQ ID NO: 124) and
  - (b) Xaa-Tyr-Ile-Ala-Tyr-Xaa-Thr-Thr-Ala-Gly-Ile-Val-Pro-Gly-Lys-Ile-Asn-Val-His-Leu-Val; (SEQ ID NO: 132), wherein Xaa may be any amino acid.
- 3. A polypeptide comprising an antigenic portion of a soluble *M. tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 1, 2, 4-10, 13-25, 52, 94 and 96 or a complement thereof under moderately stringent conditions.
- 4. A polypeptide comprising an antigenic portion of a *M tuberculosis* antigen, or a variant of said antigen that differs only in conservative substitutions and/or modifications, wherein said antigen comprises an amino acid sequence encoded by a DNA sequence selected from the group consisting of the sequences recited in SFQ ID NOS: 26-51, 133, 134, 158-178 and 196, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SFQ ID NOS-26-51, 133, 134, 158-178 and 196 or a complement thereof under moderately stringent conditions.
- $5.00 \times 10 \times V$  molecule comprising a nucleotide sequence encoding a polypeptide according to any one of claims 1-4.

- 6. A recombinant expression vector comprising a DNA molecule according to claim 5.
  - 7. A host cell transformed with an expression vector according to claim 6.
- 8. The host cell of claim 7 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cells.
- 9. A method for detecting M. tuberculosis infection in a biological sample, comprising:
- (a) contacting a biological sample with one or more polypeptides according to any of claims 1-4; and
- (b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting *M. tuberculosis* infection in the biological sample.
- 10. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:
- (a) contacting a biological sample with a polypeptide having an N-terminal sequence selected from the group consisting of sequences provided in SEQ ID NO: 129 and 130; and
- (b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting M tuberculosis intection in the biological sample
- 11. A method for detecting *M tuberculosis* infection in a biological sample, comprising:
- (a) contacting a biological sample with one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 41, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and ONA sequences.

- (b) detecting in the sample the presence of antibodies that bind to at least one of the polypeptides, thereby detecting M. tuberculosis infection in the biological sample.
- 12. The method of any one of claims 9-11 wherein step (a) additionally comprises contacting the biological sample with a 38 kD *M. tuberculosis* antigen and step (b) additionally comprises detecting in the sample the presence of antibodies that bind to the 38 kD *M. tuberculosis* antigen.
- 13. The method of any one of claims 9-11 wherein the polypeptide(s) are bound to a solid support.
- 14. The method of claim 13 wherein the solid support comprises nitrocellulose, latex or a plastic material.
- 15. The method of any one of claims 9-11 wherein the biological sample is selected from the group consisting of whole blood, serum, plasma, saliva, cerebrospinal fluid and urine.
- 16. The method of claim 15 wherein the biological sample is whole blood or serum.
- 17. A method for detecting *M tuberculosis* infection in a biological sample, comprising:
- (a) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotide primers is specific for a DNA molecule according to claim 5, and
- (b) detecting in the sample a DNA sequence that amplifies in the presence of the objectuation thereby detecting M subcreation

- 18. The method of claim 17, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule according to claim 5.
- 19. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:
- (a) contacting the sample with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotide primers is specific for a DNA sequence selected from the group consisting of SEQ ID NOS. 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and
- (b) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers, thereby detecting *M. tuberculosis* infection.
- 20. The method of claim 19, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.
- 21. The method of claims 17 or 19 wherein the biological sample is selected from the group consisting of whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine.
- 22. A method for detecting *M tuberculosis* infection in a biological sample, comprising:
- (a)—contacting the sample with one or more oligonucleotide probes specific for a DNA molecule according to claim 5, and
- (b) detecting in the sample a DNA sequence that hybridizes to the obsonucleotide probe, thereby detecting M tuberculosis infection.

- 23. The method of claim 22 wherein the probe comprises at least about 15 contiguous nucleotides of a DNA molecule according to claim 5.
- 24. A method for detecting *M. tuberculosis* infection in a biological sample, comprising:
- (a) contacting the sample with one or more oligonucleotide probes specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and
  - (b) detecting in the sample a DNA sequence that hybridizes to the oligonucleotide probe, thereby detecting M. tuberculosis infection.
- 25. The method of claim 24 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.
- 26. The method of claims 22 or 24 wherein the biological sample is selected from the group consisting of whole blood, sputum, serum, plasma, saliva, cerebrospinal fluid and urine.
- 27. A method for detecting *M tuberculosis* infection in a biological sample, comprising:
- (a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide according to any one of claims 1-4; and
- (b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M tuberculosis* infection in the biological sample.
- 28. A method for detecting *M nuberculos*, suffection in a biological sample, comprising.

Į,

- (a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide having an N-terminal sequence selected from the group consisting of sequences provided in SEQ ID NO: 129 and 130; and
- (b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M. tuberculosis* infection in the biological sample.
- 29. A method for detecting M tuberculosis infection in a biological sample, comprising:
- (a) contacting the biological sample with a binding agent which is capable of binding to a polypeptide encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and
- (b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting *M. tuberculosis* infection in the biological sample.
- 30. The method of any one of claims 27-29 wherein the binding agent is a monoclonal antibody.
- 31. The method of any one of claims 27-29 wherein the binding agent is a polyclonal antibody.
  - 32 A diagnostic kit comprising:
  - (a) one or more polypeptides according to any of claims 1.4; and
  - (b) a detection reagent.
  - 33 A diagnostic kit comprising:
  - (a) one of more polypeptides having an N terminal seating explicit 4 from

- 34. A diagnostic kit comprising:
- (a) one or more polypeptides encoded by a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198, the complements of said sequences, and DNA sequences that hybridize to a sequence recited in SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198; and
  - (b) a detection reagent.
- 35. The kit of any one of claims 32-34 wherein the polypeptide(\$) are immobilized on a solid support.
- 36. The kit of claim 35 wherein the solid support comprises nitrocellulose, latex or a plastic material.
- 37. The kit of any one of claims 32-34 wherein the detection reagent comprises a reporter group conjugated to a binding agent.
- 38. The kit of claim 37 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G. Protein A and lectins.
- 39. The kit of claim 37 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dve particles
- 40. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a DNA molecule according to claim 5.

- 41. A diagnostic kit according to claim 40, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotide of a DNA molecule according to claim 5.
- 42. A diagnostic kit comprising a at least two oligonucleotide primers, at least one of the primers being specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.
- 43. A diagnostic kit according to claim 42, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotide of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198.
- 44. A diagnostic kit comprising at least one oligonucleotide probe, the oligonucleotide probe being specific for a DNA molecule according to claim 5.
- 45. A kit according to claim 44, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule according to claim 5.
- 46. A diagnostic kit comprising at least one oligonucleotide probe, the oligonucleotide probe being specific for a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151–155, 184–188, 194–195 and 198.
- 47. A kit according to claim 46, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA sequence selected from the group consisting of SEQ ID NOS: 3, 11, 12, 135, 136, 151-155, 184-188, 194-195 and 198
  - 48. A monoclonal antibody that bands to a polypeptide according to any of

- 49. A polyclonal antibody that binds to a polypeptide according to any of claims 1-4.
- 50. A fusion protein comprising two or more polypeptides according to any one of claims 1-4.
- 51. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and ESAT-6 (SEQ ID NO: 99).
- 52. A fusion protein comprising a polypeptide having an N-terminal sequence selected from the group of sequences provided in SEQ ID NOS: 129 and 130.
- 53. A fusion protein comprising one or more polypeptides according to any one of claims 1-4 and the *M. tuberculosis* antigen 38 kD (SEQ ID NO: 150).
  - 54. A diagnostic kit comprising:
  - (a) one or more fusion proteins according to any one of claims 50-53; and
  - (b) a detection reagent.





Fig. 1A-1



110. 1.1-







Fig. 1B-1



F 4



F 1



CPM Incorporated







Fig. 3B







SUBSTITUTE CHEETACHLE 26)





Fig 7



717 (J




Fig. 9



